VOL. 36. Ser. A. Title-page and Index. [Issued December, 1949.]

THE REVIEW OF APPLIED ENTOMOLOGY.

SERIES A: AGRICULTURAL.

ISSUED BY THE COMMONWEALTH INSTITUTE OF ENTOMOLOGY



LONDON:

COMMONWEALTH INSTITUTE OF ENTOMOLOGY,

41, QUEEN'S GATE, S.W.7.

Price 8s. net.

All Rights Reserved.

Commonwealth Agricultural Bureaux.

Executive Council.

Sir PATRICK R. LAIRD, C.B., F.R.S.E., Chairman, United Kingdom.

A. V. HUBBARD, Vice-Chairman, Southern Rhodesia.

Lieutenant-Colonel J. G. ROBERTSON, Canada.

J. E. Cummins, Australia.

E. MARSDEN, C.M.G., C.B.E., M.C., D.Sc., F.R.S., New Zealand.

A. P. VAN DER POST, Union of South Africa.

J. W. DULANTY, C.B., C.B.E., High Commissioner, Eire.

The Deputy High Commissioner, India.

The High Commissioner for Pakistan.

Sir OLIVER GOONETILLEKE, K.C.M.G., K.B.E., High Commissioner, Ceylon.

J. G. HIBBERT, C.M.G., M.C., Colonies, Protectorates and Mandated Territories.

Sir Herbert Howard (Secretary), 2, Queen Anne's Gate Buildings, London, S.W.1.

COMMONWEALTH INSTITUTE OF ENTOMOLOGY.

Director and Editor. W. J. Hall, M.C., D.Sc.

Assistant Director.
T. H. C. Taylor, D.Sc.

Cssistant Editor. H. S. Bushell, M.A.

Head Office—British Museum (Natural History), Cromwell Road, London, S.W.7.

Publication Office and Library-41, Queen's Gate, London, S.W.7.

INDEX OF AUTHORS

A reference in heavy type indicates that a paper by the author has been noticed.

Abdul Ghani, M., 312. Acree jr., F., 25. Afanasiev, M., 158, 388. Afzal Husain, M., 122, 312. Agacino, E. Morales, 205. Aguirre, A. Ego-, 133. Ahlberg, O., 408. Ahmad, Muzaffer, 119, 121. Ahmad, Taskhir, 119, 123. Alexander, C. P., 100. Alfaro, A., 196, 202, 203, 204, 293. Alibert, H., 67, 389. Alkan, B., 3, 352. Allen, N., 371. Allen, N. N., 153. Allen, T. C., 160. Alliot, H., 317. Allman, S. L., 306. Almeida, P. R., 104. Ananda Rau, S., 250. Anderson, L. D., 326, 330. Anderson, R. F., 220. Anderssen, E. E., 45. Ansari, M. Atiqur Rahman, 123. Armitage, H. M., 79, 184, 396. Ashby, D. G., 240, 417. Atiqur Rahman Ansari, M., 123. Austin, M. D., 315. Ayoutantis, A., 292,

Back, E. A., 146. Baker, A. C., 221. Baker, E. W., 148. Balch, R. E., 35, 207. Ball, W. S., 188. Baranyovits, F., 61, 62, 63, 64. Barbier, J., 411. Bare, C. O., 85, 86. Barnes, H. F., 308. Barnes, M. M., 57. Bartlett, K. A., 53. Basinger, A. J., 80. Beacher, J. G., 395. Beckley, V. A., 404. Beckwith, C. S., 137. Bégué, H., 296. Bellod y Bellod, M., 201. Benlloch, M., 199, 204, 293.

Benne, E. J., 134, 135. Bennett, S. H., 132. Beran, F., 292. Besemer, A. F. H., 111, 285, 295. Besson, J., 238. Betheil, J., 153. Bhalla, H. R., 372. Bibby, E. F., 156. Bigger, J. H., 227. Bishop, C. F., 231. Bishopp, F. C., 12. Bissell, T. L., 25. Bjornseth, E. H., 350. Blackburn, N. D., 23. Blake, S. F., 12. Blanch, G. E., 127. Blanchard, E. E., 109. Blanchard, R. A., 157. Blattny, C., 278, 288. Blauvelt, W. E., 96. Blunck, H., 406. Blunt, D. L., 403. Boczkowska, M., 42, 43, 349. Bodenstein, O. F., 374. Bohstedt, G., 153. Bolivar Pieltain, C., 37. Bondy, F. F., 370. Bonnemaison, L., 32, 277. Borg, A., 408. Börner, C., 65, 66. Bottger, G. T., 13, 22. Boudru, M., 353. Bourne, A. I., 159. Bowen, C. V., 13, 93. Bowser, P. H., 350. Box, H. E., 110. Boyce, H. R., 167. Boyd, W. M., 136. Bradley, G. A., 240. Brande, J. van den, 286. Brann jr., J. L., 57. Brannon, L. W., 229. Breakey, E. P., 222. Bredo, H. J., 294. Brett, C. H., 9, 94, 99, 154, 183. Briand, L. J., 77. Brimblecombe, A. R., 68, 248. Brindley, T. A., 61. Broadbent, L., 195, 381.

Brooks, J. W., 326, 330.
Bromley, S. W., 335.
Brouwer, W. M. T. J. de, 239.
Brown, A. W. A., 99, 341.
Brown, N. R., 33, 239, 255.
Brown, W. B., 355.
Brown, W. J., 104.
Bruce, W. G., 93.
Bruce, W. N., 342.
Bruel, W. E. van den, 353.
Brues, C. T., 427.
Brunson, M. H., 152.
Buchanan, L. L., 301.
Burkhardt, G., 325.
Burnet, I. M., 280.

Cagle, L. R., 374. Cakebread, E. J. N., 287. Caldwell, N. E. H., 47, 68, 69, 180, 307, 309. Callan, E. McC., 126, 424. Callens, J. Y., 291. Cañadell Vidal, J. M., 272. Cañizo Gómez, J. del, 198, 199, 385. Cannon, R. C., 69, 249, 308. Capella, A. M. Cid, 168. Carimini, M., 38. Carlson, E. C., 222. Carlson, F. W., 98. Carpenter, S. J., 235. Cassil, C. C., 154. Castagne, E., 308. Castro, A. Ruiz, 198. Cerny, J., 355. Chamberlin, T. R., 157. Chandler, S. C., 95, 127, 322. Chapman, P. J., 271. Chaturvedi, P. L., 122. Chauvin, R., 32. Chevalier, A., 405. Chiaromonte, A., 66, 67. Childers, N. F., 53. Chiu (Shin-foon), 298. Choudhri, R. S., 22. Cid Capella, A. M., 168. Clancy, D. W., 396. Clausen, R., 279. Clausen, R. T., 351. Clemente, F. Gómez, 196, 201. Cobin, M., 54. Collenette, C. L., 26. Common, I. F. B., 362. Compere, H., 32, 97, 109. Conde Jahn, E., 49.

Connell, W. A., 224. Cooley, C. E., 233. Coon, B. F., 18. Cory, E. N., 157. Cotton, R. T., 12, 20. Cox, J. A., 325. Crespy, 280. Crisp, D. J., 272.

Dabrowski, Z., 117, 254. Daguerre, J. B., 50. Daniels, L. B., 79. Darling, H. S., 366. Davatchi, A., 385. Daviault, L., 34, 240, 319, 320. Davis, D. E., 171. Dayal Mathur, Ishwar, 119. De, R. K., 120. Dean, F. P., 19, 98, 149. Dean, R. W., 59, 271, 324. De Bach, P., 145, 160. de Brouwer, W. M. T. J., 239. Decker, G. C., 81, 162, 223, 225, 227, 228. del Cañizo Gómez, J., 198, 199, 385. Deonier, C. C., 168. Dethier, V. G., 298. de Toledo, A. A., 5, 76. Dicker, G. H. L., 314, 359. Dionis'ev, D. E., 118. Ditman, L. P., 95, 325. Dobreanu, E., 41. Dobrovol'skii, B. V., 118. Domenjoz, R., 38, 39. Domínguez, García-Tejero, F., 204. Donnelly, U., 206. Doucette, C. F., 142. Douglas, T. A., 223. Dowden, P. B., 36. Drachovská-Simanová, M., 355. Draize, J. H., 162. Dresden, D., 174. Duffy, E. A. J., 214. Dumbleton, L. J., 309. Dunavan, D., 369. du Pree, M., 25. Duval, G., 8.

Eastwood, H. W., 46. Ebeling, W., 152. Eccleston, K., 39. Eghlidi, S., 387. Ego-Aguirre, A., 133.

Eide, P. M., 221. Elliot, M. I., 88. Emery, G. A., 283. Engel, H., 349. English, L. L., 20, 308. Esfandiari, E., 386. Evans, J. W., 266. Evestropof, E., 387.

Faggioli, D., 194. Fahringer, J., 116, 117. Farrar, M. D., 19, 290. Farwick, S., 406. Fauré, G. Olalquiaga, 108. Favard, A., 269. Fayette, L. J., 154. Fenaroli, A., 252. Fennah, R. G., 71. Fenton, F. A., 158, 388. Feytaud, J., 237. Ficheroulle, H., 42. Finney, D. J., 381. Fisher, E. H., 160. Flanders, S. E., 78. Folsom, D., 209. Forbes, R. S., 240. Forte, P. N., 74. Fox, C. J., 165. Frankenfeld, J. C., 12. Franklin, H. J., 100. Fransen, J. J., 112, 270. Frazier, N. W., 92. Frear, D. E. H., 297. Freiberg, M. A., 142. Frezal, P., 293. Frodsham, J., 378. Fuller, G., 289. Fulmek, L., 270. Fulton, B. B., 148. Fulton, R. A., 160.

Gabotto, L., 355.
Gadd, C. H., 251, 368.
Gaddis, C. H., 336.
Gahan, A. B., 315.
Gaines, J. C., 231.
Gallay, R., 280, 296.
Ganda Ram Sharma, 312.
García-Tejero, F. Domínguez, 204.
Gasser, R., 38.
Gauthier, G., 167.
Gay, F. J., 181.
Gayner, F. C. H., 315.

Geering, Q. A., 261. Gerberg, E. J., 98. Gertler, S. I., 24, 93. Ghani, M. Abdul, 312. Ghesquière, J., 316. Giannotti, O., 300. Gil, G. Piédrola, 60. Gillham, E. M., 380, 381. Girdler, G. W., 168. Glover, L. C., 147. Glover, R. S., 419. Goble, H. W., 163, 166. Goidanich, A., 191. Golfari, L., 191. Gómez, J. del Cañizo, 198, 199, 385. Gomeza Ozámiz, J. M., 215. Gómez Clemente, F., 196, 201. Gómez-Menor Ortega, J., 202. Gonggrijp, J., 276. Goodall, F. L., 323. Gooden, E. L., 340. Goodhue, L. D., 17, 336. Gorton, T. F., 323. Govaerts, J., 375. Graham, C., 157. Grandjean, P. H. Vercammen-, 282. Grandori, L., 346. Grandori, R., 268, 346. Grayson, J. M., 337. Greaves, T., 363. Greenwood, D. E., 183. Greis, 355. Grison, P., 349. Groves, K., 71. Gruber, M., 273. Guido, A. Silveira, 49. Guilhon, J., 237. Gunn, D. L., 205, 375. Günthart, E., 273, 279, 281. Gunther, F. A., 88. Gupta, K. M., 122. Gupta, S. N., 120. Gurcharn Singh Sohi, 123.

Haddock, M. J., 132.
Häflinger, E., 60.
Hallemans, A., 278, 279.
Haller, H. L., 25, 393.
Hamilton, D. W., 58, 334.
Hammond, G. H., 163.
Hansberry, R., 147, 351.
Hargreaves, H., 388.
Harman, S. W., 338.
Haroon Khan, M., 120, 310, 312.

Harris, L. S., 188. Harrison, A., 39. Harrison, P. K., 159. Hartzell, F. Z., 255. Hassan, A. S., 409. Hastings, E., 223. Hathaway, W. B., 159. Haydak, M. H., 343. Hegarty, A., 308. Heinrich, C., 125. Helson, G. A. H., 177, 178. Hely, P. C., 46, 181, 344. Hem Singh Pruthi, 310. Hensill, G. S., 154. Hepting, G. H., 333. Heringa, J. W., 283. Herklots, G. A. C., 3. Hewlett, P. S., 420. Hicks, C. S., 297. Hill, R. E., 236. Hille Ris Lambers, D., 66, 348. Hinton, H. E., 261. Hixson, E., 236. Hodson, A. C., 243. Hodson, W. E. H., 32. Hoerner, J. L., 236. Hoffmann, C. H., 333. Holenstein, R., 273. Holmes, E., 280. Horton, J. R., 250. Hough, W. S., 256, 372. Hu (Ching-yung), 298. Hull, W. B., 153, 229. Hunt, C. R., 329. Hurtig, H., 341. Husain, M. Afzal, 122, 312. Hussein, M., 412. Hutson, R., 134, 135.

Ingle, L., 339. Isely, D., 24. Ishwar Dayal Mathur, 119. Ivy, E. E., 183.

Jackson, M. L., 393.
Jacobson, M., 25.
Jahn, E. Conde, 49.
James, H. C., 302, 304.
Jannone, G., 108, 382, 384.
Jaynes, H. A., 219.
Jenkins, C. F. H., 74, 106.
Jeppson, L. R., 155.
Jepson, W. F., 426.

Johnston, R. N., 33. Johnstone, H. F., 92. Jones, B. M., 28, 208. Jones, M. A., 51. Jones, P. R., 147.

Kamal, M., 410. Kaminski, E., 117. Kapur, A. P., 426. Kapuścinski, S., 267. Kartman, L., 155. Kaussari, M., 387. Kearns, C. W., 92. Kearns, H. G. H., 131. Keenan, W. N., 163. Keifer, H. H., 80, 82, 104. Kelsey, J. M., 414. Kemp, H. K., 246. Kevan, D. K. McE., 106. Khan, M. Haroon, 120, 310, 312. Khristov, A., 354. Kierstead, L. G., 21. Kiriukhin, G., 385. Klarenberg, K. A., 190. Klein, H. Z., 175. Kleine, R., 72. Knowlton, G. F., 11, 24, 388. Komárek, J., 65. Kovache, A., 42. Kovachevski, I. Kh., 4. Krijgsman, B. J., 174. Krogh, P. M. D., 107. Kuenen, D. J., 129. Kurdyumov, N. V., 114. Kurir, A., 104, 115, 116, 117, 128. Kusters, C., 275. Kuwana, S. I., 410.

Ladha Ram Mohindra, 312.
LaDue, J. P., 88, 188.
Lal, K. B., 122.
Lambers, D. Hille Ris, 66, 348.
Lambert, R., 240, 319, 320.
Langford, G. S., 339, 347.
Lardy, H. A., 153.
Lathrop, F. H., 21.
Latta, R., 87, 142.
Lavine, I., 9.
Lazzari, M., 30.
Lea, D. E., 422.
Leach, J. G., 231.
Leclercq, J., 375.
Leech, H. B., 240.

Lefèvre, P. C., 295. Lejeune, R. R., 240. Lepage, H. S., 300. Le Pelley, R. H., 405. Lepesme, P., 177. Lepigre, A. L., 297. Lever, R. J. A. W., 266, 390, 413. Levin, C., 22. Lewis, S. E., 39, 355. Lhoste, J., 105. Lin (Sping), 298. Lindblom, A., 409. Lindgren, D. L., 88, 188. Lindinger, L., 270. List, G. M., 92, 230, 236. Little, E. L., 12. Lloyd, F. E., 96, 97. Loftin, U. C., 401. Lombardini, G., 114. Lord, F. T., 270. Loughnane, J. B., 208. Lounsky, J., 291. Lucas, C. E., 418, 419. Ludvik, G. F., 228. Lupo, V., 383.

Maan, W. J., 270. McCarthy, T., 181. MacCreary, D., 224, 321. McCrory, S. A., 154. McCulloch, E. C., 71. MacDonald, H. A., 71. McDuffie, W. C., 55. McIntosh, A. H., 376. MacKay, M., 240. McKay, R., 208. McKenzie, H. L., 188, 400. Mackie, D. B., 56. McLeod, W. S., 89, 156. Magee, C. J., 365. Maheux, G., 167. Manalo, G. D., 134, 135. Mani, M. S., 110. Manolache, C. I., 41. Manolache, F. C., 40, 41. Marí, A. Quintana y, 168. Marilli, O., 66. Markham, R., 377. Márquez, V. Moreno, 202. Marsh, R. W., 131. Marshall, Sir G. A. K., 388. Martelli, G. M., 43, 72. Martelli, M., 192.

Lyle, C., 219.

Martin, C. H., 15, 18. Martin, Hubert, 131. Martineau, R., 320. Marucci, P. E., 219. Massee, A. M., 66, 358, 359. Massibot, J. A., 250. Mathew, G. E. A., 168. Mathis, W., 83. Mathur, C. B., 122. Mathur, Ishwar Dayal, 119. Mauri, M., 346. May, A. W. S., 248. Meijer, T. M., 351. Melis, A., 31, 38, 64, 141, 191, 193. Melville, A. R., 404. Mendizabal, M., 293. Menor Ortega, J. Gómez-, 202. Michelbacher, A. E., 90, 155. Mickel, C. E., 348. Middlekauff, W. W., 155. Miles, H. W., 173. Miles, M., 173. Miller, E. J., 135. Miller jr., H. W., 258. Miller, L. W., 70, 138, 245. Miner, F. D., 24. Misra, M. P., 120. Mitchell, B. L., 137. Mohindra, Ladha Ram, 312. Monte, O., 6. Morales Agacino, E., 205. Morant, V., 169. Moreno Márquez, V., 202. Moretti, G., 346. Morris, K. R. S., 255. Morrison, A. E., 190. Morrison, F. O., 164, 421. Morstatt, H., 72. Moursi, A. A., 410, 427. Moutia, L. A., 125, 209. Muesebeck, C. F. W., 32. Müller, P., 274. Muma, M. H., 157. Munro, H. K., 176. Munro, J. A., 128, 388. Munro, J. W., 32. Muzaffer Ahmad, 119, 121.

Nagel, R. H., 34. Neal, P. A., 161, 162. Neary, M. E., 392. Negi, P. S., 120. Nel, R. G., 168. Nelson, H. D., 160. Newcomer, E. J., 19, 98, 149. Newton, H. C. F., 377. Newton, J. H., 92. Nickels, C. B., 151. Nobrega, N. R., 7. Noirot, C., 317. Nolte, H. W., 75. Norton, L. B., 329, 351. Nye, W. P., 24.

Oettingen, W. F. von, 161.
O'Kane, W. C., 233.
Olalquiaga Fauré, G., 108.
Oldham, H. T., 343.
Oosthuizen, M. J., 1.
Orelli, O. Schneider-, 407.
Orlando, A., 7, 299, 300.
Ortega, J. Gómez-Menor, 202.
Osborn, H. T., 146.
Osburn, M. R., 83.
Ossiannilsson, F., 408.
Otten, E., 239.
Owen jr., W. L., 231.
Oxley, T. A., 265, 405, 418.
Ozámiz, J. M. Gomeza, 215.

Packard, C. M., 96. Pagán, C., 51. Palmer, D. F., 190. Paquet, G., 319. Parker, D. E., 143. Parker, W. L., 224, 395. Parks, T. H., 136. Parr, W. J., 360. Patterson, N. A., 270. Paulian, R., 177. Pearson, A. G., 15. Peay, W. E., 10, 11, 12. Pelley, R. H. Le, 405. Pemberton, C. E., 249. Pepper, J. H., 223. Perron, J. P., 165. Pestel, E., 293. Peterson, A., 9, 258. Peterson, L. O. T., 240. Pettey, F. W., 171, 389. Philips, F. S., 162. Phillips, G. R., 146. Pickett, A. D., 270, 392. Pickles, A., 318, 319. Piédrola Gil, G., 60. Pieltain, C. Bolivar, 37. Pierce, W. C., 151.

Pinhey, E. C. G., 74. Planes, S., 197. Plank, H. K., 51, 53, 54, 55. Plank, J. E. van der, 45. Plummer, C. C., 148. Pogorelskin, M. A., 393. Pol, P. H. van de, 111. Poos, F. W., 337. Posnette, A. F., 110. Post, R. L., 128. Potter, C., 377, 380, 381, 420. Potts, S. F., 36, 145. Powning, R. F., 364. Pradhan, S., 310. Pree, M. du, 25. Prentice, I. W., 376. Provasoli, L., 348. Proverbs, M. D., 421. Pruthi, Hem Singh, 310. Putnam, L. G., 99.

Quayle, H. J., 80. Questel, D. D., 22. Quidet, P., 405. Quintana y Marí, A., 168.

Radford, C. D., 427. Rahman Ansari, M. Atiqur, 123. Rainey, R. C., 27. Rainwater, C. F., 370. Ram Mohindra, Ladha, 312. Ram Sharma, Ganda, 312. Rao, A. S., 110. Rau, S. Ananda, 250. Raucourt, M., 296. Reed, W. D., 86. Reeks, W. A., 207, 240. Rehn, J. A. G., 301. Reid, J. A., 418. Reinhard, H. J., 14. Rhoades, W. C., 94, 99. Rice, E. B., 363. Rice, P. L., 257, 321. Rice, R. I., 92. Richards, O. W., 210, 213, 265. Richardson, C. H., 85. Riker, A. J., 220. Ripper, W. E., 260. Risbec, J., 44, 250. Ris Lambers, D. Hille, 66, 348. Rivals, P., 317. Rivnay, E., 409. Roark, R. C., 208, 240.

Roberti, D., 193.
Roberts, F. M., 423.
Roberts, R. S., 24.
Robertson, P. L., 416.
Robles, P. Seguinot, 53, 54.
Rodionov, Z. S., 3.
Roehrich, R., 349.
Romney, V. E., 94, 96.
Rossem, G. van, 130.
Roth, E. R., 333.
Rothschild, Lord, 26.
Rubtzov, I. A., 241.
Ruiz Castro, A., 198.
Russo, G., 29, 195.
Ruszkowski, J., 117, 384, 385.
Ryan, H. J., 189.

St. John, J. L., 71. Salt, R. W., 390. Satchell, J. E., 377. Satterthwait, A. F., 83, 150, 157. Sauer, H. F. G., 76. Sauvage, F., 278. Sayed, M. Taher, 409, 411, 412. Scala, A., 238. Scaramuzza, L. C., 240. Schechter, M. S., 393. Schedl, K. E., 31, 32, 348. Schmidt, G., 72. Schneider, F., 273. Schneider-Orelli, O., 407. Schwan, B., 409. Schwitzgebel, R. B., 12. Scott, E. E., 401. Seguinot Robles, P., 53, 54. Sen, A. C., 122. Servadei, A., 30, 37, 115. Shanahan, G. J., 365. Sharma, Ganda Ram, 312. Sharp, S. S., 96. Shaw, F. R., 136, 159. Shaw, J. G., 20, 221. Shaw, M. W., 377. Sherman, F., 370. Shirck, F. H., 91. Short, J. R. T., 69. Siegler, E. H., 24, 93, 343. Silberschmidt, K., 7, 299. Silva, P., 125. Silveira Guido, A., 49. Silvestri, F., 176. Simanová, M. Drachovská-, 355. Simmonds, F. J., 28, 425. Simonetta, E., 269.

Simpson, L. J., 35. Sinclair, D. A., 297. Singh Pruthi, Hem, 310. Singh Sohi, Gurcharn, 123. Small, T., 259. Smee, C., 44. Smith, C. C., 35, 240. Smith, F. F., 325. Smith, H. S., 80. Smith, J. H., 307. Smith, K. M., 312, 357, 377, 422. Smith, R. F., 90. Smith, R. W., 166. Smith, W. A., 306. Snapp, O. I., 95, 235. Snyder, T. E., 317. Soenen, A., 277. Sohi, Gurcharn Singh, 123. Solomon, M. E., 417. Sonan, J., 177, 208. Sorenson, C. J., 12. Soyka, W., 64. Speyer, E. R., 360. Squires, D. W., 339. Srivastava, A. S., 153, 229. Stafford, E. M., 16, 342. Stahl, C. F., 87. Stahler, N., 23. Standish, J., 348. Starr, D. F., 20. Stearns, L. A., 224, 257. Steinweden, J. B., 79. Stephens, R. M., 179. Stevenson, F. J., 209. Stewart, K. E., 34. Stewart, W. W. A., 166. Strawińska, J., 354. Strawiński, K., 254. Strickland, A. H., 110. Struble, G. R., 301. Stultz, H. T., 270. Sukhov, K. S., 75. Summersgill, J. V., 323. Sun (Yun-pei), 420. Suranyi, P., 254. Swain, R. B., 83. Swanson, C., 155. Sweetman, H. L., 378.

Taher Sayed, M., 409, 411, 412. Taskhir Ahmad, 119, 123. Taylor, H., 378. Taylor, T. H. C., 26. Tejero, F. Domínguez García-, 204. Telford, H. S., 379.
Tenhet, J. N., 85, 86.
Terpstra, P., 112, 270.
Theobald, F. V., 66.
Thomas, H. E., 401.
Thompson, R. W., 163, 164, 166.
Thompson, W. R., 348.
Thorne, F. T., 190, 343.
Thorpe, W. H., 272.
Tilemans, E., 292.
Todd, F. E., 10.
Toledo, A. A. de, 5, 76.
Tooke, F. G. C., 107.
Tudor, P., 260.
Turner, N., 328, 401.
Turnipseed, G. F., 308.
Twinn, C. R., 167.

Ullyett, G. C., 172. Underhill, G. W., 374, 394. Urquijo, P., 199, 200.

Van den Brande, J., 286. van den Bruel, W. E., 353. van de Pol, P. H., 111. van der Plank, J. E., 45. van Rossem, G., 130. Vansell, G. H., 10. Vasil'ev, I. V., 241, 242. Vayssière, P., 316. Venkataraman, T. V., 120. Venturi, F., 30, 37, 114. Vercammen-Grandjean, P. H., 282. Verma, P. M., 120. Verma, R. M., 122. Vidal, J. M. Cañadell, 272. von Oettingen, W. F., 161. Voûte, A. D., 352. Vovk, A. M., 75. Vrydagh, J. M., 1, 295. Vukasovič, P., 251.

Wagner, R. P., 15. Walkden, H. H., 12, 20. Wallace, C. R., 73, 138. Wallis, R. L., 144. Waloff, N., 210, 213, 264. Waloff, Z., 123, 205. Walton, R. R., 183. Wason, E. J., 247. Waterston, J. M., 48. Watkins, T. C., 81, 329. Watson, E. B., 240. Watson, M. A., 423. Watson, T. Y., 403. Webber, H. H., 15. Webster, R. L., 222. Weinman, C. J., 223, 225, 227, 243. West, T. F., 277. Westenberg, L., 112, 270, 295. Westlake, W. E., 19. Wheeler, E. H., 56. White, D. G., 54. Whitehead, T., 312, 376. Whitten, R. R., 143. Wickens, G. M., 422. Wiesmann, R., 353. Wilkes, A., 78. Wille, J. E., 133. Wilson, F., 181, 182, 364, 365. Wilson, H. F., 153, 229, 393. Wilson, J. K., 22. Winteringham, F. P. W., 39. Winters, H. F., 53. Wipprecht, R., 231. Wisecup, C. B., 168. Wishart, G., 77, 391. Wolcott, G. N., 169, 232. Wolfenbarger, D. O., 8, 17. Wood, C. A., 312, 376. Wood, S. L., 24. Woodruff, N., 328. Woodside, A. M., 334, 373. Wright, D. W., 240, 261, 417. Wright, J. M., 19.

Yetter jr., W. P., 341. Yuill, J. S., 168. Yust, H. R., 160.

Zacher, F., 130. Zimmerman, O. T., 9. Zinkernagel, R., 38. Zobrist, L., 282.

GENERAL INDEX

In the case of scientific names, the page reference is cited only under the heading of the generic name.

Abbella, Gir. nec v. Heyd. (see Abbellisca). Abbellisca, n.n. for Abbella, Gir., 316. abbreviata, Aphis (see A. rhamni).

Abies balsamea (Balsam Fir), pests of, in Canada, 33, 34, 35, 78, 208, 240, 255, 319; type of, favourable to Harmologa fumiferana, 207.

abietina, Pristiphora (Lygaeonematus).

abietis, Cephaleia (Lyda).

abstrusus, Pseudaphycus. Abutilon, Macrosiphum solanifolii on, in greenhouses in Britain, 361.

Abutilon avicennae, Podagrica spp. on, in Rumania, 41.

Abutilon indicum, Earias insulana on, in Punjab, 312.

Abutilon striatum var. spurium, Bemisia tabaci transmitting virus to, in Brazil,

Abyssinia, Schistocerca gregaria in, 124, 403.

Acacia (see Robinia).

Acaena, use of Antholcus varinervis against, in New Zealand, 244.

Acalypha, Ferrisia virgata on, in Gold Coast, 111.

Acantholyda pinivora, identification of excreta of, in Germany, 76.

Acanthoscelides (see Bruchus).

Acaphylla steinwedeni, on Camellia in California. 187.

Acer (see Maple).

Acer glabrum, new Aphid on, in Utah,

Acer negundo, Hyphantria cunea on, in Hungary, 254.

acerata, Acraea.

Aceria camelliae, on Camellia in California, 187.

Aceria essigi, on blackberry in U.S.A., 83. Aceria mangiferae, sp. n., on mango in Egypt, 409.

Aceria sheldoni (on Citrus), in Australia, 181; in U.S.A., 83, 181, 188; measures against, 181.

Aceria tulipae, in Liliaceous bulbs in

U.S.A., 83.

aceris, Trioxys (see T. cirsii).

Acetic Acid, survival of Tyroglyphus nadinus in, 114; Celcure (q.v.) containing, 414.

4-Acetodiphenyltrichlorethane, 275.

Acetone, uses of, as a solvent, 9, 51, 90, 95, 131, 135, 154, 164, 224, 226, 230, 274, 325, 341, 342; in aerosol formula. 325.

Achaetoneura samiae, developing in Loxostege sticticalis after Meteorus loxostegei, 29.

Achras zapota, Rhodoneura myrsusalis on, in Ivory Coast, 67.

Achrysocharella ruforum, parasite of Diprion pini in Germany, 239.

Achrysopophagus aegyptiacus, hosts of, in Palestine, 410.

Acraea acerata, on sweet potato in Uganda, 367.

Acraea alciope, bionomics of, Boehmeria nivea in Ivory Coast, 67.

Acridiophaga (see Sarcophaga).

Acritocera negligens, on coconut in Fiji, 267.

Acrobasis caryae, sprays controlling, on pecan in Texas, 152.

Acrosternum hilare, food-plants and control of, causing deformation of peaches in Virginia, 373.

Acrotylus, on Eleusine in Uganda, 367. Acrylonitrile, tests and uses of, as a fumigant, 12, 20, 62; and carbon tetrachloride, 12.

Actia interrupta, hosts of, in Quebec, 321.

acuminatus, Coccus.

acupunctatus, Scyphophorus, Acutaspis, characters of, 401.

Acutaspis subnigra, sp. n., on avocado in Peru, 401.

Acutaspis tingi, sp. n., intercepted in California on coconut from Mexico.

Acyrthosiphon (see Macrosiphum).

Adelencyrtus simmondsi, sp. n., parasite of Coccus hesperidum in S. Australia, 32.

Adelges (see Chermes).

Adelphocoris rapidus, on cotton in U.S.A., 156, 184.

Adenostoma, Tachardiella ferrisi on, in California, 187.

Adhesives, banding with, 345, 404; other uses of, for catching insects, 123, 195.

Adiantum cuneatum, insecticides injuring,

adonidis, Entomoscelis. adonidum, Pseudococcus. Adoretus, soil treatments against, in tobacco fields in S. Rhodesia, 137.

Adoretus tenuimaculatus, A. versutus recorded as, in Fiji, 266.

Adoretus versutus, food-plants of, in Fiji, **266**, **390**; DDT against, **266**.

adornatus, Calacarus.

Aëdes concolor, insecticide tests on larvae of, 298.

Aegeria bibionipennis, on raspberry and Boysenberry in California, 399.

Aegeria exitiosa, treatments against, on peach in Delaware, 321.

aegopodii, Cavariella.

aegyptiacus, Achrysopophagus.

aemula, Camptylochila (Epizeuxis).

aeneovirens, Rhynchites (see R. minutus). aenescens, Naranga.

aeneus, Chalcodermus; Corymbites (Selatosomus).

aequatus, Rhynchites. aerea, Phyllotreta.

Aerosols, uses of DDT as, 18, 87, 88, 92, 95, 147, 161, 162, 230, 292, 325, 326, 327, 331, 336, 337; other insecticides used in, 147, 228, 230, 236, 331, 336, 337; equipment for dispersing, 17, 87, 88, 92, 95, 147, 325; dispersed from aeroplanes, 292, 293; against pests in aircraft, 233, 336, 337; factors affecting efficiency of, 17, 18.

Aeschynomene sensitiva, tests of toxicity

of, to insects, 52.

Aesculus californica, Eulecanium kunoense on, in California, 399.

aethiops, Caliroa.

Affinin, source and structure of, **25**. *affinis*, *Agria* (*Pseudosarcophaga*). *afra*, *Spalangia*.

Africa, French Equatorial (see Cameroons).

Africa, French West, pests of stored and field ground-nuts in, 44, 250; Lepidoptera on guava in, 67, 389, 390; miscellaneous pests in, 67, 68, 250; breeding and migrations of Schistocerca gregaria in, 205, 206, 207; new form of Tribolium indicum in, 261; beneficial insects in, 44, 67, 390.

Africa, Portuguese East, *Dacus bivittatus* cucumarius in, 176; digest of plant-quarantine restrictions in, 105.

Africa, South, Aonidiella aurantii on Citrus in, 172; Dysdercus spp. on cotton in, 27; Dacus bivittatus cucumarius in, 176; Desmeocraera varia on Mimusops spp. in, 389; new weevil on sweet potato in, 388; thrips and virus disease of tobacco in, 45; Coleoptera in timber in, 1, 107; pests

of stored products in, 1, 2, 168; factors affecting control of prickly pear by insects in, 171, 172, 389; Dactylopius opuntiae attacking spineless cactus in, 389; natural enemies and biological control of insects in, 1, 27, 389; plant quarantines in, 105, 209; attempted introduction of Anaphoidea nitens into Mauritius from, 125; Cydia pomonella intercepted in Kenya in apples from, 405.

africana, Gryllotalpa.

africanus, Coccus (Lecanium); Lyctus.
Agallia albidula, food-plants of, transmitting virus of tomato in Brazil, 77.
Agasphaerops nigra, bionomics and con-

trol of, on Lilium longiflorum in U.S.A.,

Agathis bicolor, parasite of Harmologa fumiferana in Quebec, 321.

Agathis buttricki, hosts of, in U.S.A., 84. Agathis pumila, utilisation of, against Coleophora laricella in New Brunswick, 208.

Agave americana, 389. agavis, Chrysomphalus.

Agicide, 395.

Agonoscelis pubescens (versicolor), parasite of, 27.

Agonoxena argaula, on coconut in Fiji, 390.

Agral W.E.M., 28.

Agria affinis, utilisation of, against Harmologa fumiferana in Quebec, 78. agricola, Anisoplia.

Agrilus, on Pistacia in Persia, 386.

Agrilus anxius, relation of, to die-back of birch in Canada, 208, 319, 320.

Agrilus hyperici, establishment of, against Hypericum perforatum in Australia, 216; imported into California, 216.

Agriotes, in Belgium, 287; in Britain, 281, 381, 382; effect of populations of, on crop yields, 281, 381, 382; effect of benzene hexachloride against, 281; susceptibility of, to stomach poisons, 287.

Agriotes lineatus, in Hungary, 64.

Agriotes obscurus, in Hungary, 64. Agriotes sputator, in Hungary, 64.

Agriotes ustulatus, in Hungary, 64.

Agrocide 3, 303.

Agromyza phaseoli (on beans), in Australia, 46, 47, 73, 177; in Mauritius, 210; insecticides against, 46, 47, 73, 210.

Agromyza theae, bionomics of, on tea in Ceylon, 368.

Agropyrum spp., Cephus cinctus on, in N. Dakota, 128.

Agrotis segetum, possibly transmitting tobacco mosaic in Crimea, 75; bait for, damaging castor in Hungary, 62.

Agrotis ypsilon, on cranberry in Massachusetts, 100; outbreak of, in Queensland, 307; in Virginia, 333; measures and experiments against, 307, 308, 333.

Aiolopus, in Mauritius, 210.

Aircraft, carriage and control of insect pests in, 106, 233, 249, 336, 337; insecticides distributed from, 33, 34, 35, 36, 90, 99, 143, 228, 236, 237, 292, 294, 327, 328, 396, 401; equipment for applying insecticides from, 33, 34, 92; factors concerned in effectiveness of insecticides distributed from, 17, 33, 35. 146. (See Helicopters.)

abama, Anticarsia gemmatalis on ground-nuts in, 20; Fiorinia theae Alabama, on Camellia in, 188; quarantine against white-fringed beetles in, 184. Albania, Ceresa bubalus on fruit trees in.

Alberta, Meteorus loxostegei in. 28: Melanoplus mexicanus in, 99.

albidula, Agallia. albifrons, Decticus.

albitarsus, Xorides.

Albizzia, Tydeus womersleyi on, in Ceylon, 369; pests of, in U.S.A., 143.

Albizzia lebbek, Anychus orientalis on, in Egypt, **412**.

albizziae, Homadaula. albolineana, Taniva.

Alcidodes dentipes, on ground-nuts in Tanganyika, 426.

alciope, Acraea.

Alcohol, survival of Tyroglyphus nadinus in, 114; uses of, as a solvent, 24, 93, 94, 135, 140, 164. (See Methylated Spirit.)

alecto, Telenomus (Prophanurus).

parasite Aleochara bipustulata, of Hylemyia spp. in Britain, 174.

Aleochara sparsa, parasite of Psila rosae in Britain, 417.

Aleurites, pests of, in Nyasaland, 44. Aleurodes citri (see Dialeurodes).

Aleurodids (see Whiteflies).

Algeria, Quadraspidiotus perniciosus in, 293; breeding and migrations of Schistocerca gregaria in, 206, 207.

Alizarin, test of, against Kalotermes brevis, 233.

Allotropa burrelli, parasitising Pseudococcus comstocki in Japan and U.S.A., 410; not parasitising mealybugs known as P. comstocki in Palestine,

Alloxysta, hyperparasite of Toxoptera graminum in Uruguay, 50.

Allyl Pentachlorphenyl Ether, 224. Almond, pests of, in U.S.A., 152, 156,

185, 398, 399.

Alocasia, Papuana huebneri on, in Ellice Is., 390.

Aloe, American, 389.

Alophora nasalis (parasite of Dysdercus spp.), bionomics and distribution of, in Africa, 27; synonymy of, 27.

Alophora nigeriensis (see A. nasalis).

Alophora subcoleoptrata, bionomics of, parasitising Eurygaster integriceps in Tadzhikistan, 241.

Alsophila pometaria, on shade trees in New Brunswick, 35, 208; in U.S.A., 36, 143; insecticides against, 35, 36,

Alstonia congensis, new Coccid on, in Gold Coast, 110.

Alternanthera sessilis, Marasmia venilialis on, in Fiji, 267.

Althaea officinalis, Podagrica spp. on, in Rumania, 41.

Althaea rosea, Earias spp. on, in Punjab. 312; Podagrica spp. on, in Rumania, 41.

Altica (see Haltica).

altitudinis, Chelonus.

Alumina (see Aluminium Oxide).

Aluminium, suitability of, for containers for Toxaphene, 395.

Aluminium Arsenate, in sprays and dusts against Prays oleellus on olive, 141.

Aluminium Hydroxide Gel, decreasing efficiency of lead-arsenate sprays against Cydia pomonella, 372.

Aluminium Oxide, dust of, against Calandra oryzae, 217; as carrier for dust, 218; types of, 217, 218.

Aluminium Stearate, effect of, in DDT

sprays, 88, 89.

Aluminium Sulphate, in spray formulae. 98, 149.

alutacea, Physonota.

alveolatifrons, Pseudaphycus.

Alydus, on Lespedeza in Virginia, 374.

amabilis, Eublemma.

Amarantus retroflexus, Lygus oblineatus on, in Virginia, 334.

ambiguella, Clysiana (Clysia).

Amblymerus subfumatus, parasite and hyperparasite of Diprion pini Germany, 239.

Amblymerus verditer, hosts of, in Quebec, 320, 321.

Amblypelta cocophaga, on cassava in Solomon Is., 390.

Ambrosia, Psallus seriatus reared on, in Oklahoma, 184.

Ambrosia artemisifolia, Lygus oblineatus on, in Virginia, 334.

ambulans, Onychiurus.

Ameloctonus christenseni, sp. n., parasite of Thyreion gelotopoeon in Argentina, 109.

americana, Malacosoma; Periplaneta.

americanum, Mezium.

americanus, Pigmeophorus.

amictorius, Exenterus.

amicus, Eupelmus cyaniceps.

Ammonia, as fumigant against cheese mites, 245; preventing fermentation of sunflower-seed oil emulsion, 354.

Ammonium Dinitro-o-cresylate, 111.

Ammonium Phosphate, experiments with, as bait for *Dacus oleae*, **201**.

Amorpha, toxic principle of Amorpha

fruticosa, 9.

Amorpha fruticosa, pests of, in Oklahoma,
9; tests of extracts of, against insects,
9, 10, 154; toxicity of, to mammals,
10; toxic principle of, 9.

amorphella, Walshia.

Amradiplosis echinogalliperda, gen. et sp. n., on mango in India, 110.

Amyl Alcohol, thiocyanoacetic ester of, 252, 253, 254. (See Thiocyanoacetates.)

Amyl Pentachlorphenyl Ethers, 224.

Amyris elemifera, timber of, repellent to termites, 169.

Anabasine, tests of toxicity of, to insects and *Tetranychus*, 13, 93.

Anabasine Sulphate, spraying with, ineffective against *Polychrosis botrana*, **4.** Anacardic Acid, repellency of chlorinated,

to Kalotermes brevis, 232.

Anacardium occidentale, new Mirid on, in Uganda, 26.

Anachaetopsis tortricis, parasite of Homoeosoma electellum in U.S.A., 84.

Anagrus, parasitising Apanteles glomeratus in Germany, 407.

Anagrus armatus var. nigriventris, use of, against Typhlocyba froggatti in Tasmania, 70.

Anagrus epos, DDT apparently not harmful to, in U.S.A., 147.

Anagyrus, hyperparasite affecting value of, against *Pseudococcus kenyae* in Kenya, 404.

Anagyrus kivuensis, 404; parasite of mealybugs in Palestine, 410; parasite of, 410.

analis, Bruchus.

Anantha lateralis (see Helomyia).

Anaphoidea nitens, establishment of, against Gonipterus scutellatus in Kenya and Mauritius, 106, 125, 405; lifecycle and rearing of, 106.

Anaphothrips orchidii, insecticides against, on Cinchona in Porto Rico, 53.

Anasa tristis (Squash Bug), insecticides against, in U.S.A., 12, 101, 234, 370.

anastativorus, Torymus.

Anastatus disparis (bifasciatus, auct.), bionomics of, in Jugoslavia, 116, 117. Anastrepha, tests of insecticides against, in Peru, 134.

Anastrepha ludens, on mangos and Citrus in Mexico, 20, 148; effects of sub-lethal doses of tartar emetic on, 148;

repelled by smoke, 20.

Anastrepha mombinpraeoptans, spray against, on mango in Porto Rico, **54.**Anastrepha serpentina, host-fruits and distribution of, in Mexico and Texas, **221.**

Ancylis comptana, 178.

ancylivorus, Macrocentrus. ancylus, Diaspidiotus.

andreae, Dysdercus.

Andrector ruficornis, tests of insecticides on, in Porto Rico, 52.

Andromedotoxin, in flowers of *Rhododen-dron molle*, **299**.

Andropogon, Phenacoccus on, in Maryland, 315.

Angitia, attempted introduction of, into Cuba against Homoeosoma electellum, 84.

Angitia cacoeciae, parasite of Harmologa fumiferana in Canada, 207.

Angitia cerophaga, introduced parasite of Plutella maculipennis in New Zealand, 244.

Angitia chrysosticta, parasite of Ephestia elutella in Britain, 212.

Angitia obliterata, technique of rearing, from Cydia molesta in peach terminals in Ohio, 23.

angusi, Datana.

angustatus, Hylastes.

angustifrons, Pseudaphycus.

Anilastus, parasite of Athalia rosae in Italy, 193.

Anisandrus (see Xyleborus).

Anisoplia spp., bionomics and control of, on grasses and wheat in Persia, 387.

Anisota rubicunda, spray against, on shade trees in U.S.A., 143.

Anisota virginiensis, spray against, on shade trees in U.S.A., 143.

Annona cherimolia, 134.

annularis, Chlorophorus.

annulata, Myzocallis.

annulatus, Phytodietus.

Anobium punctatum, tests with timber preservatives against, in S. Africa and New Zealand, 107, 245, 414-416.

Anomala, soil treatments against, in tobacco fields in S. Rhodesia, 137.

Anomis texana, dusts ineffective against. on cotton in Peru, 134.

Anomocaulus fulvovestitus, on pineapple and Pandanus spp. in Fiji, 266, 390.

Antestia (on coffee), in Belgian Congo, 295; in Kenya, 404; parasite of, in Uganda, 367; dusts against, 295, 404.

Antestia faceta, on coffee in Uganda, 367. Antestia lineaticollis (on coffee), in Fr. Cameroons, 295; in Uganda, 367.

Anthemis tinctoria kelwayi, Homoeosoma electellum on, in Iowa, 84.

Antheraea pernyi (Tussore Silk Worm), birds preventing establishment of, in Italy, 67.

Anthocoris nemorum, predacious on Paratetranychus pilosus in England, 288; dinitro compound not harmful to, 288.

Antholcus varinervis, use of, against Acaena in New Zealand, 244; natural enemies of, 244.

Anthonomus grandis, on cotton in U.S.A., 13, 14, 231, 232, 370, 371, 401; not found in California, 398; bionomics of, 371; measures and experiments against, 13, 14, 231, 232, 370, 371,

401; history of, 401.

Anthonomus musculus, sprays against, on blueberry in New Jersey, 137.

Anthonomus piri (on pear), bionomics of, in France, 277; in Holland, 286; insecticides against, 277, 286.

Anthonomus pomorum (on apple), in Britain, 284, 314, 315, 359; in France, 293; in Holland, 276, 285, 286; in Spain, 198; in Switzerland, 296; bionomics of, 314; insecticides against, 198, 276, 284, 286, 293, 296, 315, 359.

Anthonomus rubi, on strawberry, etc., in Belgium, 278, 279, 353; bionomics

and control of, 278, 279.

Anthonomus vestitus, dusts ineffective against, on cotton in Peru, 134.

Anthrenus vorax, not attacking fleeces from goats dipped in DDT, 146; Toxaphene protecting cloth from, 395. Anticarsia gemmatalis, dusts against, on

groundnuts in Alabama, 20.

Antimony Fluoride, repellency of wood treated with, to Kalotermes brevis, 232.

antiopa, Nymphalis.

66736

antiqua, Hylemyia (Chortophila); Orgyia. Antirrhinum, selenium treatment of, against pests in Illinois, 291.

Ants, associated with Aphids, 24, 247; associated with Coccids, 74, 110, 111, 404; destroying other insects, 111, 120, 172, 220, 305, 315, 389, 425; lizards and birds destroying, 24; measures and experiments against, 52,

74, 94, 218, 234, 253, 292, 305, 404; electrically-charged barrier for confining, 342.

Ants, Harvester (see *Pogonomyrmex*).

Ants, Leaf-cutting (see Atta).

Anuraphis chaerophylli, sp. n., on apple and Chaerophyllum in Europe, 66; considered identical with A. devecta (q.v.), 66.

Anuraphis communis (see A. devecta).

Anuraphis crataegi, apple Aphid erroneously recorded as, 66; food-plants of,

Anuraphis devecta, bionomics of, on apple in Europe, 66; synonymy of,

Anuraphis persicae-niger, bionomics and control of, on stone fruits in New South Wales, 247.

Anuraphis roseus (on apple), in Britain, 131; in Nova Scotia, 392; in U.S.A., 92, 335; sprays against, 92, 131, 335. anxia, Lachnosterna (Phyllophaga).

anxius, Agrilus.

Anychus latus, in Sudan, 412; A. orientalis not identical with, 412.

Anychus orientalis, on Citrus, etc., in Egypt and Palestine, 412; not identical with A. latus, 412.

Aonidiella aurantii (on Citrus), in South Africa, 172; in Australia, 74, 177, 216; in California, 79, 88, 89, 160, 187, 188; precautions against introduction of, into Arizona, 79; introduced into Persia from Palestine, 387; laneous food-plants of, 188; bromide against, 79, 80, 187; sprays against, 74, 88, 89; oil sprays not decreasing resistance of, to hydrocyanic acid gas, 160; natural enemies and biological control of, 172, 216; fumigation preventing natural control of, 172.

Aonidiella inopinata (see Melanaspis). Aonidiella perniciosa (see Quadraspidiotus).

aonidum, auct., Chrysomphalus (see C. ficus).

Apanteles, parasite of Vanessa cardui in Eritrea, 383; parasite of Laphygma exempta in Nyasaland, 45; parasite of Cirphis unipuncta in Queensland, 307; parasites of, 45.

Apanteles crocidolomiae, sp. n., parasite of Crocidolomia binotalis in Bihar, 119.

Apanteles diparopsidis, parasite of Bucculatrix in Nyasaland, 44.

Apanteles euproctisiphagus, sp. n., parasite of Euproctis lunata in Delhi, 119.

Apanteles fumiferanae, hosts of, in Canada, 240, 321; bionomics of, 240

В

Apanteles glomeratus, establishment of, against Pieris rapae in Australia, 215; parasite of Pieris brassicae in Europe, 406, 407; utilisation of, against P. rapae in New Zealand, 244; parasite of, 406, 407.

Apanteles gracilariae, sp. n., parasite of Gracilaria theivora in Formosa, 177;

a preoccupied name, 177.

Apanteles heichinensis, sp. n., parasite of Homona coffearia in Formosa, 177.

Apanteles homoeosonae (parasite of Homoeosoma electellum), in Cuba, 83; in U.S.A., 84.

Apanteles maculitarsus, parasite of Laphygma exempta in Nyasaland, 45;

parasites of, 45.

Apanteles ruficrus, parasite of Laphygma exempta in Nyasaland, 45.

Apanteles taiwanensis, sp. n., parasite of Euproctis taiwana in Formosa, 177.

Apanteles theae, sp. n., parasite of Homona coffearia in Formosa, 177. Apanteles transvaalensis, 45.

Apatite, 329.

Apechthis ontario, parasite of Harmologa fumiferana in Quebec, 320.

apfelbecki, Crypturgus (see C. cinereus). Aphaenogaster fulva aquia, predacious on

Cydia pomonella in W. Virginia, 220; effect of spraying on, 220.

Aphaereta muscae, parasite of Harmologa fumiferana in Quebec, 320.

Aphanotus (see Tribolium).

Aphanus spp., natural enemies and control of, attacking harvested groundnuts in Senegal, 44.

aphelinoides, Eurymyiocnema.

Aphelinus, parasite of Quadraspidiotus perniciosus in Italy, 65.

Aphelinus flavus, utilisation of, against Myzocallis annulata in Tasmania, 70.

Aphelinus mali (parasite of Eriosoma lanigerum), in Australia, 70, 247; winter storage of, in Britain, 359, 360; Washington, 98; parasitising Toxoptera graminum in Uruguay, 50; effects of DDT on activity of, 98, 247.

Aphidencyrtus, hyperparasite of Toxoptera graminum in Uruguay, 50.

aphidimyza, Phenobremia.

Aphidius, parasite of Anuraphis persicaeniger in New South Wales, 247.

Aphidius platensis, parasite of Aphids in S. America, 50.

Aphids, and plant virus diseases, 4, 5, 7, 46, 79, 93, 209, 267, 313, 355, 366, 376, 388, 413, 422, 423, 424, 426; factors affecting transmission of plant viruses by, 422-424; resistance of potato varieties to. 209; ants associated with, 24, 247; natural enemies of, 50, 66. 70, 93, 98, 192, 247, 273, 410, 411, 414, 427; tests of insecticides on, 9, 111, 228, 298, 300; electrically - charged barrier for confining, 342; classification and new species of, 65, 66, 348, 388.

Aphis abbreviata (see A. rhamni). Aphis brassicae (see Brevicoryne).

Aphis citri (see A. gossypii).

Aphis citricidus, on Citrus in Fiji, 413. Aphis durantae, experiment with Leis conformis and, in Egypt, 411.

Aphis fabae, flights of, in Britain, 196; transmitting beet mosaic in Czechoslovakia, 355; on beans in Italy, 254; on nasturtium in New York, 421.

Aphis gossypii (frangulae, auct.), in Australia, 177, 366; in Bulgaria, 4; in Egypt, 411; in Fiji, 413; in Italy, 193; in Peru, 134; in U.S.A., 9, 231, 331; in Uruguay, 50; on cotton, 134, 231, 415; on cucurbits, 4, 177, 193, 331; relation of, to cucumber mosaic in Capsicum and Passiflora edulis, 4, 366; on other plants, 413; natural enemies of, 50, 193, 411; test of dusts against, 9, 134, 331; calcium arsenate favouring infestation by, 231.

Aphis hederae, on Hedera helix in Utah,

388.

Aphis heraclella, on celery in Utah, 388. Aphis laburni, transmitting groundnut rosette in Africa, 426; sprays against, on beans in Spain, 199.

Aphis maidis, on maize in Australia and Fiji, 177, 413; parasite of, in Uruguay.

Aphis nerii, experiment with Leis conformis and, in Egypt, 411; on Asclepias in Fiji, 413.

Aphis pomi (Apple Aphis), in Bulgaria, 354; in Holland, 111; in Italy, 253; in New Hampshire, 234; in Turkey, 352; tests of insecticides against, 111, 234, 253, 352, 354.

Aphis pseudobrassicae (see Rhopalosiphum).

Aphis rhamni, insecticides against, on potato in New Hampshire, 234.

Aphis ribi-gillettei, on black currant in Utah, 388.

Aphis rumicis, auct. (see A. fabae).

Aphis spiraecola, insecticides against, on Spiraea in U.S.A., 234, 335.

Aphis, Apple (see *Aphis pomi*).

Aphis, Black Cherry (see Myzus cerasi).

Aphis, Black Peach (see Anuraphis persicae-niger).

Aphis, Cabbage (see Brevicoryne brassicae).

Aphis, Melon (see Aphis gossypii). Aphis, Oak (see Myzocallis annulata). Aphis, Pea (see Macrosiphum onobrychis). Aphis, Potato (see Macrosiphum solani-

Aphis, Red Leaf Apple, identity of, 66. Aphis, Rosy Apple (see Anuraphis roseus). Aphis, Spiraea (see Aphis spiraecola).

Aphis, Strawberry (see Capitophorus fragariae).

Aphis, Tea (see Toxoptera aurantii). Aphis, Woolly Apple (see Eriosoma lanigerum).

Aphrastobracon flavipennis, parasite of *Eublemma amabilis* in Bihar, **120**.

Aphrophora saratogensis, effect of, on pines in U.S.A., 220, 221; fungus associated with, 220, 221; DDT against, 143, 220.

Aphthona, on crucifers in England, 132. Aphthona euphorbia (on flax), in Belgium, 291; bionomics of, in Italy, 268; DDT against, 268, 269, 291.

Aphytis bovelli, parasite of Quadraspidiotus perniciosus in Italy, 191.

Aphytis mytilaspidis (parasite of Coccids), in Belgium, 49; in Bermuda, 49; in Canada, 392; sulphur sprays destroying, 392.

apicalis, Aphanus. apicicornis, Coryna. apicifera, Archytas.

Apiomorpha egeria, sp. n., bionomics of, on Eucalyptus gomphocephala in W. Australia, 69, 70.

Apion, DDT against, on Lotus corniculatus in Czechoslovakia, 278.

Apion damryi, bionomics and control of, on artichoke in Italy, 31.

Apion elegantulum, sprays against, on lucerne in Spain, 203.

Apion ulicis, establishment of, against Ulex europaeus in New Zealand, 244.

Aplomya (see Zenillia).

Apophyllum anomalum, 217.

Aporia crataegi, spray against, in Bulgaria, 354.

appalachianus, Neopristomerus (Pristomerus).

Apple, Ceresa bubalus on, in Albania, 37; pests of, in Australia, 70, 74, 106, 179, 180, 218, 245, 246, 247; pests of, in Belgium, 277, 279; pests of, in Britain, 66, 131, 132, 284, 285, 314, 315, 358, 359; Aphis pomi on, in Bulgaria, 354; pests of, in Canada, 167, 270, 271, 392, 393; new mite on, in Egypt, 412; pests of, in France, 66, 238, 293; pests of, in Holland, 66, 285, 286; pests of, in Italy, 64, 66736

191, 194, 253, 269; Anastrepha serpentina on, in Mexico, 221; Paratetranychus pilosus on, in New Zealand, 427; pests of, in Spain, 198; Argyresthia conjugella on, in Sweden, 409; pests of, in Switzerland, 296; pests of, in Turkey, 352; Eriosoma lanigerum on. in U.S.S.R., 118; pests of, in U.S.A., 12, 13, 19, 57-59, 80, 81, 92, 93, 98, 100, 101, 126, 127, 128, 149, 150, 152, 157, 188, 219, 225, 234, 256, 257, 258, 271, 322, 324, 335, 336, 338, 340, 372, 374, 399, 400, 410; Eriosoma lanigerum on, in Uruguay, 50; identity of red leaf aphis on, 66; identity of Eriophyids on, 82; not attacked by mealybugs known as Pseudococcus comstocki in Palestine, 410; effects of insecticide treatments on, 19, 157, 179, 218, 224, 253, 284, 401.

Apple (Fruit), Pristonychus terricola damaging stored, in England, 358; new mite infesting, in Italy, 114; Cydia pomonella intercepted in, in Kenya, 405; methyl-bromide fumigation of, 186; problems of insecticide residues on, 59, 71, 81, 135, 136, 162, 180, 209; removal of DDT residues from, 135, 136.

Apple, Astrakhan (see Pyrus malus var. astracanica).

Apple Aphis (see Aphis pomi).

Apple Aphis, Red Leaf, identity of, 66. Apple Aphis, Rosy (see Anuraphis roseus). Apple Aphis, Woolly (see Eriosoma lanigerum).

Apple Blister Mite (see Eriophyes pyri mali).

Apple Blossom Weevil (see Anthonomus pomorum).

Apple Bug, Green (see Neolygus communis novascotiensis).

Apple Maggot (see Rhagoletis pomonella). Apple Red Bug (see Lygidea mendax).

Apple Red Bug, Dark (see Heterocordylus malinus).

Apple Rust Mites (see Calepitrimerus baileyi and Vasates schlechtendali).

Apple Sawfly (see Hoplocampa testudinea). Apple Scab (see Venturia inaequalis).

Apricot, pests of, in Australia, 108, 247; new mite on, in Egypt, 412; Cydia molesta on, in France, 238; Hyalopterus arundinis on, in Turkey, 352; pests of, in U.S.A., 398, 399, 400. Aps 202, 337.

aquia, Aphaenogaster fulva.

Arabia, migrations of Schistocerca gregaria in, 124. arabidis, Asterolecanium.

B 2

Arachis hypogaea (see Groundnuts).

Arachnophaga hirtibasis, parasite of Stenoma decora in Brazil, 126.

Araecerus, in dried cassava roots in Uganda, 367.

Aralia polyscias, eradication of Halticus bracteatus on, in California, 187.

arboreus, Lygus.

Archips (see Tortrix).

Archips fumiferana (see Harmologa).

Archytas apicifera, tests of toxicity of DDT to, in Ohio, 258, 259.

Arctornis, Euproctis distinct from, 26.

chrysorrhoea, Arctornis auct. (see Euproctis similis).

argaula, Agonoxena.

Arge ochropa (rosae, auct.), on rose in Italy, 253; tests of thiocyanoacetate dusts against, 253.

argentatus, Orosius (Thamnotettix). argentifera, Plusia (Phytometra).

Argentina, Atta spp. in, 50, 51; cotton pests in, 126, 142; beneficial insects in, 109, 126; beneficial insects introduced into other countries from, 97, 133.

argentinensis, Porizon.

Argyresthia conjugella, on apple and Sorbus in Sweden, 409; sprays against,

Argyresthia ephippella, on fruit trees in Switzerland, 273.

Argyroploce hebesana, parasite of, on Iris germanica in U.S.A., 84.

Argyroploce leucotreta, bionomics of, in Ivory Coast, 67; on cotton in Uganda, 366, 367; causing fall of Citrus fruits in Eritrea, 108.

Argyroploce variegana, dormant sprays against, in orchards in Holland, 276. argyrospila, Tortrix (Archips).

Aristotelia urbaurea, outbreak of, on oak

in California, 400.

Arizona, miscellaneous pests in, 55, 147; beneficial insects and biological control in, 55, 147; precautions against introduction of Aonidiella against introduction of Aonidiella aurantii into, 79; Chilo spp. intercepted in, 147.

Arkansas, Aphids on spinach, mustard and oats in, 24, 25.

Arkotine, 276.

Armadillidium spp., tests of DDT dusts against, in greenhouse in Massachusetts, 159.

armata, Pyrgodera.

armatus, Anagrus; Dacus.

armicollis, Magdalis. armigera, Heliothis.

Armyworm, Yellow Striped (see Prodenia

ornithogalli).

Armyworms (see Cutworms).

Aroga trialbamaculella, bionomics of, on blueberry and cranberry in New Jersey, 137.

Arsenic, tests of toxicity of, to Agriotes, 287; risks of insecticides containing, 71, 72, 257, 287.

Arsenic Pentoxide, in baits for Lepidopterous larvae, 307; toxicity of, to Bombyx mori, 299.

Artichoke (Cynara), Vanessa cardui on, in Eritrea, 382, 383; Apion damryi on, in Italy, 31.

Artichoke, Jerusalem (see Helianthus tuberosus).

articulatus, Selenaspidus.

arundinis, Hyalopterus.

Arundo donax, Hyalopterus arundinis not developing on, 192.

Asaphes, hyperparasite of Toxoptera graminum in Uruguay, 50.

Asaphes vulgaris, parasite of Trioxys cirsii in Tasmania, 70.

Ascia monuste (on crucifers), insecticides against, in Peru, 133; in Trinidad, 319.

Asclepias, Aphis nerii on, in Fiji, 413.

Ascogaster quadridentata, parasite of Cydia nigricana in Britain, 263; parasite of C. pomonella in W. Virginia, 219; effect of sprays on, 220.

Ash (Fraxinus), mechanism of injury to, by Eulecanium coryli in Czechoslovakia, 65; Cossus cossus on, in France, 31.

Ash, types of, as carriers for dusts, 112, 114, 123, 295; unsatisfactory as carrier for dusts, 112, 114.

asparagi, Crioceris.

Asparagus, Crioceris asparagi on, in New Hampshire, 234; Halotydeus destructor on, in New Zealand, 309; Platyparea poeciloptera on, in Sweden, 409.

Asparagus Beetle (see Crioceris asparagi). Asparagus Fly (see Platyparea poeciloptera).

Aspergillus (Grey Mould), associated with mealybugs in Gold Coast, 111.

Asphondylia ricini, sp. n., parasites of, on castor in Hyderabad, 110.

Aspidiotiphagus, parasite of Quadraspidiotus perniciosus in Italy, 65.

Aspidiotiphagus citrinus (parasite Coccids), in Bermuda, 49; in Italy, 191; in Spain, 197.

Aspidiotiphagus lounsburyi, parasite of Chrysomphalus ficus in Florida, 83.

Aspidiotus destructor, on avocado California, 189; Mauritius, 210. on coconut

Aspidiotus perniciosus (see Quadraspidiotus).

Aster (Callistephus), Macrosteles divisus transmitting vellows virus of, Colorado, 231.

Aster vimineus, Lygus oblineatus on, in

Virginia, 334.

Asterolecanium arabidis, dust against, on Pittosporum tobira in California, 187.

Asterolecanium bambusae (Bamboo Scale). Chilocorus stigma introduced into Porto Rico from Florida against, 54.

Asterolecanium coffeae, on coffee in

Kenya, 404.

Asterolecanium miliaris (Bamboo Scale). Chilocorus stigma introduced into Porto Rico from Florida against, 54. Asterolecanium pustulans, on peach and fig in California, 189.

Asterolecanium variolosum, utilisation of Habrolepis dalmanni against, on oak

in Tasmania, 70.

Astrakhan Apple (see Pyrus malus var. astracanica).

ater, Hylastes.

Athalia colibri (see A. rosae).

Athalia proxima, pupation of, 122.

Athalia rosae, bionomics of, in Italy, 192. Athous, effect of, on oats in Britain, 381, 382.

atlanis, Sarcophaga. Atlox 1045A, 227. atra, Phyllotreta.

atrescens, Dacus bivittatus cucumarius. Atriplex spp., attempted use of, against Chortoicetes terminifera in Australia, 216, 217.

Atropa belladonna, Aphids transmitting tobacco rosette to, 422.

atrum, Colaspidema.

Atta, key to species of, in Argentina, 50. Atta laevigata saltensis, habits of, in Argentina, 50, 51.

Atta sexdens piriventris, habits of, in Argentina, 50, 51.

Atta vollenweiderei, habits of, in Argen-

tina, 50, 51.

Attagenus piceus, toxicity of benzenehexachloride deposits to, 235; treatments protecting wool and cloth from, 223, 224, 395; not attacking fleeces from goats dipped in DDT, 146; laboratory rearing of, 343.

attenuatus, Hylastes.

Aulacophora cattigarensis (Black Cucurbit Beetle), tests of Millettia pachycarpa against, in China, 299.

Aulacophora foveicollis, tests of DDT

against, in India, 123.

Aulacophora hilaris, on cucurbits in Australia, 177.

Aulacophora palmerstoni, on cucurbits in Australia, 177.

Aulacophora similis, on cucurbits in Pacific Is., 267, 390; on sweet potato, 267.

Aulacorthum pseudosolani (see Macrosiphum solani).

aurantii, Aonidiella: Toxoptera.

aurea, Lonchaea.

auricaudata, Ceromasia. auricularia, Forficula. auriferella, Stathmopoda,

Australia, grasshoppers in, 47, 48, 74, 216, 305, 344; miscellaneous pests in, 177, 218; pests of stored products in, 68, 181, 182, 217, 309, 364; pests and disease of tobacco in, 178, 219; vegetable pests in, 177, 178, 215, 216, 217, 218, 219; pests in buildings in, 364; biological control of insects and plants in, 215, 216; introduction of beneficial insects into other countries from, 216, 410; Phoracantha senio intercepted in Fiji in logs from, 267. (See also under the various States.)

Australia, South, orchard pests in, 246, 247; pests of stored wheat in, 181, 217; natural enemies biological control of insects in, 32, 216, 247; biological control of Hypericum

perforatum in, 216.

Australia, Western, outbreaks of Cydia pomonella on apple in, 106; new Coccid on Eucalyptus in, 69; miscellaneous pests in, 74, 244.

australis, Calosoma (see C. maderae);

Teleonemia.

Austria, forest pests in, 128, 407; Quadraspidiotus perniciosus in, 292; new parasite of Q. perniciosus in, 64. austriaca, Anisoplia.

Austroicetes cruciata, baits for, in W.

Australia, 74.

Autographa brassicae (see Trichoplusia ni). Autographa (Plusia) gamma, dust against, on flax in Belgium, 291; experiments with tobacco mosaic and, in Russia, 75. avellanae, Phytoptus.

avenae, Macrosiphum.

Avocado, new Coccid on, in Peru, 401; new Mirid on, in Uganda, 26; pests of, in U.S.A., 188, 189, 343.

Azalea, Pseudocneorrhinus bifasciatus on, in New York, 301; Gracilaria azaleella on, in Switzerland, 60; treatment of, with DDT dip against soil pests, 291. (See Rhododendron molle.)

Azalea Moth (see Gracilaria azaleella).

azaleella, Gracilaria.

Azobenzene, in sprays against *Tetrany-chus*, 103, 283, 284; cause of efficiency of, as a fumigant, 96; effect of, on plants, 103, 284.

azteca, Platylactista.

Azya (predacious on Coccus viridis), introduced into Porto Rico from Brazil, 54.

B

Babai (see Alocasia).

Bacteria, infecting cutworms, 210.

Bacterial Canker (of cherry) (see *Pseudomonas*).

Bagasse, as a carrier for derris, 52.

Bagrada cruciferarum (picta), tests of DDT dust against, in India, 123.

baileyi, Calepitrimerus.

Bait-sprays, against Diptera, 46, 47, 148, 248; against Teleonemia australis, 75.

Baits, for crickets, 4; for Ctenolepisma urbana, 364; for Dacus oleae, 201; for Hylemyia brassicae, 350; for locusts and grasshoppers, 4, 74, 109, 157, 186, 294, 305, 403, 412, 413; for moths, 132, 261, 293; for Lepidopterous larvae, 53, 62, 96, 307, 333, 372; for Tipulid larvae, 353, 369; formulae for, 4, 53, 96, 109, 132, 201, 294, 305, 307, 333, 350, 369, 412.

bajulus, Hylotrupes.

Baker's Mealybug (see *Pseudococcus maritimus*).

Balsam Fir (see Abies balsamea). balteatus, Syrphus (Epistrophe).

Bamboo (Bambusa), Coccids on, in Porto Rico and U.S.A., 54, 188; (culms), Coleoptera infesting, in Porto Rico and India, 54, 119, 122; factors affecting storage infestation of species of, by Dinoderus minutus, 54, 55.

Bamboo Scales (see Asterolecanium bambusae and A. miliaris).

bambusae, Asterolecanium.

Banana, *Pentalonia nigronervosa* and bunchy-top of, in Australia and Pacific Is., 46, 267, 413; thrips on, in Australia, 306; *Cerataphis lataniae* on, in Fiji, 413; fruit-fly infesting, in Hawaii, 249; *Cosmopolites sordidus* on, in Uganda, 367.

Banana Rust Thrips (see Scirtothrips

signipennis).

Bands, to study hibernation of Anthonomus pomorum, 314; against Cydia pomonella, 322; use of DDT in, against ants, 74.

Barathra brassicae (see Mamestra).

barbatus, Pogonomyrmex.

Barberry (see Berberis).

barbeyi, Crypturgus.

barbita, Magdalis.

Barbitistes constrictus, identification of excreta of, on conifers in Germany, 76.

Barium Chloride, spraying with, against Cleonus punctiventris, 63.

Barium Fluosilicate, in bait for Ctenolepisma, 364; experiment with, against Nymphula nymphaeata, 346.

Bark-beetles, relation of, to *Ceratosto-mella ulmi*, 163, 320; natural enemies of, 195; bibliography of, 72; bulletin on, in Britain, 32; classification of

Palaearctic, 31, 32, 348.

Barley, effect of wireworms on yield of, in Britain, 381; pests of, in Italy, 37, 383; pests of, in U.S.A., 91, 147; Aphid on, in Uruguay, 49; satisfactory food for *Limonius californicus*, 91; effect of benzene hexachloride on, 413.

Barley (Stored Seed), DDT against pests

in, 19, 38.

Barriers, against Blissus leucopterus, 154; against Laphygma frugiperda, 332. basalis, Polymerus (Poeciloscytus).

Basalt, tests of micronised, against insects in stored maize, 210.

Bassus (see Agathis).

Bathyplectes curculionis, attempted use of, against Hypera brunneipennis in Arizona, 55.

Beach Plum (see Prunus maritima).

Bean Beetle, Mexican (see Epilachna varivestis).

Bean Fly (see Agromyza phaseoli).

Beans, pests of, in Australia, 46, 47, 73, 177, 345; Sitona lineatus on, in Belgium, 291; Bruchus obtectus on, in Brazil, 76; pests of, in Britain, 173, 284; Coleoptera on, in Canada, 166, 167; new mite on, in Egypt, 411; pests of, in Fiji, 266, 267; B. obtectus on, in Holland, 130; pests of, in Italy, 254, Agromyza phaseoli on, in Mauritius, 210; pests of, in New Zealand, 138, 309; Empoasca fabalis on, in Porto Rico, 53; Aphis laburni on, in Spain, 199; Hylemyia cilicrura on, in Switzerland, 281; Dasus simplex on, in Uganda, 367; pests of, in U.S.A., 12, 13, 14, 94, 225, 229, 230, 234, 350, 370, 394, 399, 421; varieties of, in relation to Thrips tabaci, 138; unfavourable food for Limonius californicus, 91; effects of insecticide treatments on, 94, 224, 225, 234, 331, 421; risk of DDT residues on, 135.

Beans (Stored), pests in, and their control, 19, 21, 76, 385, 396, 397; (dried), unfavourable food for Ephestia elutella, 265; DDT impairing germination of, 19.

Beans, Broad (see Broad Beans).

Beans, Lima or Madagascar (see Lima Beans).

Beans, Tepary (see *Phaseolus acutifolius*). Beauveria bassiana, experiments with, infesting Cydia pomonella in W. Virginia, 220.

Beauveria globulifera, infesting Coleoptera in Rumania and Britain, 40, 315. beckii, Lepidosaphes (Mytilococcus).

Beech, Cryptococcus fagi on, in Canada, 208; Prionus coriarius on, in England, 215.

Bees, Bumble, toxicity of DDT dust to, 136.

Bees, Honey, efficiency of, in pollinating lucerne in Utah, 10, 11; risks and tests of toxicity of insecticides to, 13, 91, 100, 136, 221, 222, 237.

Bees, Solitary, toxicity of DDT dust to, 136.

Beet, pests of, in Australia, 73, 74, 177; Agallia albidula on, in Brazil, 77; Agriotes on, in Britain, 281; book on pests and diseases of, in Czechoslovakia, 355; benzene hexachloride protecting. against Melolontha in France, 280; Coleoptera on, in Hungary, Halotydeus destructor on, in Zealand, 309; Empoasca lybica on, in Palestine, 176; Acridids attacking, in Persia, 388; Chaetocnema tibialis on, in Spain, 204; pests of, in U.S.A., 11, 186, 398; insects and virus disease of, 186, 355, 423, 424; unfavourable food for Limonius californicus, 91; cryolite causing injury to, 186; risk of DDT residues on leaves of silver, 73.

Beet Flea-beetle (see Chaetocnema tibialis).

behrensi, Hepialus.

Belgium, cabbage pests in, 278, 292; pests of fruit trees and small fruits in, 277, 278, 279, 292, 353, 354; Chermes cooleyi on Pseudotsuga in, 353; thrips on vines in, 278, 292; miscellaneous pests in, 286, 291, 292; parasite of Diaspis visci in, 49.

Bell Peppers (see Capsicum).

Bemisia tabaci, transmitting virus diseases of malvaceous plants in Brazil, 299, 300.

Bengal, new gall-midge on mango in, 110; new Coccinellid attacking *Tetranychus* on jute in, 427.

bennigseni, Ootheca.

Bentonite, nicotine fixed on, 59, 126, 149, 335, 402; other uses of, in dusts and sprays, 222, 256, 288, 346; action of, on dinitro-o-cyclohexylphenol, 288.

Benzene, uses of, as a solvent, 51, 135, 224, 328, 339; synthesis and insecticidal action of derivatives of, 275.

Benzene Hexachloride, against ants, 94, 218, 234, 305; against Aphids, 13, 24, 25, 74, 134, 154, 199, 218, 229, 234, 278, 281, 293, 326, 327, 328, 331, 332, 336; ineffective against Coccids, 74; against Jassids, 134, 234, 325, 326, 336; against Lygaeids, 22, 74, 234; against Mirids, 235, 250; against Pentatomids, 13, 235, 333; against other Hemiptera, 134, 234, 303, 332, 235. 335; against Coleoptera, 13, 22, 73, 134, 137, 138, 139, 140, 148, 149, 196, 198, 199, 200, 203, 204, 217, 230, 234, 235, 236, 278, 279, 280, 281, 282, 286, 293, 296, 315, 331, 332, 333, 336, 338, 339, 358, 359, 408; against Collembola, 294, 296, 308, 331, 332, 333, 336; against locusts and grasshoppers, 13, 74, 99, 133, 134, 157, 217, 226, 227, 228, 234, 236, 237, 294, 305, 333, 341, 342, 413; brush application suspensions of, against locusts, 294; against mites, 218, 239, 281, 309, 359, 365; against sawflies, 234, 286, 296; termite nests treated with, 404, 405; against thrips, 234, 278, 306, 331; natural enemies of pests little affected by, 134, 234, 281; not increasing infestation by Diatraea spp., 304; toxicity of, to bees, 13, 222, 237; mode and mechanism of, on insects, 174, 175, 215, 275, 277, 369; fumigant action of, 25, 413, 421; method of testing, on Drosophila, 421; duration of effectiveness of, 94, 99, 134, 140, 141, 154, 226, 227, 236, 237, 251, 280, 281, 305, 413, 421; effect of, on plants, 13, 22, 69, 90, 134, 138, 149, 156, 236, 280, 309, 331, 338, 377, 413; effect of, on seed germination, 13, 90, 138, 148, 149, 156, 331, 338; question of tainting by, 13, 157, 234, 235, 236, 280, 281, 335, 336, 365; residues of, on pears, 335; question of risks and toxicity of, to man and other mammals, 215, 278, 294, 378, 413; comparison of toxicity of, to insects and vertebrates, 174, 175; in baits, 74, 294, 305, 333, 369, 412, 413; barriers of, 94, 332; in dusts, 13, 22, 24, 25, 69, 74, 90, 99, 133, 134, 137, 138, 140, 148, 149, 154, 156, 178, 203, 204, 218, 226, 227, 228, 229, 230, 233, 234, 236, 237, 239, 250, 264, 278, 281, 286, 294, 296, 303, 204, 205, 206, 208, 209, 300, 326, 237, 231 304, 305, 306, 308, 309, 326, 327, 331, 332, 333, 336, 358, 359, 365, 377, 408, 412, 413; carriers for, 13, 22, 25, 69,

90, 99, 137, 138, 140, 149, 156, 178, 217, 229, 233, 236, 237, 250, 294, 303, 333, 412; oil-impregnated dust of, 229; used as a fog, 236; spraying with, 13, 22, 23, 69, 74, 90, 157, 198, 199, 203, 218, 226, 227, 234, 235, 245, 247, 264, 278, 281, 282, 286, 293, 294, 305, 306, 309, 325, 326, 331, 332, 333, 335, 339, 340, 412, 413; atomised sprays of, 36, 341; seed treatment with, 90, 156, 279, 280, 281, 331; used as smokes, 236, 404; soil treatment with, 13, 73, 137, 138, 148, 149, 156, 338, 369; uses of, against pests of stored products, 134, 139, 141 pests of stored products, 134, 139-141, 217, 235, 236; applied in wall washes, 139, 140; applied from aeroplanes, 236, 237, 327, 328; solutions of, 333, 341, 342; emulsified solutions of, 69, 139, 218, 227, 245, 264, 281, 306, 309, 340; in wettable powder suspensions, 13, 22, 199, 227, 233, 235, 236, 281, 305, 325, 331, 335, 340, 412; acetonewater suspensions of, 90, 226; and Bordeaux mixture, 325; and neutral copper compound, 332; and DDT, 218, 264; and molasses, 90; nicotine, 326, 327; and oil emulsions, 325, 331, 335; and piperonyl cyclohexanone, 99; and pyrethrum, 99, 235; and rotenone, 154, 229; and sulphur, 148, 149, 237, 250; wetting agents for, 90, 199, 226; isomer of, 13, 22, 24, 25, 69, 90, 94, 99, 133, 134, 137, 138, 139, 140, 148, 149, 157, 174, 175, 199, 200, 215, 217, 226, 227, 228, 230, 234, 235, 236, 237, 247, 264, 277, 278, 281, 293, 294, 296, 303, 304, 305, 306, 308, 309, 325, 326, 327, 331, 332, 333, 335, 336, 338, 339, 365, 377, 378, 412, 413, 421; toxicity of other isomers of, 175, 215, 237, 278; surveys of data on, 60, 215, 280, 296, 297; proprietary preparations of, 133, 204, 237, 303, 369; derivatives of (see Polychlorcyclane Sulphide).

Benzenesulphonic Acid, toxicity of esters of, to Cydia pomonella, 93.

Berberis, Pseudocneorrhinus bifasciatus on, in New York, 301.

Berecynthus, Stal, 316.

Berecyntiscus, n.n. for Berecyntus, How., 316.

Berecyntus, How. (see Berecyntiscus). bergrothi, Helopeltis. berlesei, Caloglyphus.

Bermuda, miscellaneous pests and biological control work in, 48, 49; Diabrotica duodecimpunctata intercepted in, 49; digest of plant pest legislation in, 209.

Bermuda Cedar (see Juniperus bermudiana).

Retula (see Birch)

Betula (see Birch).

betulae, Byctiscus. bibax, Biprorulus.

bibionipennis, Aegeria.

biclavis, Howardia.

bicolor, Agathis (Bassus); Triaenodes. Bidens pilosa, Homoeosoma electellum on, in Cuba, 84.

Bidens pilosa radiata, Tingid on, in Florida, 23, 24.

bifasciata, Comperiella.

bifasciatum, Rhagium. bifasciatus, Pseudocneorrhinus.

bifasciatus, auct., Anastatus (see A. disparis).

bifidus, Stethorus. bifrons, Ootheca.

Bihar, Eublemma amabilis predacious on Laccifer lacca in, 120; beneficial insects in, 119, 120.

bimaculatus, Tetranychus. bimarginata, Haltica (Altica). binotalis, Crocidolomia.

Biological Control, of insects, suggested technique and potentialities of, in U.S.S.R., 241, 242; of noxious plants, 181, 188, 216, 244, 390.

bipartitus, Dacus (see D. bivittatus).
Biprorulus bibax, sprays against, on

Citrus in Queensland, 248. bipunctalis, Psara (Pachyzancla).

bipunctatus, Calocoris (see C. norvegicus). bipustulata, Aleochara.

bipustulatus, Chilocorus.

Birch, Cosmia trapezina on, in Belgium, 277; Prionus coriarius on, in Britain, 215; relation of Agrilus anxius to die-back of, in Canada, 208, 319, 320; Coleophora salmani on, in New Brunswick, 208.

Birch Borer, Bronze (see *Agrilus anxius*). Birds, destroying noxious insects, **24**, **109**, **167**, **255**, **312**; destroying useful insects, **67**, **244**, **425**.

Birdsfoot Trefoil (see Lotus corniculatus). Biscirus lapidarius, utilisation of, against Smynthurus viridis in Tasmania, 70.

biselliella, Tineola.

bispinosus, Dendrothrips.

bivittata, Diabrotica.

bivittatus, Dacus; Melanoplus. Black Leaf 10 Dust Base, 395.

Black Leaf 155, 101, 395, 402; preparation of DDT and, 58.

Blackberry, *Rhynchites germanicus* on, in Britain, 358; *Pantomorus godmani* on, in New South Wales, 345; pests of, in U.S.A., 83, 399.

Bladan (see Hexaethyl Tetraphosphate).

blancardella, Lithocolletis.

Blastophagus (see Myelophilus).

Blattella germanica, 342; DDT dust against, in Australia, 364.

Blissus hirtus, benzene-hexachloride dust against, in New Hampshire, 234.

Blissus leucopterus, dusts and dust barriers against, in U.S.A., 9, 12, 154; cereals, 12.

Blondelia nigripes, parasite of Athalia rosae in Italy, 193.

Blondelia virilis, parasite of Athalia rosae

in Italy, 193.

Blood Albumin, uses and effects of, in sprays, 58, 89, 189, 272.

Blueberry, pests of, in New Jersey, 137: DDT residues on fruits of, 135.

Boehmeria cylindrica drummondiana, Tingid on, in Florida, 24.

Boehmeria nivea, Acraea alciope on, in Ivory Coast, 67.

Bogosiella fasciata (parasite of Dysdercus), bionomics and distribution of, in Africa, 27.

Bolivia, digest of plant quarantine restrictions in, 105.

Bombotelia jocosatrix, new parasite of, in Formosa, 177.

Bombus, pollinating lucerne in Utah, 10, 11.

Bombyx mori, tests of insecticides on, 113, 270, 299.

Bone Dust, unsatisfactory as carrier for dusts, 112, 114.

boninsis, Pseudococcus.

Boracure, 414.

borbonica, Pelopidas.

Bordeaux Mixture, ineffective against Empoasca lybica, 176; against Epitrix cucumeris, 102; and arsenicals, 102, 137, 141, 165, 332, 373; and DDT, 102, 127, 165; possibility of reducing effectiveness of DDT, 166; facilitating removal of DDT residues from apples, 257; effect of, on dinitro-o-cresol sprays, 112: as emulsifier for oils, 90, 272; sunflower-seed oil emulsion compatible with, 354; and sulphur, 181; toxaphene compatible with, 224; effects of adjuvents on wetting power and deposits of sprays of, 42; formulae for, 90, 102, 137, 165, 181, 272, 373; formulae containing, 90, 102, 137, 141, 165, 181, 332.

borealis, Eriococcus.

Boric Acid, treatment of timber with, against Lyctus, 248; Celcure (q.v.), sometimes containing, 414.

bosai, Lissorhoptrus. botrana, Polychrosis. Botrytis, 361.

bovelli, Aphytis. bowringi, Colaphellus.

Box (see Buxus).

Boysenberry, pests of, in California, 399. Brachiaria mutica, Marasmia venilialis on, in Fiji, 267.

Brachymeria ovata, 30.

Brachymeria tachardiae, parasite of Lepidoptera in India, 120, 121; bionomics of, 121.

Brachyrhinus (see Otiorrhynchus).

Bracon, parasite of Asphondylia ricini in Hyderabad, 110.

Bracon gelechiae, utilisation of, against Gnorimoschema operculella in Bermuda and Australia, 49, 216; parasite of Recurvaria in Quebec, 321.

Bracon greeni, utilisation of, against Eublemma amabilis in Bihar, 120.

Bracon hebetor (parasite of Pyralids in stored products), in Britain, 212; Senegal, 44.

Bracon lefrovi, parasite of cotton bollworms in Punjab, 120, 121; bionomics of, 121.

Bracon mellitor, hosts of, in U.S.A., 84, 151.

Bracon nuperus, parasite of Homoeosoma electellum in U.S.A., 84.

Bracon politiventris, parasite of Tortrix packardiana in Quebec, 321.

Bracon vulgaris (see Cremnops).

bracteatus, Halticus.

Bran, Tribolium castaneum in, 68; in baits, 62, 74, 96, 109, 157, 158, 305, 307, 333, 353, 369, 412, 413.

brasiliensis, Eutinobothrus (Gasterocercodes).

Brassica campestris, Calocoris norvegicus on, in Italy, 383.

brassicae, Brevicoryne (Aphis); Charips; Hylemyia (Chortophila); Mamestra (Barathra); Pieris; Plusia (Autographa) (see Trichoplusia ni).

Brazil, Stenoma decora on cacao, etc., in, **125, 126;** *Coelosternus* on cassava in, **6**; new Curculionid on rice in, 8; miscellaneous pests in, 5, 76, 104; insects and virus diseases of plants in, 7, 76, 77, parasite of S. decora in, 126; beneficial insects introduced into Porto Rico from, 53.

Brevicoryne brassicae (on crucifers), in Australia, 74, 218; in Italy, 254; in New Hampshire, 234; in Trinidad, 319; in U.S.S.R., 118; parasite of, in Uruguay, 50; insecticides against, 74, 118, 218, 234, 254, 300, 421; effects of insecticides on production of young by, 421.

Brevipalpus obovatus, importance of, on tea in Ceylon, 368.

Brevipalpus oudemansi, B. pyri recorded as, in Egypt, 412.

Brevipalpus pyri, sp. n., on fruit trees in Egypt, 412.

brevipes, Pseudococcus.

brevis, Hoplocampa; Kalotermes (Cryptotermes).

brevistylus, Dacus (see D. ciliatus).

Bridal Wreath (see Spiraea vanhouttei). brillians, Harrisina.

Britain, forest pests in, 32, 214, 315; pests of greenhouse and ornamental plants in, 32, 284, 288, 360, 361, 362; pests of hops in, 287, 288, 358; Sciara on mushrooms in, 284; orchard pests in, 66, 131, 132, 173, 284, 287, 288, 314, 315, 358, 359; pests of small fruits in, 284, 358, 359, 408; Aphid and virus diseases of strawberry in, 312, 313, 314, 376; pests of stored products in, 210, 264, 282, 355, 358, 417, 418; pests of vegetable and root crops in, 132, 173, 190, 195, 196, 261, 281, 284, 288, 360, 361, 362, 377, 417; Phyllotreta spp. transmitting new virus disease of turnip in, 377; wireworms in, 281, 381; books on pests in, 173, 357; beneficial insects in, 174, 212, 215, 263, 288, 314, 315, 359, 417; plant quarantine legislation in, 209; Ceratostomella ulmi possibly introduced into Canada in crates from, **163.** (See also Jersey.)

britannicus, Dynaspidiotus.

British Columbia, Cecidophyes ribis in, 83; pests of forest and shade trees in, 78, 240; Epitrix tuberis in, 167; beneficial insects introduced into eastern Canada from 78.

Broad Beans, new mite on, in Egypt, 411; Halotydeus destructor on, in New

Zealand, 309.

Broccoli, *Plutella maculipennis* on, in Porto Rico, **52**; pests of, in U.S.A., **159**.

Brochymena quadripustulata, sprays against, on peach in S. Carolina, 235.

Brome Grass (see Bromus).

Bromo-DDT [1, 1, 1-trichlor-2, 2-bis (parabromophenyl)ethane], tests with, as an insecticide, 14, 237, 274, 275, 421, 422; liberation of HCl by, 275.

α-Bromo-naphthalene, as greenhouse fumigant against *Tetranychus*, 103.

Bromus, Cephus cinctus on, in N. Dakota, 128.

Brontispa longissima, on coconut in New Hebrides and Solomon Is., 390. brouni, Kalotermes (Calotermes).

Brown-tail Moth (see Euproctis chrysor-rhoea).

Bruchidius (see Bruchus). bruchocida, Lathromeris.

Bruchophagus, species of, infesting leguminous seeds in Sweden, 408.

Bruchophagus gibbus (funebris), 408; on Lotus corniculatus in New York, 71.

Bruchus analis, in stored soy beans in Uganda, 367.

Bruchus chinensis, experiments with types of petrol against, in stored seeds in S. Africa, 2.

Bruchus floridae, parasites of, on Amorpha fruticosa in Oklahoma, 9.

Bruchus maculatus, insecticides against, in stored cowpeas in S. Carolina, 369.

Bruchus obtectus (obsoletus, auct.), in field and stored beans in Brazil, 76; in field beans in Holland, 130; introduced and eradicated in Poland, 385; in U.S.A., 21; bionomics of, 76, 130; measures against, in stored beans, 21, 76; parasites of B. pisorum reared on, in Hungary and Ukraine, 62, 242, 243.

Bruchus pisorum (Pea Bruchid), bionomics of, in Hungary, 61, 62; in Russia, 242, 243; in U.S.A., 12, 14; natural enemies of, 62, 242, 243; measures and experiments against, 12, 14, 62.

Bruchus unicolor, new parasite of, on Onobrychis sativa in Russia, 242.

brumata, Operophtera. brunneipennis, Hypera.

brunneus, Lyctus.

Brussels Sprouts, Hylemyia spp. on, in Britain, 173; pests of, in U.S.A., 159.

Bryobia praetiosa (on apple), in Australia, 74, 179, 180, 218, 246, 247; in Nova Scotia, 393; on pear, 179, 246, 247; sprays against, 180, 218, 246, 393; effect of DDT on infestation by, 74, 179, 218, 246, 247; not affected by copper sprays, 393.

bubalus, Ceresa.

Bucculatrix, parasite of, on sweet potato in Nyasaland, 44.

Bucculatrix ruficoma, 44.

Buckeye, California (see Aesculus californica).

Buddleia lindleyana, insecticidal properties of, 299.

Budworm, Spruce (see *Harmologa fumi-ferana*).

Budworm, Tobacco (see *Heliothis vires-cens*).

Building Timbers, Coleoptera in, 107, 414; termites in, 169, 249, 414.

Bulgaria, Aphids and virus diseases of *Capsicum annuum* in, 4, 5; orchard pests in, 354.

Bumelia laetevirens, Anastrepha serpentina on, in Mexico, 221.

buoliana, Rhyacionia.

Bupalus piniarius, identification of excreta of, on pine in Germany, 76.

Burma, parasite of Scirpophaga nivella on sugar-cane in, 119.

burrelli, Allotropa.

Bursera simaruba, used in tests of repellents against termites, 233.

Busseola fusca, on maize in Kenya, 405. Butea frondosa, host-plant of Laccifer lacca in Bihar, 120.

 β -Butoxy- β -thiocyanodiethyl Ether (see Thiocyanates).

Butter, determination of DDT in, 393. buttricki, Agathis (Bassus).

n-Butyl Pentachlorphenyl Ether, 224. buxi, Monarthropalpus.

Buxus (Box), pests of, in U.S.A., 143, 187, 400.

sempervirens, Chrysomphalus Buxus dictyospermi on, in Persia, 386.

Byctiscus betulae, bionomics and control of, on vines in Spain, 198.

Byctiscus populi, characters distinguishing B. betulae from, 198.

Cabbage, pests of, in Australia, 74, 177, 178, 218, 305; pests of, in Belgium, 278, 292; pests of, in Britain, 132, 173; Phyllotreta aerea on, in Canada, 167; Lixus latro on, in Eritrea, 384; Crocidolomia binotalis on, in Fiji, 390; pests of, in Holland, 285, 286; Phyllotreta cruciferae on, in India, 122; pests of, in Italy, 253, 254, 383; pests of, in New Zealand, 244; Pierids on, in Peru, 133; flea-beetles on, in Spain, 204; pests of, in Switzerland, 281; Brevicoryne brassicae on, in U.S.S.R., 118; pests of, in U.S.A., 13, 91, 102, 154, 159, 160, 222, 234; pests of, in West Indies, 52, 53, 319; not attacked by Athalia rosae, 192; unfavourable food for Limonius californicus, 91; benzene hexachloride not affecting flavour of, 234; risks of DDT deposits on, 74, 81, 135; not injured by Velsicol AR-60, 421.

Cabbage, Chinese, Phyllotreta spp. transmitting new turnip virus to, in Britain, 377; injured by Velsicol AR-60, 421. Cabbage Aphis (see Brevicoryne bras-

sicae).

Cabbage Bug, Harlequin (see Murgantia histrionica).

Cabbage Looper (see Trichoplusia ni).

Cabbage Maggot (see Hylemyia brassicae).

Cacao, Coccids on, in W. Africa, 110; Stenoma decora on, in Brazil, 125; pests of, in Fiji, 266, 413; mealybugs transmitting swollen-shoot disease of, 111, 112.

experiments Cacao (Stored), with Ephestia elutella and, 213, 265.

Cacoecia (see Tortrix).

cacoeciae, Angitia (Horogenes).

Cactoblastis cactorum, reduction importance of, on *Opuntia* in S. Africa, 171, 172, 389.

Cadmium Fluoride, repellency of wood treated with, to Kalotermes brevis, 232.

Caesalpinia sepiaria, new Mirids on, in Uganda, 26.

caesar, Zenillia (Aplomya, Epimasicera).

caespitum, Tetramorium.

Cages, for rearing and studying insects, 77, 78, 258; for sampling of grasshopper populations, 166.

Cajanus cajan (indicus), Mirids on, in Uganda, 26.

Calacarus adornatus, on Camellia California, 187.

Calamotropha leptogramella, bionomics and control of, on *Paspalum* in Queensland, 308.

Calandra, tests of benzene hexachloride against, 139, 217, 234, 235; tests and uses of DDT against, 19, 28, 38, 39, 134, 139, 217, 292, 324, 369.

Calandra granaria (in stored cereals), in Australia, 182; in Italy, 30, 38; in Switzerland, 38, 39; measures and experiments against, 19, 30, 38, 39, 182.

Calandra oryzae, on field maize in New Hampshire, 234; (in stored cereals), in S. Africa, 2; in Australia, 182, 217, 309, 310; in Mauritius, 210; in Peru, 134; in Uganda, 367; in U.S.A., 14, 85, 369; in dried cassava roots, 367; strains of, 182, 217, 310; tests and use of fumigants against, 2, 20, 85, 182, 217; other measures and experiments against, 14, 19, 28, 52, 134, 210, 217, 234, 235, 369.

calceolariae, Pseudococcus. Calcite, 330. (See Calcium Carbonates.) Calcium Arsenate, against pests crucifers, 41, 204; against cotton pests, 14, 104, 184, 231, 232, 370, 371, 401; favouring infestation of cotton by Aphids and Heliothis, 231, 371; against pests of potato, 102, 165, 196, 332, 359; against insects on other plants, 4, 10, 141; toxicity of, to bees, 222; in cutworm baits, 96; formulae for, in dusts, 232, 332; adhesive qualities of dusts of, 114; in mopping mixture, 370; spraying with, 41, 42, 102, 104, 141, 165, 196, 332; applied from aircraft, 401; and Bordeaux mixture, 102, 141, 165, 332; and yellow cuprous oxide, 165; and DDT, 232, 371; and nicotine, 371; oils improving tenacity of spray deposits of, 42; sunflower-seed oil emulsion compatible with, 354; and rotenone, 371; and soap, 104; and sulphur, 184, 232; Toxaphene compatible with, 224; other insecticides compared with, 14, 96, 104, 165, 299; oxides reducing effect on plants of soil containing, 371.

Calcium Carbonates, as diluents for dusts, 218, 329, 330; tests of effect of, on larvae of *Epilachna varivestis*, 330. (See Chalk and Limestone.)

Calcium Chloride, and oil emulsion, spraying with, against mites, 361.

Calcium Cyanamide, experiment with, against Nymphula nymphaeata, 346; used to defoliate cotton, 401.

Calcium Cyanide, soil treatment with, against *Rhabdopterus picipes*, 137; dusting with, against *Tomaspis flavilatera*, 304.

Calcium Hydroxide (see Lime).

Calcium Oxide (see Quicklime).

Calcium Sulphate (see Gypsum).

Calepitrimerus baileyi, on apple in U.S.A., 83.

Calepitrimerus vitis, on vines in U.S.A., 83.

California, Coccids on almond in, 156, 398, 399; orchard pests in, 83, 92, 185, 188, 189, 397, 398, 399, 400; pests of *Citrus* in, 79, 88, 89, 145, 155, 160, 187, 188, 189, 343; pests of other sub-tropical fruits in, 188, 189, 343, 399; pests of bush-fruits in, 83, 399; forest pests in, 301, 400; Coccids on olive in, 16, 187, 188, 342, 399, 400; pests of ornamental plants in, 142, 154, 185, 187, 188, 399, 400; *loftini* on cereals and sugar-cane in, 146, 147; Desmoris spp. on sunflower in, 150; pests of vegetable and root crops in, 12, 56, 97, 187, 396; Eutettix tenellus and curly-top of beet and vegetables in, 186, 397, 398; pests of vines in, 147, 185, 190, 397, 398; walnut pests in, 80, 155, 188, 189, 399; grasshoppers in, 186, 398; Lygus spp. affecting seed production of lucerne and Parthenium in, 90, 94; miscellaneous pests in, 187, 399; enemies and biological control of insects in, 91, 97, 145, 147, 156, 160, 187, 343, 397; parasites of *Chrysopa* in, 396; Coleoptera imported into, against *Hypericum perforatum*, 188, 216; beneficial insects imported into other countries from, 49, 216; pests intercepted in quarantine in, 147, 187, 188; measures to prevent introduction of pests into, 79, 398; *Diaspis visci* probably introduced into Bermuda from, 48; precautions against introduction of *Aonidiella aurantii* into Arizona from, 79.

California Buckeye (see Aesculus californica).

californicus, Cybocephalus; Limonius.

Caliroa aethiops, spray against, on rose in U.S.A., 335.

Caliroa limacina, tests of thiocyanoacetates against, on pear in Italy, 253; Pentatomid predacious on, on cherry in New South Wales, 108.

Calliceras (see Ceraphron).

Calliphora vomitoria, tests of DDT and related compounds against, 274.

Calliptamus italicus, damaging crops in Persia, 387, 388.

Calliptamus turanicus, damaging crops in Persia, 387, 388.

Callipterus juglandis (see Chromaphis).

Callosobruchus (see Bruchus).

Calocoris norvegicus (bipunctatus), on flax in Eire, 208; bionomics and control of, on beans, etc., in Italy, 383.

Caloglyphus berlesei, in copra in Fiji, 266. Caloglyphus mycophagus, C. berlesei recorded as, 266.

Calomel (see Mercurous Chloride).

Calophyllum antillanum, tests of toxicity of, to insects, 52.

Calophyllum inophyllum, Stictoptera sub obliqua on, in Fiji, 266.

Calopogonium coeruleum, tests of toxicity of seeds of, to insects, **52**.

Calosoma maderae (australis), predacious on Cirphis unipuncta in Queensland, 307.

Calotermes (see Kalotermes).

Calymnia (see Cosmia).

Camelina sativa, not attacked by Athalia rosae in Italy, 192.

Camellia, Chrysomphalus dictyospermi on, in Persia, 386; pests of, in U.S.A., 187, 188, 190.

camelliae, Aceria.

Cameroons, Antestia lineaticollis on coffee in, 295; Dacus bivittatus and its parasite in, 176, 177.

campestris, Lygus.

Camphor (see Cinnamomum camphora).

Camponotus compressus, predacious on Eublemma amabilis in Bihar, 120.

Camponotus picipes, predacious Eutermes in Br. Guiana, 305.

Campoplex haywardi, sp. n., parasite of Gnorimoschema operculella in Argentina, 109.

Campoplex sulcatellus, Meteorus loxostegei destroying, in Loxostege sticti-

calis, 29.

Campsomeris erythrogaster, utilisation of, against Clemora smithi in Mauritius, 210.

Campsomeris lachesis, utilisation of, against Clemora smithi in Mauritius, 210.

Campsomeris phalerata, utilisation of against Clemora smithi in Mauritius, 210.

Campsomeris pilosella, parasite of Clemora smithi in Mauritius, 210.

Camptoptera, 407.

Camptylochila aemula, parasites of, on

spruce in Quebec, 321.

Canada, cereal pests in, 390, 391; forest pests in, 240, 255; grasshoppers in, 166, 341; mite on plum in, 83; Laemophiceus in stored grain in, 418; beneficial insects and biological control in, 28, 255, 391; beneficial insects introduced into New Zealand from, 244. (See also under the various Provinces.)

canadensis, Polistes.

cancellata, Schistocerca.

Canella winteriana, toxicity of bark of, to Andrector, 52.

canella, Diatraea.

canescens, Nemeritis.

Cankerworm, Fall (see Alsophila pometaria).

Cannabis sativa (see Hemp).

Cantaloupe (see Melon).

capitata, Ceratitis.

capitella, Incurvaria.

Capitophorus fragaefolii, on strawberry in Belgium, 353.

Capitophorus fragariae, relation of, to virus diseases of strawberry in Britain, 313, 376; on Potentilla, 313; measures against, 314.

Capitophorus potentillae, on Potentilla anserina in Britain, 313; not transmitting strawberry crinkle virus, 313.

Capitophorus tetrarhodus, on rose in Britain, 313; strawberry virus experimentally transmitted by, 313.

Capnodis spp., on Pistacia in Persia, 386. Capsicum, Agallia albidula on, in Brazil, 77; Empoasca lybica on, in Palestine, 176; Scapteriscus vicinus damaging, in U.S.A., 94; effect of benzene hexachloride on flavour of, 234.

Capsicum annuum, Aphids and virus diseases of, in Bulgaria, 4, 5; Diptera on, in Libya, 72.

Capua (see Tortrix). Carbinol DDT, 421.

Carbolineums (see Tar Distillates).

Carbon Bisulphide, soil fumigation with, against Leptinotarsa decemlineata, 259; against pests of stored products, 2, 62, 76, 85, 182, 217; mixtures of other fumigants with, 85; as a solvent, 51.

Carbon Dioxide, tolerance of Rhizopertha

dominica to, 364, 365.

Carbon Tetrachloride, fumigation with, against pests of stored products, 12, 20, 85, 217; mixture of other fumigants with, 12, 20, 85, 217; not reducing germination of seed maize, 85; study of absorption of, by wheat, 223; as solvent for organic insecticides, 94, 161; risks of using, 85, 161.

carbonaria, Capnodis.

cardinalis, Dysdercus; Rodolia (Novius). Cardol, repellency of chlorinated, to Kalotermes brevis, 232.

Cardoon, Vanessa cardui on, in Eritrea,

cardui, Vanessa (Pyrameis).

caridei, Coccophagus; Sarcophaga (Acridiophaga).

carinatus, Eriophyes. cariosa, Capnodis.

Carnation (in greenhouses), forms of *Macrosiphum solanifolii* on, 361, 362; *Tetranychus* on, 288, 290; not injured by azobenzene, 284; selenium treatment of, 290.

Carolina, North, Chermes on Pinus strobus in, 333, 334; Diabrotica duodecimpunctata on maize in, 148; Protoparce spp. on tobacco in, 87.

Carolina, South, miscellaneous pests in, 13, 369–372; Pantomorus in, 184; pests of peach in, 235, 370; biological control of Cydia molesta in, 370.

Carpet Beetle, Black (see Attagenus piceus).

carpini, Tetranychus (Eotetranychus). Carpinus betulus, pests of, in Switzerland,

Carpocapsa (see Cydia).

carpocapsae, Mastrus.

Carpophilus, in shelled ground-nuts in U.S.A., 25.

Carpophilus dimidiatus, in stored maize in Uganda, 367.

Carrot, Diptera on, in Britain, 173, 377, 417; Melolontha on, in France, 280; pests of, in Switzerland, 280, 281;

Cercopids on, in Wisconsin, 160; Anuraphis crataegi on, 66; unfavourable food for Limonius californicus, 91; question of effect of benzene hexachloride on flavour of, 235, 280.

Carrot Fly (see Psila rosae). carvae, Acrobasis; Curculio.

casei, Tyrolichus.

Casein, used for feeding insects, 148, 343; in spray formulae, 28, 148, 181.

Cashew (see Anacardium occidentalis). Cashew-nut Shell Oil, repellence of crude and chlorinated fractions of, to

Kalotermes brevis, 232.

Cassava. Coelosternus on, in Brazil and W. Indies, 6; Zonocerus elegans on, in Nyasaland, 44; Amblypelta cocophaga on, in Solomon Is., 390; pests in stored roots of, in Uganda, 367.

Cassava Flour, as carrier for locust bait,

294.

Cassia alata, tests of toxicity of parts of, to insects. 52.

Cassia fistula, Toxoptera aurantii on, in Fiji, 413.

Cassia spectabilis, tests of toxicity of parts of, to insects, 52.

cassiae, Pachymerus.

Cassida, not parasitised by Lathromeris bruchocida, 242.

Cassida nebulosa, egg parasite of, in U.S.S.R., 242. castaneum, Tribolium.

Castnia licoides (licus, auct.), on sugarcane in Br. Guiana, 304.

Castor, Xyleborus fornicatus on, in Ceylon, 251; Anychus orientalis on, in Egypt, 412; pests of in Hungary, 62; pests of, in India, 110, 123; leafhoppers on, in Palestine, 176; pests of, in Uganda, 26, 367; not attacked by Athalia rosae, 192.

Castor Oil, Dehydrated (see Sostroil).

Casuarina equisetifolia, new Coccid on, in Gold Coast, 110; Celosterna scabrator on, in Réunion, 317.

Catalpa, pests of, in U.S.A., 143.

catalpae, Ceratomia.

Catantops melanostictus, on Eleusine in Uganda, **367.**

Catolaccus, parasite of Bruchus floridae in Oklahoma, 9.

cattigarensis, Aulacophora.

caudata, Desiantha; Lepidiota; Tettigonia.

caudatus, Ephialtes.

Cauliflower, pests of, in Australia, 74, 218; Lixus latro on, in Eritrea, 384; Helophorus on, in Holland, 190; Ceuthorrhynchus quadridens on, in Switzerland, 281; pests of, in Trinidad,

319; pests of, in U.S.A., 159; not injured by Velsicol AR-60, 421.

cautella, Ephestia.

Cavariella aegopodii, flights of, in England, 196.

cavus, Dibrachys.

Cecidomyiids, book on, 308.

Cecidophyes ribis, on currant in Br. Columbia, 83.

Cedar Oil, repellence of chlorinated, to Kalotermes brevis, 232.

Cedrela, new Mirid on, in Uganda, 26.

Ceiba pentandra, Stenoma decora on, in Brazil, 125; new Coccid on, in Gold Coast, 110; Zygaenid on, in Ivory Coast, 67.

Celcure, tests of, for treating timber against Anobium, 414, 415; composi-

tion of, 414.

Celery, Anuraphis crataegi on, in Europe, 66; pests of, in U.S.A., 102, 388; virus disease of, 388; DDT deposits on, 135.

Celosterna scabrator, bionomics of, in Réunion, 317; not occurring

Mauritius, 317.

Celtis occidentalis, Lachnosterna spp. on, in Texas, 14.

Cemiostoma (see Leucoptera).

Centella asiatica, insecticidal properties of, 299.

Centipedes, predacious on Prionus coriarius in Britain, 215.

cepae, Petrobia.

Cephaleia abietis, characters of excreta of, on spruce in Germany, 76.

cephalonica, Corcyra.

Cephalonomia, parasite of Lygaeids in Senegal, 44.

Cephalosporium lecani, infesting Saissetia oleae in Persia, 385.

Cephus cinctus (on wheat), in Canada, in N. Dakota, 128; factors affecting diapausing larvae of, 390, 391; on grasses, 128.

Ceraphron, parasite of Apanteles transvaalensis in Nyasaland, 45.

cerasi, Myzus.

cerasivorella, Coleophora.

Cerasus laurocerasus (see Prunus).

Cerataphis lataniae, food-plants of, in Fiji, **413.**

Ceratia (see Aulacophora).

Ceratitis capitata, causing fall of Citrus fruits in Eritrea, 108; on Capsicum annuum in Libya, 72; eradication of, in New South Wales, 248.

Ceratomia catalpae, spray against, on

Catalpa in U.S.A., 143.

Ceratostomella ulmi (causing Dutch Elm Disease), introduction and possible vectors of, in Quebec, 163, 320; Scolytids transmitting, in U.S.A., 320; measures against, 163.

Cercyonia citri, on orange in Ivory Coast,

68.

cerealella, Sitotroga.

Cereals, Laphygma spp. on, in E. Africa, 426; Eurygaster integriceps on, in Tadzhikistan, 241; Blissus leucopterus on, in U.S.A., 12; Acridids and Tettigoniids damaging, 45, 48, 109, 388. (See Maize, Wheat, etc.)

Cereals (Stored), pests in, and their control, 2, 30, 38, 39, 68, 133, 134, 181, 182, 210-214, 217, 309, 310, 319, 348, 364, 367, 405, 417, 418; machine for destroying insects in, 282; high frequency electric fields against insects in packages of, 15; studies of sorption of fumigants by, 39, 223; methods and equipment for sampling, in silos, 419; book on storage of, 405.

Ceresa bubalus, on fruit trees in Albania,

37.

Ceresan-nieuw, seed treatment with, increasing injury to onion by *Hylemyia antiqua*, 283.

Cermatulus nasalis, predacious on insects in New South Wales, 108.

Cerococcus coffeae, sp. n., on coffee in Belgian Congo and E. Africa, 316, 317; type of injury caused by, 317.

Cerococcus hibisci, C. coffeae recorded as, on coffee in E. Africa, 316, 317.

Ceromasia auricaudata (parasite of Harmologa fumiferana), in Br. Columbia, 78; introduction of, into eastern Canada, 78.

cerophaga, Angitia.

Ceroplastes destructor, sprays ineffective against, on Citrus in W. Australia, 74.

Ceroplastes sinensis, introduction and eradication of, on Citrus in Persia, 387. Ceroputo yuccae (see Puto).

Cerotoma ruficornis (see Andrector).

Cerotoma trifurcata, test of amorpha dust against, in Oklahoma, 9.

cervinus, Polydrusus.

Ceuthorrhynchus napi, benzene hexachloride against, on rape in Switzerland, 281.

Ceuthorrhynchus pleurostigma, insecticides against, on cabbage in Switzerland, 281.

Ceuthorrhynchus quadridens, insecticides against, on cabbage and rape in Switzerland, 281.

Ceylon, pests of tea, etc., in, 251, 368, 369; parasites of Agromyza theae in, 368.

chaerophylli, Anuraphis (Yezabura).

Chaerophyllum, new Aphid on, in Central Europe, 66.

Chaetocnema concinna, on crucifers in England, 132.

Chaetocnema tibialis, DDT against, on beet in Spain, 204.

Chaetonodexodes marshalli, parasite of Schematiza cordiae in Trinidad, 425, 426; parasite of, 425.

Chaetoptelius vestitus (see Hylesinus).

chalcites, Plusia.

Chalcodermus aeneus, dusts against, on cowpeas in S. Carolina, 370.

Chalcoides (see Crepidodera).

Chalcophora, on Parthenium argentatum in Mexico, 97.

Chalepus dorsalis, spray against, on Robinia in U.S.A., 143.

Chalk, as carrier for dust, 250; in insecticide wall washes, 139.

Charips spp., hyperparasites of Toxoptera graminum in Uruguay, 50.

Cheese, mites infesting, in New Zealand and Queensland, 244, 363, 416; experiment with new Tyroglyphid and, 114.

Cheiloneurus, parasite of Pseudococcus njalensis in Gold Coast, 111.

Chelonus altitudinis, hosts of, in U.S.A., 84.

Chelonus phthorimaeae, utilisation of, against Gnorimoschema operculella in Bermuda and Australia, 49, 216.

Chenopodium, Pantomorus godmani on, in New South Wales, 345.

Chenopodium album, Lygus oblineatus on, in Virginia, 334.

Chermes, on conifers in Britain, 315; types of damage to *Pinus strobus* by species of, in N. Carolina and Canada, 333, 334; spray against, 334.

Chermes cooleyi, on Pseudotsuga taxifolia in Belgium, 353.

Chermes piceae, on Abies balsamea in New Brunswick, 208.

Chermes pinifoliae, on Pinus strobus in New Brunswick, 334.

Chermes strobi, on Pinus strobus in N. Carolina, 333, 334.

Cherry, pests of, in Britain, 358; Cydia molesta on, in France, 238; Operophtera brumata on, in Holland, 276; Hyphantria cunea on, in Hungary, 254; Caliroa limacina on, in New South Wales, 108; Neurotoma nemoralis on, in Poland, 254, 255; pests of, in U.S.A., 152, 234, 335, 399; effect of soil treatment with benzene hexachloride on, 280.

Cherry Aphis, Black (see Myzus cerasi). Cherry Sawfly (see Caliroa limacina).

Chicory, Agallia albidula on, in Brazil, 77. Chile, grasshoppers in, 108, 109.

Chilo, intercepted in U.S.A., 147.

Chilo loftini, on sugar-cane and cereals in Mexico and U.S.A., 146, 147.

Chilocorus bipustulatus (predacious on Coccids), in Persia, 385; in Spain, 197. Chilocorus stigma, introduced into Porto

Rico from Florida against Asterole-

canium spp., 54.

Chilonectria cucurbitula, Cercopid associated with, on pines in U.S.A., 220, 221.

China, tests of indigenous plants as insecticides against miscellaneous insects in, 298, 299; book on vegetable pests in, 3.

China Clay, as carrier for dusts, 137, 140, 303; spray powder of azobenzene and, 283, 284.

Chinch Bug (see Blissus leucopterus). Chinch Bug, Hairy (see Blissus hirtus). chinensis, Bruchus (Callosobruchus).

2-Chloracetonitrile, tests with, against insects in stored wheat, 20; and carbon tetrachloride, 20.

2-Chloracrylonitrile, tests with, against insects in stored wheat, 20; mixtures, 20.

Chlordane, ants' nests treated with, 94; against grasshoppers, 99, 225, 226, 227, 228, 237, 341; median lethal doses and deposits of, to grasshoppers, 226, 227, 342; not affecting crickets or *Epicauta* in pasture, 228; against Lepidoptera, 331, 332, 333; against *Peregrinus maidis*, 332; against *Popillia* japonica, 339, 340; ineffective against Rhopalosiphum pseudobrassicae, 332; possibly not toxic to bees, 222; mode of action of, on insects, 226, 277; duration of effectiveness of, on plants, 226, 227, 228, 237, 340; question of toxicity of, to vertebrates, 94, 227, 339; and injury to plants, 228, 331; in aerosols, 228; in baits, 333; in dusts, 99, 225, 226, 227, 237, 331, 332; in dust barrier 322; in approach 225, 226 dust barrier, 332; in sprays, 225, 226, 227, 228, 237, 339, 340; applied from aeroplanes, 228, 237; solutions of, 94, 228, 333, 341, 342; emulsified solutions of, 226, 227, 228, 237, 340; acetonewater suspensions of, 226; in wettable powders, 226, 227.

4-Chlor-3', 4'-dimethyldiphenyltrichlorethane, 275.

4-Chlordiphenyltrichlorethane (1-Phenyl-1-p-chlorphenyl-2, 2, 2-trichlorethane), 275, 421.

Chloretone, ineffective against Nymphula nymphaeata, 346.

Chlorinated Naphthalene (see Naphthalene, Chlorinated).

Chlorita flavescens (see Empoasca). Chlorita signata (see Empoasca lybica).

4-Chlor-4'-methyldiphenyltrichlorethane,

bis-(4-Chlor-2-methylphenyl)-dichlorethane, 275.

Chlorochroa juniperina, on Juniperus communis in Poland, 268.

Chloroclystis rectangulata, dormant sprays against, in orchards in Holland, 276.

Chloroform, toxic to Tyroglyphus nadinus,

Chlorophora excelsa, Phytolyma lata on, in Ivory Coast, 68.

Chlorophorus annularis, infesting bamboos in India, 119; digestive enzymes of larvae of, 119.

1-(4-Chlorphenyl)-2, 2, 2-trichlorethanol, 421.

4 - Chlor - 4' - n - propyldiphenyltrichlorethane, 275.

Chortoicetes terminifera (in Australia), outbreaks of, 47, 48, 216; parasites of, 344; measures and experiments against, 48, 216, 217.

Chortophaga viridifasciata, water exchange between air and, 375.

Chortophila (see Hylemyia).

christenseni, Ameloctonus.

Chromaphis juglandicola, DDT spray controlling, on walnut in California,

Chromaphis juglandis, tests with Prangos against, on walnut in Kirghizia, 118.

Chrotogonus rotundatus, attacking ground-nuts in Tanganyika, 426.

Chrotogonus turanicus, damaging crops in Persia, 387, 388.

chrysanthemi, Diarthronomyia.

Chrysanthemum, pests of, in U.S.A., 188, 289, 290, 291; treatment of, with selenium compounds, 289, 290, 291.

Chrysanthemum cinerariaefolium, Lamellicorns on, in Uganda, 367; quality of pyrethrum obtained from, in Holland, 270.

Chrysanthemum leucanthemum, Lygus oblineatus on, in Virginia, 334.

Chrysanthemum Midge (see Diarthronomyia chrysanthemi).

chrysocephala, Psylliodes.

Chrysocharis laricinellae, utilisation of, against Coleophora laricella in New Brunswick, 208.

Chrysolina (see Chrysomela).

Chrysomela gemellata, Rossi (see C. quadrigemina).

Chrysomela hyperici (attacking Hypericum perforatum), use of, in Australia, 216; establishment of, in California, 188, 216; spread of, in New Zealand, 244; proposed use of, against H. androsaemum, 244.

Chrysomela lapponica (see Melasoma). Chrysomela quadrigemina (attacking Hypericum perforatum), use of, in Australia, 216; establishment of, in

California, 188, 216.

Chrysomphalus agavis, parasite of, on Juniperus bermudiana in Bermuda, 49. Chrysomphalus aonidum, auct. (see C.

ficus).

Chrysomphalus dictyospermi, treatment of limes imported into California from Mexico against, 79; natural enemies, food-plants and control of, in Persia, 385, 386, 387.

Chrysomphalus ficus, measures to prevent introduction of, into California from Mexico, 79; in Egypt, 411; in Florida, 83; in Trinidad, 319; on Citrus, 79, 83, 319; cultivation increasing infestation by, 83; natural enemies of, 83; experiment with Leis conformis and, 411.

Chrysomphalus obscurus (see Melanaspis). Chrysopa, parasites of, in California, 396; predacious on Aphids in Holland and New South Wales, 66, 247; in Peru, 134; not affected by DDT and benzenehexachloride dusts, 134.

Chrysopids, list of parasites of, 396. chrysorrhoea, L., Euproctis (Phalaena). chrysorrhoea, auct., Euproctis (Arctornis) (see E. similis).

chrysosticta, Angitia.

Cicindela punctulata, killed by benzenehexachloride dust in Oklahoma, 100.

ciliatus, Dacus.

cilicrura, Hylemyia (Chortophila).

cilium, Spodoptera.

Cimex lectularius, water exchange between air and, 375.

Cinchona, pests of, in Porto Rico, 53.

cinctus, Cephus.

cinereus, Crypturgus. cingulata, Rhagoletis. cingulatus, Dysdercus.

Cinnamomum camphora, Chrysomphalus dictyospermi on, in Persia, 386.

circumflexus, Myzus.

Cirphis unipuncta, in Fiji, 390; outbreaks and natural enemies of, in Queensland, 307; in U.S.A., 96, 100; on cranberry, 100; on graminaceous crops, 96, 307, 390; measures and experiments against, 22, 96, 307.

cirsii, Trioxys.

citri, Aphis (see A. gossypii); Cercyonia; Dialeurodes (Aleurodes); Paratetranychus; Pseudococcus; Unaspis (Prontaspis).

citricidus, Aphis.

citrina, Xanthopimpla. citrinus, Aspidiotiphagus.

Citrullus vulgaris (see Watermelon).

Citrus, Aonidiella aurantii on, in S. Africa, 172; pests of, in Australia, 74, 108, 177, 181, 248, 249, 344, 345; Diaprepes esuriens on, in Bermuda, 49; Anychus orientalis on, in Egypt, 412; Aphis citricidus on, in Fiji, 413; not attacked by Quadraspidiotus perniciosus in Italy, 64; Papilio demodocus on, in Mauritius, 210; pests of, in Mexico, 79, 148; Pseudaonidia trilobitiformis on, in Nyasaland, 44; pests of, in Palestine, 409, 410, 412; Coccids on, in Persia, 386, 387; Coccids on, in Peru, 134; Pseudococcus filamentosus on, in Punjab, 123: Coccids on, in Spain, 196; Coccids on, in Trinidad, 319; pests of, in U.S.A., 13, 14, 79, 83, 89, 145, 155, 160, 181, 187, 188, 189, 221, 343; persistence of DDT deposits on, 89; insecticide treatments and injury to. 152, 400, 413; fumigation of transported fruits of, against pests, 79.

Citrus Bud Mite (see *Aceria sheldoni*). Citrus Red Mite (see *Paratetranychus*

citri).

Citrus Rust Mite (see *Phyllocoptruta olei-vorus*).

Cladius pectinicornis, bionomics and control of, in Belgium, 278, 279. claripalpis, Theresia (Paratheresia).

claripennis, Elasmus; Exenterus.

Clausenia purpurea (parasite of Pseudococcus spp.), in Japan, 410; bionomics and utilisation of, in Palestine and U.S.A., 410; parasites of, 410.

Clays, as carrier for dusts, 22, 112, 114, 329, 330, 402; types of, 329, 330; action of, on larvae of *Epilachna varivestis*, 330.

Clemora smithi, parasites and biological control of, on sugar-cane in Mauritius, 209, 210.

Cleonus punctiventris, insecticides against, on beet in Hungary, 63.

Clidemia hirta, utilisation of Liothrips urichi against, in Fiji, 390.

Cloanthanus heldoranus, on Parthenium argentatum in Mexico, 97.

Closterocerus, parasite of Diprion pini in Germany, 239.

Closterocerus insignis, parasite of Agromyza theae in Ceylon, 368. Clothes Moths, treatments protecting cloth, etc., against, 224, 253, 324, 395; resistance to, of fleeces from goats dipped in DDT, 146; rearing of, 343.

Clover, pests of, in Italy, 30, 115; Hypera meles on, in Ontario, 166; Nomophila noctuella on, in Rumania, 41; insects on, in U.S.A., 55, 136, 160, 187, 227, 228, 373, 395; effect of border plants on damage to, by insects, 374; experiment with Limonius californicus and, 91; insecticides and injury to, 228, 413.

Clover, Egyptian (see Trifolium alexandrinum).

Clover, Hubam (see Melilotus alba). Clover, Sour (see Melilotus indica).

Clover, Subterranean (see Trifolium subterraneum).

Clover, Sweet (see Melilotus).

Clover Mite (see Bryobia praetiosa).

Clover Weevil (see Sitona cylindricollis).

Clysiana (Clysia) ambiguella, sprays against, on vines in Switzerland, 296. Clytus floralis (see Plagionotus).

Cneorrhinus, on vines in Spain, 198.

coachella, Litoprosopus.

Coccidella spelaea, sp. n., on cacao in Gold Coast, 110.

Coccinella repanda var. transversalis (predacious on Aphids) in Fiji, 414; in New South Wales, 247.

Coccinella septempunctata, destroying eggs of Leptinotarsa decemlineata in Spain, 196.

Coccophagus caridei (parasite of Coccus viridis), introduction of, into Porto Rico from Brazil, 54.

Coccophagus fallax (parasite of Coccus viridis), introduction of, into Porto Rico from Brazil, 54.

Coccophagus heteropneusticus (see C. caridei).

Coccus acuminatus, fumigation against, on Gardenia imported into California from Mexico, 187.

Coccus africanus, relations of other insects to, on coffee in Kenya, 404.

Coccus discrepans, food-plants of, in Punjab, 123.

Coccus hesperidum, new parasite of, in S. Australia, 32; on Citrus in Persia, 387.

Coccus pseudomagnoliarum, on orange in Persia, 387.

Coccus viridis, natural enemies introduced from Brazil into Porto Rico against, 54; on gardenia, 54.

cockerelli, Dikraneura; Paratrioza.

Cockroaches, DDT dusts against, in buildings, 292, 364; use of, for insecticide tests, 10, 52, 175, 378; electrically-charged barrier for confining, 342.

Coconut, new Coccid intercepted in California from Mexico on, 401; pests of, in Fiji, 267, 390, 413; Aspidiotus destructor on, in Mauritius, 210; pests of, in New Hebrides and Solomon Is., 390.

cocophaga, Amblypelta.

Cooling Moth (see Cydia pomonella).

Coelophora inaequalis, predacious on Aphids in Fiji, 414.

Coelosterna scabrator (see Celosterna).

Coelosternus, list of species of, on cassava in Brazil and W. Indies, 6.

Coelosternus granicollis, bionomics of, on cassava in Brazil, 6.

Coenosia tigrina, predacious on Hylemyia spp. in England, 174.

coeruleas, Zicrona. coeruleus, Rhynchites. Coffea (see Coffee).

coffeae, Asterolecanium; Cerococcus; Diarthrothrips; Lygus.

coffearia, Homona.

Coffee, 110; estimation of infestation of, by Stephanoderes hampei in Brazil, 5; Antestia on, in Fr. Cameroons, 295; pests of, in Belgian Congo, 295, 316; pests of, in Kenya, 316, 317, 404; Stephanoderes on, in Porto Rico, 300; Coccid on, in Tanganyika, 316; pests of, in Uganda, 26, 367; book on pests of, 405.

Coffee Husks, in bait for locusts, 403. *Cola cordifolia*, new Coccid on, in Gold Coast, 110.

Colaphellus bowringi, tests of Millettia pachycarpa against, in Chile, 299.

Colaspidema atrum, sprays against, on lucerne in Spain, 203, 204, 293.

Cold Storage, of Aphelinus mali, 360; of potato tubers against Gnorimoschema operculella, 56.

Coleophora cerasivorella (fletcherella), in nests of Odynerus tigris in Ontario, 167. Coleophora fuscedinella, dormant sprays

against, in orchards on Holland, 276. Coleophora laricella, utilisation of parasites against, on larch in New Bruns-

wick, 208.

Coleophora salmani, on birch in New

Brunswick, 208.

Colias eurytheme, killed by benzenehexachloride dust on lucerne in Oklahoma, 100.

colibri, Athalia (see A. rosae).

Collards, pests of, in Louisiana, 159. collaris, Thyraeella.

Colocasia, pests of, in Pacific Is., 390,

Colombia, Sacadodes pyralis on cotton in, 126.

Colorado, Melanoplus spp. in, 237; orchard pests in, 79, 92, 185; potato pests in, 230; insects and virus diseases of plants in, 79, 93, 230, 231; Meteorus loxostegei in, 28.

Colorado 9, 237.

Colorado Potato Beetle (see Leptinotarsa decemlineata).

columbariella, Tinea. comes. Ervthroneura. commixtalis, Loxostege,

communis, Anuraphis (Dentatus) (see A.

devecta); Neolygus.

Comperiella bifasciata (parasite of Aonidiella aurantii), unlikely to become established in S. Australia, 216.

completa, Rhagoletis suavis. composita, Persectania,

compressus, Camponotus.

Compsolacon gracilis, on groundnuts in Solomon Is., 390.

comptana, Ancylis.

comstocki, Pseudococcus.

Comstockiella sabalis, parasites of, on Sabal bermudiana in Bermuda, 49.

concinna, Chaetocnema.

concolor, Aëdes.

conformis, Leis.

confusum, Tribolium.

Congo, Belgian, coffee pests in, 295, 316; Dacus spp. in, 176; pests of groundnuts in, 426; Schistocerca gregaria in, 207, 366; insects in timber in, 1, 295; Clerid predacious on Lyctus in, 1.

Coniopteryx tineiformis, predacious on Paratetranychus pilosus in Britain, 359; not affected by dormant sprays; 359.

conjugella, Argyresthia.

Connecticut, miscellaneous pests in, 36, 183, 301, 336; Pyrausta nubilalis on maize in, 401-403.

Conophthorus lambertianae, twig injury to pine by, in California, 301.

Conotrachelus, DDT against, on tomato in Peru, 134.

Conotrachelus nenuphar (in U.S.A.), on apple, 59, 100; reduced emergence of. from sun-scorched apples, 100; peach, 95, 127, 225, 373; causing deformation of peaches, 373; measures and experiments against, 59, 95, 100, 127, 225, 373.

conquisitor, Itoplectis. consocialis, Oechalia,

constrictus, Barbitistes; Desmoris.

Contarinia nasturtii (torquens), timing of control of, on cabbage in Holland, 285, 286,

contractus, Thylodrias. convergens, Hippodamia.

Convolvulus arvensis, Aphid Bulgaria, 5.

cooki, Cremastus.

cooleyi, Chermes.

Copa occidentalis, bionomics of, on cucurbits in Ivory Coast, 68.

Copidosoma koehleri, bionomics, introduction and value of, against Gnorimoschema operculella in Australia, 216.

Copper Arsenate, Basic, tests with, as an insecticide, 371, 372; and DDT, 371,

372; and rotenone, 371.

Copper Carbonate, in mixed dusts and sprays, 282, 362, 363; effectiveness of DDT, 363. reducing

Copper Dimethyldithiocarbamate, repellency of wood treated with, to Kalo-

termes brevis, 233.

Copper Fluoborate, repellency of wood treated with, to Kalotermes brevis, 232.

Copper Fluoride, repellency of wood treated with, to Kalotermes brevis, 232. Copper Fluosilicate, repellency of wood

treated with, to Kalotermes brevis, 232. Copper Naphthenate, wood treated with, against Coptotermes formosanus, 249.

Copper Oxide, reducing effect on plants of calcium arsenate in soil, 371. (See Cuprous Oxide.)

Copper Oxychloride, uses and effects of, in insecticide sprays and dusts, 112,

273, 362, 363.

Copper Pentachlorphenate, repellency of wood treated with, to Kalotermes brevis, 233.

Copper Phosphates, repellency of wood treated with, to Kalotermes brevis, 232.

Copper Sebacate, in spray formulae, 131, 132.

Copper Sulphate, in dust against Epitrix cucumeris on potato, 332; and oil emulsion, against Quadraspidiotus perniciosus and sooty mould, 293; Celcure (q.v.) containing, 414; treatment of wood with, against termites, 169, 232. (See Bordeaux Mixture.)

Copra, Caloglyphus berlesei infesting, 286. Coptosoma punctatissima (Bean Pentatomid), tests of Millettia pachycarpa against, in China, 299.

Coptotermes formosanus, in building timber in Hawaii, 249; copper naphthenate protecting wood from, 249.

Coranus pallidus, predacious on Lygaeids in Senegal, 44.

Corcyra cephalonica (in stored products), in Mauritius, 210; in Queensland, 309, 310; in Senegal, 44; in Uganda, 367; parasite of, 44; measures and experiments against, 28, 44, 210.

Cordia macrostachya, investigations relating to proposed introduction of insects into Mauritius from Trinidad

against, 424, 425. cordiae, Schematiza.

Cordylomera nitidipennis, bionomics and control of, in Khaya senegalensis in Senegal, 250.

core, Euploea.

Coreopsis spp., Homoeosoma electellum on, in U.S.A., 84.

coriarius, Prionus. corinna, Euploea core.

Cork Oak (see Quercus ilex var. suber).

Corn Borer, European (see *Pyrausta nubilalis*).

Corn Earworm (see *Heliothis armigera*). Corn Rootworm, Southern (see *Diabrotica duodecimpunctata*).

corni, Eulecanium (Lecanium).

cornuta, Tachardiella.

cornutus, Gnathocerus; Vasates.

corollae, Syrphus.

Corrosive Sublimate (see Mercuric Chloride).

corsicus, Crypturgus (see C. cinereus).

coryli, Eulecanium (Lecanium).

Corylus, Eriophyid on, in N. America, 83; Polydrusus cervinus on, in Britain, 358.

Corymbites, effect of, on yields of oats in England, 381, 382.

Corymbites aeneus, in Hungary, 64.

Corymbites cupreus, in England, 381, 382. Coryna apicicornis, on leguminous crops in E. Africa, 367, 426.

Coryneum, 192.

Corythaica monacha, on egg-plant in Trinidad, 319.

Corythucha pergandei, test of benzene hexachloride against, in New Hampshire, 234.

Cosmia spp., on apple in Belgium, 277; bionomics of, 277.

Cosmopolites sordidus, failure to establish Plaesius javanus against, on banana in Uganda, 367.

Cossus cossus, measures against, on ash in France, 32.

costalis, Eutermes (Nasutitermes).

Cotton, Argyroploce leucotreta on, in Fr. W. Africa, 67; Dysdercus spp. on, in S. Africa, 27; pests of, in Argentina, 126, 142; Agallia albidula on, in Brazil, 77, 104; Earias cupreoviridis on, in China, 312; Sacadodes pyralis on, in

Colombia, 126; new mite on, in Egypt, 411; Aphis gossypii on, in Fiji, 413; Sacadodes pyralis on, in Br. Guiana, 126; pests of, in India, 120, 121, 310, 312; Dysdercus on, in Paraguay, 142; Acridids attacking, in Persia, 388; pests of, in Peru, 134; Earias insulana on, in Spain, 197; pests of, in Uganda, 26, 27, 366, 367; Tetranychus turkestani on, in U.S.S.R., 118; pests of, in U.S.A., 12, 13, 14, 156, 183, 184, 231, 370, 371, 401; Sacadodes pyralis on, in Venezuela, 126; pests of, in West Indies, 126, 300; list of insects and mites recorded on, 388; effect of insecticides on germination and growth of, 338; calcium cyanamide for defoliation of, against Anthonomus grandis, 401.

Cotton Boll Weevil (see Anthonomus

grandis).

Cotton Borer (see Eutinobothrus brasiliensis).

Cotton Fabric, insects damaging, 324. Cottonseed, as food for *Dysdercus*, 27; treatments of, against *Platyedra gossypiella*, 66, 105.

Cottonseed Cake or Husks, in baits, 4,

Cottonwood Leaf Beetle (see *Melasoma* lineatopunctata).

Cowpea Curculio (see Chalcodermus

Cowpeas, Chalcodermus aeneus on, in S. Carolina, 370; insects on, in Fiji, 266, 413; new Jassid transmitting leaf crinkle of, 266; effect of soil treatment with benzene hexachloride on germination and growth of, 13.

Cowpeas (Stored), experiments against

Bruchids in, 2, 3, 369.

Crambus hortuellus, dusts against, on cranberry in Massachusetts, 100.
Crambus malacellus, on rice in Mauritius,

210.

Cranberry, pests of, in U.S.A., 100, 137. Crapemyrtle (see *Lagerstroemia*).

crassifemur, Eulimneria (Limnerium). crassus, Cylindrocopturus.

crataegellum, Rhopalosiphum.

crataegi, Anuraphis; Aporia; Eriosoma. Crataegus, weevils on, in Britain, 314, 358; Aphids on, 66, 336.

Cremastus, parasite of Marasmia venilialis in Fiji, 267; tests of toxicity of DDT to, 258, 259.

Cremastus cooki, in Ohio, 258.

Cremastus epagoges, hosts of, in U.S.A., 84.

Cremastus forbesi, in Ohio, 258.

Cremnops vulgaris, parasite of Loxostege sticticalis in N. America, 28, 29; loxostegei Meteorus intrinsically superior to, 29.

Creosote, treatments of logs or timber with, against Coleoptera, 103, 107, 248; against termites, 317; and oils,

103, 107.

Crepidodera, DDT against, on willow in U.S.A., 143.

Cresols, insecticides produced by chlorin-

ation of, 275, 276.

Crioceris asparagi (Asparagus Beetle), benzene hexachloride against, in New Hampshire, 234.

Crocidolomia binotalis, new parasite of, in Bihar, 119; on cabbage in Fiji, 390.

crocidolomiae, Apanteles.

Crotalaria mucronata, Aphis gossypii on, in Fiji, 413.

Croton, Psallus seriatus reared on, 184. Croton capitatus, Mirids on, in Missouri, 156.

crouani, Graeffea.

cruciata, Austroicetes.

cruciferae, Phyllotreta.

cruciferarum, Bagrada.

Cryolite, against insects on fruit and nut trees, 19, 151, 345; not affecting populations of Tetranychus pacificus on apple, 149; against pests of leguminous plants, 21, 230, 330, 370, 395, 396; against Lepidoptera on solanaceous crops, 10, 87, 178, 372; against Harrisina brillians on vines, 185, 397; against Lepidoptera on other crops, 100, 126, 178, 331, 332; against grasshoppers, 186; ineffective against Popillia japonica, 101; toxicity of, to bees, 222; in baits for cutworms, 62, 96; dusting with, 10, 21, 87, 100, 126, 178, 186, 230, 330, 331, 332, 370, 395, 396, 397; carriers for, 87, 178, 395; effect of carriers on efficiency of, 330; spraying with, 19, 101, 149, 150, 151, 345, 372; adhesives for, 19; applied from aircraft, 396; Alorco form of, 21; formulae containing, 19, 87, 96, 126, 149, 150, 151, 178, 331, 345, 395; and DDT, 150, 372; and Lethane, 396; and oil emulsions, 19, 149, 151; and sulphur, 21, 126, 230, 395; effects of, on plants, 19, 186, 331.

Cryptoblabes gnidiella, on mango in Egypt, 409.

Cryptococcus fagi, on beech in New Brunswick, 208.

Cryptolaemus montrouzieri, use of, against Dactylopius opuntiae on spineless cactus in S. Africa, 389.

Cryptotermes (see Kalotermes).

Crypturgus, key to Palaearctic species of, 31.

Crypturgus apfelbecki (see C. cinereus).

Crypturgus barbeyi, probably a synonym of C. numidicus, 31.

Crypturgus cinereus, synonymy of, 31. Crypturgus corsicus (see C. cinereus).

Crypturgus cylindricollis (see C. pusillus). Crypturgus maulei (see C. pusillus).

Crypturgus numidicus, C. barbeyi probably a synonym of, 32.

Crypturgus parallelocollis (see C. pusillus

parallelocollis).

Crypturgus pusillus, in felled pine in Italy,

195; synonymy of, 31.

Crypturgus pusillus parallelocollis, status of, 31.

Cryptus inornatus, developing in Loxostege sticticalis after Meteorus loxostegei, 29.

Cryptus sexannulatus, introduced from Canada into New Zealand against Cydia pomonella, 244.

CS431, 403.

Ctenolepisma longicaudata, in buildings in Australia, 363, 364; measures against, 364.

Ctenolepisma urbana, in buildings in Australia, 364.

Cuba, parasites and attempted biological control of Homoeosoma electellum on sunflower in, 83, 84; pests of sugarcane in, 240; new parasite in mealybugs intercepted in U.S.A. on pineapples from, 315; failure to establish Lixophaga diatraeae in Br. Guiana from, 304; plant quarantine restrictions in, 105.

Cubé (Lonchocarpus), dusting with, 315, 319, 326, 327, 395, 420, 421; •oil-impregnated dust of, 326, 327; carriers for, 395, 420; spraying with, 17, 160, 326, 327, 334, 335; adhesives and spreaders for, 160, 327, 335; formulae containing, 160, 189, 319, 326, 327, 334, 335, 421; and DDT, 189; and oil emulsions, 17, 160, 189, 335; and thiocyanate, 327; and Velsicol AR-60, 17, 420, 421; rotenone contents of, 17, 160, 315, 326, 334, 395, 421; bibliography of pests of stored roots of, 272. cucumarius, Dacus bivittatus.

Cucumber, Dacus bivittatus on, in W. Africa, 176; pests of, in Britain, 173, 284, 288; Aphis gossypii on, in Bulgaria, 4; Diaphania spp. on, in Trinidad, 319; pests of, in U.S.A., 94, 101, 225, 331; unfavourable food for Limonius californicus, 91; insecticides and injury to, 94, 224, 225, 331; benzene hexachloride not affecting flavour of, 234; DDT mixed with stored seed of, against insects, 19.

Cucumber Beetle, Spotted (see *Diabrotica duodecimpunctata*).

Cucumber Beetle, Striped (see *Diabrotica melanocephala*).

cucumeris, Epitrix.

Cucumis, Dacus bivittatus on, in W. Africa, 176.

Cucumis melo (see Melon).

Cucurbit Beetle, Black (see Aulacophora cattigarensis).

Cucurbita, susceptibility of species of, to Melittia cucurbitae, 101. (See Pumpkin and Squash.)

cucurbitacearum, Tetranychus (Eotetranychus).

cucurbitae, Melittia.

Cultural Practices, influence of, on insect pests, 3, 4, 41, 45, 115, 151.

cunea, Hyphantria. cunicularius, Hylastes. cupreoviridis, Earias.

cupreus, Corymbites.

Cupric Compounds (see under Copper). Cuprous Oxide, Yellow, and calcium arsenate, 165; compatible with DDT, 165, 166.

Curculio caryae, bionomics and control of, on pecan in Texas, 151.

Curculio, Cowpea (see Chalcodermus aeneus).

Curculio, Plum (see Conotrachelus nenuphar).

curculionis, Bathyplectes.

Currant, Cecidophyes ribis on, in Br. Columbia, 83; Incurvaria capitella on, in Holland. 285; soil treatment protecting, against Melolontha in Switzerland, 280; distribution of Rhopalosiphoninus ribesina on, 408.

Currant, Black, *Dasyneura tetensi* on, in Britain, **284**; Aphid on, in Utah, **388**. *currei*, *Plectops*.

custator, Thyanta.

Cut Flowers, legislation against introduction of pests and diseases with, into U.S.A., Hawaii and Porto Rico, 71, 209.

cuthbertsoni, Hyalomya (see Alophora nasalis).

Cutworm, Sunflower Head (see *Stibadium* spumosum).

Cutworms, baits for, 53, 62, 96, 307; other measures against, 96, 307, 308.

Cyamopsis psoraloides, 310.

cyaniceps, Eupelmus. cyanocephala, Lebia.

Cyanogas, 304.

Cybocephalus, predacious on Nilotaspis halli in California, 156; predacious on Quadraspidiotus perniciosus in Italy, 65; feeding habits and distribution of, 155.

Cybocephalus californicus, in U.S.A., 155, 156; possibly identical with C. nigritulus, 156.

Cybocephalus nigritulus, in U.S.A., 155; C. californicus possibly identical with, 156.

Cyclohexanol, thiocyanoacetic ester of, as an insecticide, 252, 253, 254.

Cyclohexanone, as a solvent for DDT, 87, 325.

Cyclohexylamine Dodecyl Sulphate, as an emulsifier, 131, 263.

Cydia leguminis, DDT ineffective against, on soy beans in Peru, 134.

Cydia molesta (Oriental Peach Moth), in France, 238, 239; in Ontario, 167; in U.S.A., 23, 56, 127, 152, 184, 185, 186, 225, 341, 370, 397; in Victoria, 178; bionomics and miscellaneous food-plants of, 152, 153, 238, 239; parasites and biological control of, 23, 56, 127, 178, 185, 370, 397; sprays against, 56, 179, 238, 341; effect of DDT on parasites of, 56; other measures against, 186.

Cydia nigricana, bionomics and control of, on peas in England and Sweden, 261–264, 409.

Cydia pomonella, in Australia, 106, 179, 218, 245, 246, 247; eradication of outbreaks of, in Western Australia, 106; in England, 132; in Italy, 194; intercepted in Kenya from S. Africa, 405; in New Zealand, 244; in Nova Scotia, 271, 392; in Switzerland, 296; in U.S.A., 12, 13, 19, 57, 80, 81, 92, 100, 126, 127, 128, 150, 155, 219, 225, 256, 257, 258, 322, 338, 372, 399; on peach and plum, 399; on pear, 80, 179, 194, 245; on walnut, 80, 155, 194, 399; question of walnut race of, 80; market losses caused by, 128; natural enemies and biological control of, 219, 220, 244; effect of sprays on parasites of, 219; experiments with Beauveria bassiana against, 220; bait for timing flights of, 132; 256, 257, 258, 296, 322, 323, 338, 372; strain of, resistant to lead arsenate, 323; laboratory tests of insecticides against, 12, 24, 93; experimental use

of, in tests of spray programmes, 271; other measures against, 322, 323; use of, for rearing *Meteorus loxostegei*, 29.

Cylas formicarius (on sweet potato), in Fiji, 390; insecticides against, in Louisiana, 12.

cylindricollis, Crypturgus (see C. pusillus); Sitona.

Cylindrocopturus crassus, in chrysanthemum in California, 188.

Cyllene robiniae (see Megacyllene).

Cymbidium (see Orchids).

Cynara cardunculus (see Cardoon).

Cynara scolymus (see Artichoke).

Cynodon dactylon, Läphygma spp. on, in E. Africa, 426.

Cyphomandra, Aphid on, in greenhouses in Britain, 361.

Cypress, repellence of chlorinated gum of, to *Kalotermes brevis*, 232.

Cytox, 352.

Czechoslovakia, insects and virus diseases of beet in, 355; miscellaneous pests in, 65, 278, 288.

D

D-4, 101, 102.

Dacnusa gracilis, parasite of Psila rosae in Britain, 417.

Dacrydium cupressinum, used in tests of preservatives against insects, 414.

Dactylopius opuntiae (in S. Africa), use and value of, against prickly-pear, 171, 172, 389; control of, damaging spineless cactus, 389; bionomics of, 389; natural enemies of, 172, 389.

Dacus, introduced parasites of, in Fiji, 266.

Dacus armatus, identity, subspecies and distribution of, in Africa, 176.

Dacus armatus lulongaensis, status and distribution of, in Africa, 176.

Dacus bipartitus (see D. bivittatus).

Dacus bivittatus, bionomics, synonymy and subspecies of, in Africa, 176, 177; parasite of, 177.

Dacus bivittatus cucumarius, synonymy, host fruits and distribution of, in Africa, 176; parasite of, 177.

Dacus bivittatus cucumarius form atrescens, n., 176.

Dacus brevistylus (see D. ciliatus).

Dacus ciliatus, causing fall of Citrus fruits in Eritrea, 108; cucurbits suggested as trap-crop for, 108.

Dacus ferrugineus dorsalis, host-fruits of, in Hawaii, 249; probably introduced from Saipan, 249.

Dacus ferrugineus tryoni, bait-spray for, in New South Wales, 248.

Dacus oleae, measures against, on olive in Spain and Greece, 201, 292.

Dacus pectoralis, Wlk. (African Species) (see D. bivittatus cucumarius).

Dacus pectoralis, Wlk. (Oriental Species), 176.

Dahlbominus fuscipennis, parasite and hyperparasite of Diprion pini in Germany, 239; utilisation of, against Gilpinia hercyniae in Quebec, 320.

Dahlia, pests of, in U.S.A., 335.

Dakota, North, pests of cereals in, 128; *Hylemyia antiqua* in, 388.

Dalbergia sissoo, Coccus discrepans on, in Punjab, 123.

dalmanni, Habrolepis.

damryi, Apion.

Damson, Scolytus rugulosus on, in Britain, 358.

dardanus, Papilio.

Dasus simplex, plants attacked by, in E. Africa, 367, 403.

Dasyneura tetensi, sprays against, on black currant in Britain, 284.

Datana angusi, on blueberry in New Jersey, 137.

Dates (see Fruit, Dried). Datura stramonium, 7.

D-D Mixture (dichlorpropene and dichlorpropane), soil treatment with, against wireworms and *Otiorrhynchus*, 187.

DDA, 162.

DDD [1, 1-dichlor-2, 2-bis (parachlor-phenyl) ethane], against Erythroneura comes, 325, 326; against Lepidoptera, 13, 403; against other insects, 14; tests of toxicity of, to Diptera, 275, 421; dusting with, 14, 403; in sprays, 13, 325, 326; derivatives of, 275.

DDT, against Aleurodids, 292; against Aphids, 14, 74, 92, 98, 112, 123, 131, 134, 143, 153, 155, 192, 218, 229, 247, 251, 292, 325, 326–328, 331, 332, 352, 371, 421; favouring infestation by Eriosoma lanigerum, 98, 296; against Cercopids, 143, 220; against Coccids, 17, 74, 88, 89, 101, 134, 143, 145, 187, 189, 194, 284, 292, 342; favouring infestation by Coccids, 155; against Coreids, 108, 370; against Jassids, 47, 74, 81, 100, 123, 127, 133, 134, 143, 147, 165, 219, 225, 246, 325, 326, 328, 344, 350, 351, 363, 386, 398, 421; against Lygaeids, 22, 74, 154; against Mirids, 11, 12, 90, 91, 94, 184, 235, 250, 272, 284, 394, 395; against Pentatomids, 123, 235, 248, 249, 295; against Psyllids, 230, 231, 335; against Tingids, 75, 123; against other Hemiptera, 134, 143,

332; against ants, 74, 218, 292, 404; 352; against ants, 74, 218, 292, 404; against cockroaches, 292, 364; against Coleoptera, 19, 22, 28, 35, 38, 39, 59, 62, 63, 73, 81, 98, 100, 101, 102, 103, 104, 123, 133, 134, 138, 139, 140, 143, 144, 146, 148, 165, 187, 196, 198, 199, 203, 204, 209, 210, 217, 225, 229, 230, 232, 266, 268, 269, 276, 278, 279, 281, 282, 284, 286, 291, 292, 293, 296, 297, 315, 324, 328, 331, 332, 333, 336, 337, 338, 339, 340, 345, 346, 352, 358, 359 338, 339, 340, 345, 346, 352, 358, 359, 369, 370, 371, 372, 380, 398, 408, 421; ineffective against Collembola, 187, 305; against Cydia pomonella, 13, 57, 58, 59, 81, 92, 100, 126, 127, 150, 155, 179, 180, 218, 225, 245, 246, 247, 256, 257, 258, 296, 323, 339; against other Lepidoptera, 10, 13, 19, 21, 22, 28, 33, 34, 35, 36, 46, 56, 60, 61, 69, 74, 81, 324, 331, 332, 333, 336, 341, 344, 346, 347, 352, 363, 369, 370, 371, 372, 388, 402, 403, 409, 421; increasing infestation by Eulia velutinana, 323; against Diptera, 46, 47, 54, 59, 73, 90, 110, 134, 143, 187, 210, 223, 270, 283, 284, 286, 291, 292, 324, 328, 350, 353, 377, 388; against Forficula auricularia, 187, 399; against Gryllids, 93, 94, 372; 132, 149, 150, 155, 180, 218, 225, 246, 249, 257, 282, 296, 341; mite populations little affected by, 59, 147, 276, 359; against sawflies, 279, 286, 296; against termites, 169, 233, 324, 404, 405; against thrips, 12, 102, 126, 232, 278, 292, 306, 307, 331, 338; against woodlice, 159, 292; effect of, on insect parasites and predators, 56, 91, 98, 134, 145, 147, 150, 225, 258, 259, 285, 323, 402; risks of, to bees and other pollinating insects, 12, 91, 100, 136, 222, 237, 336; mode and mechanism of action of, on insects, 174, 175, 215, 274, 277, 339; method of testing, on Drosophila, 164; effect of temperature and humidity on toxicity of, 259, 278, 378, 379, 380; relation of crystals to toxicity of suspentions of, 376, 377; repellent action of, 88, 143, 340; comparison of action of other compounds and, on insects, 22, 274, 275,

277, 278, 295, 421; duration of effectiveness of, on plants, 11, 34, 35, 36, 63, 88, 89, 133, 134, 148, 154, 159, 189, 226, 251, 256, 324, 326, 340, 394, 404; duration of effectiveness of, on other surfaces, 28, 39, 233, 337; and injury to plants, 34, 36, 60, 69, 94, 96, 100, 101, 102, 133, 147, 149, 179, 180, 225, 230, 263, 267, 323, 326, 331, 350, 352, 370; not impairing germination of seeds, 19, 138, 148, 270, 282, 338; decreasing germination of seeds, 331; residues of, on fruits, vegetables, etc., 57, 58, 73, 81, 82, 96, 134, 135, 136, 153, 162, 209, 257, 323, 339; removal of residues of, from apples, 135, 136; question of risks and toxicity of, to man and other mammals, 13, 39, 73, 81, 82, 91, 135, 153, 158, 161, 162, 272, 285, 339, 378, 379, 393, 396; in milk and fat of animals, 153, 162, 379, 393; excretion of, by man and rabbits, 162 forest fauna little affected by, 34 effect of, on fish and other aquatic fauna, 347; comparison of toxicity of, to insects and vertebrates, 174, 175; in aerosols, 87, 88, 92, 95, 147, 161, 162, 230, 292, 325, 326, 327, 331, 336, 337; distribution of, from fog generators, 17, 18, 87, 88, 147; in baits, 158, 333, 350, 353; in bands and barriers, 74, 154, 332, 402; in dusts, 10, 11, 12, 21, 22, 35, 38, 39, 46, 47, 57, 58, 59, 62, 63, 69, 74, 75, 81, 87, 91, 94, 96, 100, 101, 102, 109, 123, 126, 133, 134, 138, 140, 143, 145, 153, 154, 158, 159, 161, 162, 165, 178, 184, 187, 204, 209, 210, 217, 218, 223, 225, 226, 227, 229, 230, 231, 232, 237, 244, 250, 251, 266, 268 231, 232, 237, 244, 250, 251, 266, 268, 269, 272, 278, 279, 285, 286, 291, 292, 294, 295, 305, 306, 307, 308, 309, 324, 326, 327, 328, 331, 332, 336, 337, 338, 342, 344, 346, 347, 350, 351, 358, 359, 363, 364, 370, 371, 372, 377, 379, 386, 394, 395, 398, 399, 402, 403, 408, 421 carriers for, 12, 19, 22, 47, 57, 69, 87, 91, 94, 96, 123, 126, 138, 140, 143, 145, 158, 178, 187, 210, 217, 218, 229, 237, 244, 266, 324, 326, 332, 337, 338, 371, 379, 420; oil-impregnated dusts of, 57, 58, 229, 326, 370; effect of other solvents on efficiency of dusts of, 326, 327, 420, 421; equipment for wetting dusts containing, 57; in sprays, 13, 22, 23, 28, 33, 34, 35, 36, 39, 46, 47, 54, 57, 58, 59, 60, 61, 63, 69, 73, 74, 81, 88, 89, 92, 96, 98, 100, 101, 102, 103, 104, 108, 109, 112, 126, 127, 133, 134, 143, 144, 145, 155, 157, 158, 159, 161, 162, 165, 178, 179, 187, 189, 198, 203, 208, 210, 217, 218, 219, 220, 223, 225, 226,

230, 235, 245, 246, 247, 248, 249, 256, 257, 258, 259, 263, 264, 267, 268, 269, 272, 276, 278, 279, 281, 284, 285, 286, 291, 292, 293, 294, 296, 297, 306, 307, 308, 309, 323, 324, 325, 326, 327, 331, 332, 335, 336, 337, 338, 339, 340, 341, 344, 345, 350, 351, 352, 359, 363, 371, 386, 394, 404, 400, a tamised sprayer 386, 394, 404, 409; atomised sprays of, 34, 35, 36, 37, 147, 158, 327, 336, 398; application of, from aircraft, 35, 90, 143, 292, 327; treatment of maize silks or ears with, 13, 158; seed treatments with, 90, 148, 210, 270, 283, 331, 350; soil treatments with, 22, 73, 90, 93, 94, 138, 148, 149, 187, 291, 338, 353, 388, 398; not affecting microbial processes in soil, 22; uses of, against pests of stored products, 12, 19, 28, 38, 39, 46, 87, 139, 140, 209, 217, 249, 324, 369; application of, to textiles, 323, 324; resistance to insects of fleeces from goats dipped in, 146; in wall washes, 139; suggested use of, at airports, international 233 ; with acetone, 95, 131, 164, 226, 230, 325; with alcohol, 28, 140, 164; with aluminium stearate, 88, 89; APS 202, 337; with arsenates, 58, 59, 98, 127, 218, 225, 232, 256, 257, 371, 372; with benzene, 131, 328, 339; with benzene hexachloride, 218, 264; with Bordeaux mixture, 102, 127, 165, 257, 325; with other copper compounds, 131, 165, 166, 225, 332, 363, 371, 372; with organic fungicides, 131, 165, 350; with carbitol, 131; with cryolite, 372; with cyclohexanone, 87, 325; with dichlordifluoromethane, 325, 327, 337; with diethyl phthalate, 146; with dinitro compounds, 112, 150, 218, 276, 284; with fish glue, 235; with hydroxypentamethylflavan, 13, 341; with isophorone, 230; with lime, 139, 165; reaction of lime with, 139, 140; with lime-sulphur, 46, 47, 359; with methyl chloride, 95, 230, 325, 327; with methylated naphthalene (see also with Velsicol preparations), 33; with solvent naphtha, 28, 46, 47, 69, 73, 178, 217, 218, 219, 263; with nicotine, 57, 58, 98, 127, 218, 306, 371, 395; with mineral oils, 17, 22, 34, 35, 36, 46, 47, 58, 81, 96, 98, 103, 126, 127, 131, 150, 155, 158, 187, 189, 225, 230, 256, 257, 258, 276, 284, 285, 293, 309, 323, 325, 333, 336, 337, 342, 380, 404; with vegetable oils, 54, 139, 396; with oil of Eucalyptus dives, 218; with soluble pine oil, 146; with orthodichlorbenzene, 337; with pyrethrum, 157, 336; with coumarone resin, 131;

with rotenone insecticides, 154, 157, 189, 218, 229, 230, 371, 395; with sulphur, 47, 57, 58, 91, 180, 187, 225, 229, 230, 231, 232, 266, 324, 338, 342, 363; fused dust of sulphur and, 91; with tetralin, 89; with thiocyanates, 218, 284, 327, 395; with tobacco residues in spray, 58; with toluene, 139, 218; with Velsicol preparations, 89, 95, 165, 292, 325, 420, 421; with xanthone, 58, 98; with xylene, 34, 144, 158, 218, 220, 267, 333, 336, 340, 342; with zinc sulphate, 165; wetting and spreading agents for, 28, 57, 58, 131, 226; powders used with, in sprays, 57, 58, 98, 143, 150, 235, 246, 328, 371; effect of particle size of spray powders containing, 328; emulsifiers solutions of, 28, 46, 54, 69, 89, 131, 139, 146, 165, 189, 263, 337, 339, 340; colorimetric determination of, 168, 393; book on, 9; surveys of data on, 208, 297; proprietary preparations of, 58, 60, 61, 62, 63, 101, 102, 103, 104, 112, 133, 134, 158, 179, 204, 217, 218, 237, 239, 258, 268, 269, 276, 324, 325, 326, 346, 347, 352; toxicity of isomers of, to insects, 275, 421.

DDT Analogues, and related compounds, tests of, as insecticides, 274, 275, 276, 421; of no value against *Nymphula nymphaeata*, 346. (See Bromo-DDT, DDD, Fluoro-DDT and Methoxy-DDT.)

dearnessi, Rhizaspidiotus (Targionia).

Decahydronaphthalene, as solvent for benzene hexachloride, 413.

decemlineata, Leptinotarsa.

decora, Stenoma.

decorus, Oedaleus.

Decticus albifrons, damaging cereals in Persia, 388; predacious on Leptinotarsa decemlineata in Spain, 196.

Decticus verrucivorus, damaging cereals in Persia, 388.

Deenate 25W, 58, 258, 325, 326.

Deenate 50P, 324.

Deenate 50W, 324.

defoliaria, Erannis.

Delaware, orchard pests in, 225, 257, 321; vegetable pests in, 225.

Delhi, parasite in Lepidoptera in, 119.

delicatus, Macrocentrus.

Delonix regia (see Poinciana).

Demerara Sugar-cane Froghopper (see *Tomaspis flavilatera*).

demodocus, Papilio.

Dendrolimus pini, identification of excreta of, or pine in Germany, 75, 76.

Dendrosoter protuberans, parasite of Myelophilus piniperda rubripennis in Italy, 195.

Dendrotettix quercus, outbreak of, on oaks in New Jersey, 301, 302; parasite of, 302.

Dendrothrips bispinosus, bionomics of, on tea in Travancore, 251.

Dentatus communis (see Anuraphis devecta).

dentipes, Alcidodes.

deplana, Eilema (Lithosia).

Dermestes, in dried meat in Queensland, 310.

Dermestes maculatus (vulpinus), infesting salt fish in Uganda, 367; tests with benzene hexachloride against, 235.

Derris, against Aphids, 247, 282; against Coccids, 16, 17; against other Hemiptera, 75, 295; against Coleoptera, 14, 52, 315, 345; against Lepidoptera, 52, 102, 112, 249, 282, 402; against Locusta migratoria, 237, 238; against mites, 282; against sawflies, 112, 286; against thrips, 53, 102, 282; predacious insects possibly destroyed by, 402; toxicity of, to bees, 222; in aerosol, 230; dusting with, 52, 53, 112, 113, 237, 282, 295, 315, 402; carriers for, 52, 112, 113, 237; effect of carriers on efficiency of 112, 113; spraying with, 102, 247, 286; stored potatoes treated with, 249; formulae containing 102, 113, 230, 237, 247; and copper compound, 282; with mutual solvent in oils, 16, 17; and soap, 247; and sulphur, 282; rotenone contents of, 10, 16, 17, 52, 53, 102, 112, 249, 286, 295, 315, 402; other insecticides compared with, 10, 14, 238, 295, 402; effect of storage on, 51, 238; bibliography of pests of stored roots of, 272.

Derris fordii, insecticidal properties of, 299.

Desert Locust (see Schistocerca gregaria). desertus, Gryllulus (Gryllus).

Desiantha caudata, on peas in New South Wales, 73.

Desmeocraera varia, bionomics of, on guava in Ivory Coast, 389, 390; on Mimusopa spp. in S. Africa, 389.

Desmoris constrictus, bionomics and control of, on sunflower in U.S.A., 150, 151.

Desmoris fulvus, bionomics and control of, on Helianthus spp. in U.S.A., 150, 151.

destructor, Aspidiotus; Ceroplastes; Glycyphagus (Lepidoglyphus); Halotydeus; Mayetiola.

detectus, Iridomyrmex.

devastans, Empoasca.

devastator, Melanoplus mexicanus.

devecta, Anuraphis (Sappaphis).

Diabrotica, sprays against, on tor

Diabrotica, sprays against, on tomato in Peru, 133.

Diabrotica bivittata, tests of mamey-seed extracts against, in Porto Rico, 51.

Diabrotica duodecimpunctata, intercepted in Bermuda, 49; in U.S.A., 148, 225, 331, 337, 338; on cucurbits, 225, 331; bionomics of, on groundnuts, 337, 338; on maize, 148; measures and experiments against, 9, 148, 225, 331, 338.

Diabrotica melanocephala (vittata), tests of insecticides against, on cucurbits in U.S.A., 12, 101, 154, 225, 234, 331.

diademata, Phyllotreta.

Diaeretus plesiorapae, parasite of Aphids in Uruguay, 50.

Dialeurodes citri, elimination of, in California, 189.

diaphana, Trybliographa.

Diaphania, insecticides against, on cucurbits in Peru, 134.

Diaphania hyalinata, on cucumber in Trinidad, 319; tests of insecticides against, 22, 23, 51, 52, 319.

Diaphania nitidalis, insecticides against, on cucurbits in Trinidad and Virginia, 319, 331.

Diaprepes esuriens, on Citrus in Bermuda, 49.

Diarthronomyia chrysanthemi, experiments with sodium selenate against, on chrysanthemum in Illinois, 291.

Diarthrothrips coffeae, spray against, on coffee in Kenya, 404.

Diaspidiotus ancylus, on fruit trees, etc., in California, 399, 400.

Diaspis visci, in Belgium, 49; on Juniperus bermudiana in Bermuda, 48, 49; possibly introduced into Bermuda from California, 48; parasites of, 49.

Diathetes pandanae, on Pandanus tectorius in Fiji, 266.

Diatomaceous Earth, effect of particle size of, in DDT suspensions, 328; as carrier for dusts, 87, 294.

Diatomite, as carrier for dusts, 329; action of, on larvae of *Epilachna varivestis*, 330.

Diatraea canella, parasites of, on sugarcane and Paspalum in Br. Guiana, 304; benzene hexachloride not increasing infestation by, 304.

Diatraea grandiosella, 157.

Diatraea impersonatella, on Paspalum in Br. Guiana, 304; rare on sugar-cane, 304.

Diatraea saccharalis (on sugar-cane), in Br. Guiana, 304; in Porto Rico, 53; in U.S.A., 13; on Paspalum, 304; parasites and biological control of, 53, effect of benzene hexachloride on, 13, 304.

diatraeae, Lixophaga; Paratheresia (see

Theresia claripalpis).

Diazoaminobenzene, tests of dust of, against Protoparce spp., 87.

Dibenzyl Pyrone (see Xanthone).

Dibrachys cavus, parasite of Apanteles glomeratus in Europe, 406, 407; reared from Lepidoptera in Quebec, 321: bionomics of, 406.

Dibromodiphenyltrichlorethane (see

Bromo-DDT).

Dicerca, on Parthenium argentatum in Mexico, 97.

Dicerca horni, damage to plums by, in California, 92.

Dichlordifluoromethane, use of, for producing aerosols, 325, 327, 337.

Dichlordiphenyldichlorethane (see

Dichlordiphenyltrichlorethane DDT).

Dichlorethane (see Ethylene Dichloride). Dichlorethyl Ether, against cheese mites, 245, 363; (emulsions of), against Hylemyia brassicae, 223; unsatisfactory for soil treatment against Conotrachelus nenuphar, 95.

1, 1-Dichlor-1-nitroethane, in fumigant mixtures against insects in stored grain, 85; addition of, to oils against

Parlatoria oleae, 17.

Di(4-chlorphenoxy)methane, against Paratetranychus citri, 155.

Dichlorpropane and Dichlorpropene (see D-D Mixture).

Dichocrocis punctiferalis, measures against, on guava and castor in Punjab, 123.

Dichomeris marginella, eradication of, on

juniper in California, 187.

Dichroplus spp., outbreak of, in Chile, 108, 109; bionomics and control of, 109.

dictyospermi, Chrysomphalus.

Dicyclohexylamine Salt, of Dinitro-ocyclohexylphenol (q.v.), 101, 102, 150, 155, 218, 222, 272, 288, 335, 372, 394. diemenalis, Nacoleia.

Diethyl Phthalate, in DDT dip formula, 146.

Diethylene Glycol Monobutyl Ether. not increasing toxicity of oil to Parlatoria oleae, 17.

Diethylene Glycol Monoethyl Ether, in DDT spray formula, 131.

Dieuches patruelis, natural enemies and control of, attacking harvested groundnuts in Senegal, 44.

Differential Grasshopper (see Melanoplus

differentialis).

differentialis, Melanoplus.

Difluorodiphenyltrichlorethane (see Fluoro-DDT).

cockerelli, Dikraneura experiments against, on vines in Arizona, 147.

Dimethoxydiphenyltrichlorethane (see Methoxy-DDT).

Dimethyldiphenyldichlorethane, 274, 275.

bis(Dimethylphenyl)dichlorethane, 275.

dimidiatus, Carpophilus.

Dinitro-o-cresol, against Anthonomus pomorum, 276, 296, 315; against Aphids, 111, 130, 247, 276, 285, 286, in barrier against Blissus leucopterus, 154; against Dasyneura tetensi, 284; against Lepidoptera, 276, 286; against locusts 99, 341, 342, and grasshoppers, 403; ineffective against Lygidea mendax, 272; against Paratetranychus pilosus, 101, 130, 276, 287, 288, 289; against Psyllids, 287; tests of ovicidal action of, 111, 273, 274, 276; dusting with, 315, 403; formulae for, in sprays, 99, 101, 289; applied from aircraft, 99; effect of pH on effectiveness of, 111, 112, 273, 274, 287; effect of temperature on toxicity of, 112, 285, 289, 380; timing of dormant sprays of, 112, 130; effects of, on plants, 99, 287, 315; and copper oxychloride, 112, 273; DDT, 112, 276, 284; and oil emulsions, 130, 272, 273, 274, 276, 284, 287; and tar distillates, 273, 274; solutions of, 99, 341, 342; salts of, 111, 272, 274, 285.

Dinitro-o-cyclohexylphenol, against Anthonomus spp., 278, 315; against Halticus bracteatus, 394; tests of, on Leptinotarsa decemlineata, 296; effective against Lygidea mendax, 272; against mites, 82, 101, 103, 150, 155, 218, 287, 288; ineffective against Psylla pyricola, 335; against Quadraspidiotus perniciosus, 296; against Thrips tabaci, 102; not harmful to Anthocoris nemorum, 288; toxicity of, to bees, 222; dusting with, 82, 101, to bees, 222; dusting with, 82, 101, 102, 278, 315, 394; effects of carriers on action of, 288; formulae for, in sprays, 103, 150, 278, 296, 355, 372, 394; and DDT, 150, 218; effect of, in lead-arsenate sprays, 101, 372, 373; and oil emulsions, 272, 296; pyrethrins, 394; and sulphur, 101, 394;

and injury to plants, 101, 102, 288; dicyclohexylamine salt of, 101, 102, 150, 155, 218, 222, 272, 288, 335, 372, 394.

Dinitrophenol, Tanalith (q.v.) containing, 414.

Dinoderus minutus, effect of age, species and treatment of bamboo culms on infestation by, in Porto Rico, 54, 55.

Dinoderus ocellaris, bionomics of, in stored bamboo in India, 122; digestive enzymes of larvae of, 119.

Dioctyl Sodium Sulphocuccinate, not increasing toxicity of oil to Parlatoria oleae, 17.

Dioryctria reniculella, parasites of, on spruce in Quebec, 321.

Dioscorea macroura, Trypetid on, in Tanganyika, 176.

1, 4-Dioxane, as a solvent for benzene hexachloride, 342.

diparopsidis, Apanteles.

Diphenyl Mercury, repellent to Kalotermes brevis, 233.

Diphenylamine, dusting with, against Hylemyia brassicae, 223.

Diphenyldichlorethane, derivatives of, as insecticides, 275.

Diphenyltrichlorethane, 421.

Diprion frutetorum (see Gilpinia).

Diprion pini (Pine Sawfly), parasites and hyperparasites of, in Germany, 239; experiments with dusts against, in Holland, 112, 113, 270; characters of excreta of, 76.

Diptilomiopus gigantorhynchus (prunorum), on blackberry and plum in N. America, 83.

Dirhinus, parasite of Dacus in Zanzibar. 177.

Dirhinus giffardi, introduced against Dacus in Fiji, 266.

dirhodum, Macrosiphum (Metopolophium). discedens, Lyctus.

Disodium Ethylene Bisdithiocarbamate. in DDT spray formula, 165.

Disonycha xanthomelaena (Spinach Fleabeetle), benzene hexachloride against, in New Hampshire, 234.

discrepans, Coccus.

dispar, Lymantria (Porthetria); Prosayleus; Xyleborus (Anisandrus).

disparis, Anastatus.

disstria, Malacosoma.

Dithane, 165.

 β , β' -Dithiocyanodiethyl Thiocyanates). Ether (see

divisus, Macrosteles.

DN-5, 394.

DN-111, 101, 155, 222, 372, 373, 394.

Dociostaurus kraussi, damaging crops in Persia, 387, 388.

Dociostaurus maroccanus, crops damaged by, in Persia, 387, 388; centres and phases of, in Spain, 202.

Dolomite, experiments with, as carrier for dusts, 112, 113, 114.

domestica, Musca; Thermobia.

domesticus, Glycyphagus. dominica, Rhizopertha.

Dominican Republic, new Pyralid on sugar-cane in, 29, 30.

Doralis (see Aphis).

dorsalis, Chalepus: Dacus ferrugineus. dorsocastaneus, Neopristomerus (Pristo-

merus) appalachianus.

Dreft, 135, 344.

Drepanosiphum platanoides, flights of, in England, 196.

Drosophila, DDT against, in canning factory in California, 187.

Drosophila melanogaster, tests of insecti-

cides on, 164, 421.
Dry-wood Termite, West Indian (see Kalotermes brevis).

dumerili, Epilachna.

duodecimpunctata, Diabrotica.

durantae, Aphis.

Dusts, equipment for applying, 11, 57, 303, 331, 364, 372, 402; application of, from aircraft, 17, 19, 228, 327, 328, 396, 401; effects of particle size on application of, 145, 393; factors affecting particle size of, 145, 146; electrostatic charges of, 393; effects and classification of diluents for, 329, 330; use of oil-112–114. impregnated, 58, 59, 145, 229, 326, 327, 370; sprays compared with, 63, 69, 165, 227, 363; (inert), against pests in stored products, 210, 217, 319; mode of action of, on insects, 329, 330; humidity reducing effectiveness of, against Lepisma saccharina, 379.

Dutch Elm Disease (see Ceratostomella ulmi).

DX, 157.

DX-nicotine, 100.

Dynaspidiotus britannicus, on Buxus in California, 187, 400.

Dynone, 218.

Dysdercus, distribution and importance of Tachinid parasites of, in Africa, 27, 28; bionomics of unidentified species of, on cotton in Argentina and Paraguay, 142.

Dysdercus andreae, tests of plant insecticides on, in Porto Rico, 52.

Dysdercus cardinalis, 366; parasite of, 27.

Dysdercus cingulatus, tests of Millettia pachycarpa on, in China, 298.

Dysdercus fasciatus (on cotton), parasite of, in S. Africa, 27; populations of, in Uganda, 366.

Dysdercus intermedius, parasite of, on cotton in S. Africa, 27.

Dysdercus megalopygus (see D. cingulatus). Dysdercus nigrofasciatus, parasites of, on cotton and Hibiscus calyphyllus in S. Africa, 27; populations of, of cotton in Uganda, 366.

Dysdercus pallidus, in Argentina, 142.

Dysdercus ruficollis, dusts against, on cotton in Peru, 134.

Dysdercus superstitiosus, populations of, on cotton in Uganda, 366; parasite of, 27.

E

Earias, on cotton in Uganda, 366, 367. Earias cupreoviridis, on cotton in China, 312; parasites and food-plants of, in Punjab, 120, 312.

Earias fabia, on cotton in Punjab, 120, 312; other food-plants of, 312;

parasites of, 120.

Earias insulana (on cotton), in Punjab, 120, 312; bionomics of, in Spain, 197; miscellaneous food-plants of, 312; parasites of, 120.

Earwig, European (see Forficula auricu-

laria).

ecdytolophae, Microgaster.

echinogalliperda, Amradiplosis. echinopus, Rhizoglyphus.

Economic Entomology, books on, 3, 355, 357, 427; bibliography of, in Croatia, 104.

Ectroma, parasite of Apanteles glomeratus in Europe, 407.

Ecuador, Schistocerca paranensis in, 133.

egeria, Apiomorpha. Egg Yolk, Dried, used for feeding insects, 343.

Egg-plant, new mite on, in Egypt, 411; Adoretus versutus on, in Fiji, 390; Empoasca lybica on, in Palestine, 175, 176; pests of, in Trinidad, 319.

Egypt, miscellaneous pests of mango in, 409, 411, 412; attempted introduction of Leis conformis into, 409, 410; digest of plant-quarantine restrictions in. 105.

Egyptian Clover (see Trifolium alexandrinum).

Eichhornia, Rhopalosiphum nymphaeae on, in Fiji, 413.

Eilema deplana, characters of excreta of, on conifers in Germany, 76.

Eire, pests of flax in, 208; summary of plant-quarantine restrictions in, 184.

Elasmopalpus lignosellus, dusts against, on soy beans in Peru, 134.

Elasmus claripennis, parasite of Eublemma

amabilis in Bihar, 120. Elasmus johnstoni, hosts of, in Punjab, 120; bionomics of, 120.

Elder (see Sambucus).

electellum, Homoeosoma.

Electricity, tests with, against insects in packaged cereals, 15; barrier charged with, for confining insects, 342.

Electrostatic Charges, of dusts, 393.

elegans, Zonocerus.

elegantalis, Leucinodes. elegantula, Erythroneura,

elegantulum, Apion.

Eleodes opaca, killed by benzene hexachloride dust in Oklahoma, 99.

Eleusine, Acridids attacking, in Uganda, 367.

elisus, Lygus.

Ellice Is., Papuana huebneri in, 390.

Elm, Cosmia spp. on, in Belgium, 277; pests of, in Canada, 163, 208, 320; pests of, in U.S.A., 14, 15, 103, 143, 144, 234, 320, 336; relation of Scolytids to Ceratostomella ulmi in, 163, 320; DDT solvents causing injury to, 144.

Elm Caterpillar, Spiny (see Nymphalis antiopa)

Elm Leaf Beetle (see Galerucella luteola), Elm Scale, European (see Gossyparia spuria).

elongatus, Dichroplus.

elutella, Ephestia.

Embia vayssierei, measures against, infesting shelled groundnuts in Senegal,

Emboloecia sauzalitae, in Lilium longiflorum in California, 187.

Empoasca, on Parthenium argentatum in Mexico, 97; dust against, on soy beans in Peru, 134; on cotton in Uganda, 366. Empoasca devastans, tests of DDT dust

against, in India, 123.

Empoasca fabae, in Ontario, 164, 165; in Peru, 133; in U.S.A., 71, 160, 225, 230, 234, 328, 335, 421; on dahlia, 335; on leguminous plants, 71, 160, 230; on potato, 164, 165, 225, 234, 328, 421; on tomato, 133; insecticides against, 133, 164, 165, 225, 230, 234, 328, 335, 421.

Empoasca fabalis, bean varieties resistant to, in Porto Rico, 32.

Empoasca flavescens, stated to transmit beet mosaic in Czechoslovakia, 355.

Empoasca lybica, bionomics of, in Palestine, 175, 176; insecticides ineffective against, 176; synonymy of, 175.

Empoasca maligna, benzene hexachloride ineffective against, on apple in U.S.A., 336.

terrae-reginae, insecticides Empoasca against, on tomato in Australia, 74,

Empusa, infesting Dactylopius opuntiae in S. Africa, 172.

Empusa aphidis (see Entomophthora).

Encarsia formosa, utilisation of, against Trialeurodes vaporariorum in greenhouses in Tasmania, 70.

Encoptolophus pallidus, on Parthenium argentatum in Mexico, 97.

Endothenia hebesana (see Argyroploce). Endrosis lactella, fumigation against, in bagged wheat in S. Australia, 182.

enigma, Oedaleonotus.

Entada abyssinica, new Mirids on, in Uganda, 26.

Entedon ovulorum, parasite of Diprion pini in Germany, 239. Entoleter, for destruction of insects in

grain and flour, 282.

Entomology, manual of equipment for studies on, 9; bibliography of, of Croatia, 104. (See Economic Entomology.)

entomophagus, Thyreophagus.

Entomophthora aphidis, infesting Toxoptera graminum in Uruguay, 50.

Entomoscelis adonidis, bionomics and control of, on rape in Rumania, 40. Eotetranychus (see Tetranychus).

epagoges, Cremastus.

ephemeraeformis, Thyridopteryx.

Ephestia cautella, in stored groundnuts in U.S.A. and Africa, 25, 44, 426; in other stored products in Queensland, 309, 310; natural enemies of, 44, 426; type of bag affecting infestation by, 25; measures against, 44; DDT deposits not toxic to larvae of, 28.

Ephestia elutella, investigations on, in stored wheat, etc., in England, 210-214, 264; processes affecting, in stored tobacco in U.S.A., 85, 86; effect of natural foods on development of, 264-266; natural enemies and disease of, 212; duration of effectiveness of pyrethrum films against, 168.

Ephestia kuehniella, in bagged wheat in Australia, 182; tests with frequency electric fields against, in packaged flour, 16; use of, for rearing Meteorus loxostegei, 29; tests of ovicides on, 273.

Ephialtes caudatus, introduced into New Zealand against Cydia pomonella, 244. Ephialtes ontario (see Apechthis).

Ephialtes pomorum, parasite of Anthonomus pomorum in Britain, 314.

ephippella, Argyresthia.

Epicaerus imbricatus, killed by benzenehexachloride dust, on lucerne Oklahoma, 100.

Epicauta spp., not killed by chlordane on potato in Illinois, 228.

Epigrimyia floridensis, hosts of, in U.S.A., 84.

Epilachna dumerili, tests of DDT dust

against, in India, 123.

Epilachna varivestis (on beans), in Canada, 167: measures and experiments against. in U.S.A., 12, 13, 14, 225, 229, 230, 234, 329, 330, 351, 370, 396, 421; on lima beans, 225, 370, 396.

Epilachna vigintioctopunctata, on cucurbits in Australia, 177; tests of DDT dust against, in India, 123.

Epimasicera caesar (see Zenillia).

Epiplagiops littoralis, in S. America, 97; introduction of, into California against Listroderes obliquus, 97.

Epistrophe (see Syrphus).

Epitrimerus gigantorhynchus (see Diptilomiopus).

Epitrimerus pyri (pirifoliae), bionomics and synonymy of, 82, 83.

Epitrimerus vitis (see Calepitrimerus).

Epitrix, sprays against, on tomato in Peru, 133; on egg-plant in Trinidad,

Epitrix cucumeris (on potato), in Ontario, 165; in U.S.A., 102, 234, 328, 332; insecticides against, 102, 165, 234, 328,

Epitrix hirtipennis, insecticides against. on tobacco in S. Carolina, 372,

in Br. Epitrix tuberis (on potato), Columbia, 167; insecticides against, in U.S.A., 230.

Epizeuxis aemula (see Camptylochila).

epos, Anagrus.

Erannis defoliaria, on fruit trees in Britain, 359.

Erannis leucophaearia, on apple and oak in Britain, 359.

Erannis tiliaria, disease of, on maple in Quebec, 319.

Ericolin, in flowers of Rhododendron molle, 299.

eridania, Laphygma (Prodenia).

Erigeron affinis, Heliopsis longipes (q.v.) recorded as, 12, 25.

Erigeron annuus, Hemiptera on, in Virginia, 334.

Erigeron canadensis, Hemiptera on, in Virginia, 334.

Eriobotrya japonica, Chrysomphalus dictyospermi on, in Persia, 386.

Eriococcus borealis, eradication of, on Liquidambar in California, 400.

Eriodendron anfractuosum (see pentandra).

Eriophyes, on apple in California, 400. Eriophyes carinatus, on tea in Travancore,

251.

Eriophyes essigi (see Aceria).

Eriophyes mangiferae, doubtful identity of, on mango in Egypt, 409.

Eriophyes pyri, biological forms of, on pear, 82; question of relationship of

E. p. mali to, 82.

Eriophyes pyri mali (Apple Blister Mite), question of relationship of, to E. pyri, 82.

Eriophyes quadrisetus typicus, on Juniperus

communis in Poland, 268.

Eriophyes vitis, on vines in U.S.A., 83. Eriosoma crataegi, benzene hexachloride

against, on Crataegus in U.S.A., 336. Eriosoma lanigerum (on apple), Australia, 70, 180, 247; in Britain, 359, 360; in Kirghizia, 118; in Switzerland, 296; in Uruguay, 50; Washington, 98; utilisation and value of Aphelinus mali against, 50, 70, 98, 247, 359, 360; effect of DDT on numbers of natural enemies of, 98, 180, 247, 296; effects of other sprays on, 98, 118.

pests of Citrus in, Eritrea, 108: Schistocerca gregaria in, 124; pests of vegetables in, 176, 382, 384; mass migrations of Lepidoptera across, 382.

erosus, Ips.

Erysiphe polygoni, dusts against insects and, infesting beans in Virginia, 229, 230.

erythrogaster, Campsomeris.

Erythroneura, on vines in U.S.A., 127, 147, 255, 256, 325; estimation of injury by, 255, 256; sprays and aerosols against, 127, 147, 325.

Erythroneura comes, on vines in U.S.A., 255, 325.

Erythroneura elegantula, on vines in U.S.A., 147.

Erythroneura leveri, sp. n., in Fiji, 266; transmitting leaf-crinkle of leguminous plants, 266.

Erythroneura variabilis, on vines Arizona, 147.

essigi, Aceria (Eriophyes).

esurians, Diaprepes.

Ethyl Acetate, for removing DDT deposits from apples, 135.

Ethyl Alcohol (see Alcohol). Ethyl Ether, as a solvent, 51.

Ethyl Formate, fumigation with, against Calandra oryzae, 217.

Ethylene Dibromide, in fumigant mixture against insects in stored wheat, 12.

Ethylene Dichloride, emulsion of, against Aegeria exitiosa, 321, 322; as fumigant against pests of stored cereals, 12, 85, 182, 217; other fumigants mixed with. 12, 85, 182, 217.

Ethylene Glycol, tests of dinitro-o-cresol

in, 380.

Ethylene Oxide, not a satisfactory fumigant against Quadraspidiotus perniciosus on nursery stock, 292.

Etiella zinckenella, on peas and Robinia

in Hungary, 63.

Eublemma amabilis, biological control of. predacious on Laccifer lacca in Bihar. 120.

Eucalyptus spp., Gonipterus scutellatus on, in E. Africa, 106, 367, 405; new Coccid on, in W. Australia, 69; Helopeltis theivora on, in Madras, 251; G. scutellatus on, in Mauritius. 125; differences in susceptibility of, to G. scutellatus, 106, 367; emulsified solution of DDT in oil of. 218.

Eucalyptus Weevil (see Gonipterus scutel-

eucerus, Habrocytus.

Euchalcidia, parasite of Pachymerus cassiae in Senegal, 44.

Eucosma, on chrysanthemum in California, 188; on Parthenium argentatum in Mexico, 97.

Eugenia cumini, new mite on, in Egypt, 411; Coccus discrepans on, in Punjab, 123.

Eugenia jambolana (see E. cumini).

Eugenia jambos, fruit-fly infesting, in Hawaii, 249.

Eugenia malaccensis, fruit-fly infesting, in Hawaii, 249.

Eugenia micheli (see E. uniflora),

Eugenia uniflora, Argyroploce leucotreta on, in Ivory Coast, 67.

Eulecanium corni, experiments against, on plum in Poland and Bulgaria, 354.

Eulecanium coryli, trees attacked by, in Czechoslovakia, 65; mechanism of injury to ash by, 65.

Eulecanium kunoense, on fruit trees, etc., in California, 399.

Eulecanium pruinosum, DDT spray increasing infestation of walnut by, in California, 155.

Eulecanium rugulosum, on fig and Pistacia in Persia, 386.

Eulia mariana, on apple in Nova Scotia,

Eulia velutinana, DDT increasing infestation of apple by, in Illinois, 323. Eulimneria crassifemur, parasite of Neurotoma nemoralis in Poland, 255.

euonymi, Unaspis.

Euonymus, Unaspis euonymi on, in California, 400.

Eupatorium pallescens, use of, to encourage Scoliids in Mauritius, 210.

Eupelmella vesicularis, parasite of Tortrix packardiana in Quebec, 321.

Eupelmus cyaniceps amicus, parasite of Bruchus floridae in Oklahoma, 9.

Euphorbia pulcherrima, insecticides causing injury to, 282.

euphorbiae, Aphthona.

Euphorocera omissa, developing in Loxostege sticticalis after Meteorus loxostegei, 29.

Eupista fuscedinella (see Coleophora). Euplectrus laphygmae, parasite of Laphygma exempta in Nyasaland, 45.

Euplectrus taiwanus, sp. n., hosts of, in Formosa, 177.

Euploea core corinna, food-plants of, in New South Wales, 344.

Euproctis, scope and status of, 26.

Euproctis chrysorrhoea, L., dusts against, in forests in Holland, 112; identity of, 26; type of genus, 26.

Euproctis chrysorrhoea, auct. (see E. similis).

Euproctis kargalike, on Pistacia in Persia, 386.

Euproctis lunata, new parasite of, in New Delhi, 119.

Euproctis similis, tests with DDT against, in Turkey, 352; identity of, 26.

Euproctis taiwana, new parasites of, in Formosa, 177.

euproctisiphagus, Apanteles.

Eupropacris, attacking groundnuts in Tanganyika, 426.

Eurygaster integriceps, parasites of, on cereals in Tadzhikistan, 241.

Euryischia, hosts and systematic position of, 109.

European Corn Borer (see *Pyrausta nubilalis*).

European Earwig (see Forficula auricularia).

European Elm Scale (see Gossyparia spuria).

European Red Mite (see Paratetranychus pilosus).

European Spruce Sawfly (see Gilpinia hercyniae).

Eurymyiocnema aphelinoides, gen. et sp. n., hyperparasite of *Pseudococcus* in Java and Uganda, 109.

euryptychiae, Pristomerus.

eurytheme, Colias.

Eurytoma, parasite of Asphondylia ricini in Hyderabad, 110.

Eurytoma syleptae, hyperparasite of Laphygma exempta in Nyasaland, 45.

Euscepes postfasciatus, on sweet potato in Fiji, 390.

Euschistus, measures against, causing deformation of peaches in U.S.A., 235, 334, 373; relation of weeds and cover crops to, 334, 373.

Euschistus servus, 235, 373. Euschistus tristigmus, 373. Euschistus variolarius, 373.

Eutermes, attacking sugar-cane in Br. Guiana, 304, 305; ant predactions on, 305; flooding against, 305.

Eutermes costalis, attacking sugar-cane in Br. Guiana, 304.

Eutetrapha tridentata, on elm infested with Ceratostomella ulmi in Quebec, 320.

Eutettix tenellus, food-plants and migrations of, transmitting curly-top of crops in California, 186, 397, 398; measures against, 398.

Eutinobothrus brasiliensis, sprays against, on cotton in Brazil, 104.

Euxoa segetum and ypsilon (see Agrotis). evanescens, Trichogramma.

ewingi, Persectania (see P. composita). exempta, Laphygma (Spodoptera).

Exenterus, utilisation of, against Gilpinia hercyniae in Quebec, 320.

Exenterus amictorius, utilisation of, against Gilpinia hercyniae in Quebec, 320-321.

Exenterus claripennis, value of, against Gilpinia hercyniae in New Brunswick, 208.

Exenterus vellicatus, value of, against Gilpinia hercyniae in New Brunswick, 208.

exigua, Laphygma.

exitiosa, Aegeria (Sanninoidea).

Exochomus melanocephalus, use of, against Dactylopius opuntiae on spineless cactus in S. Africa, 389.

Exochomus quadripustulatus, predacious on Anuraphis devecta in Holland, 66.

Eye-spotted Budmoth (see Spilonota ocellana).

F

fabae, Aphis (Doralis); Empoasca. fabalis, Empoasca. fabia, Earias. faceta, Antestia. fagi, Cryptococcus.

Fall Cankerworm (see Alsophila pometaria).

Fall Webworm (see Hyphantria cunea).

fallax, Coccophagus.

False Ipecacuanha (see Asclepias).

False Wireworm (see *Psammodes similis*). falsus, Protostrophus.

farinae, Tyroglyphus.

Farinococcus loranthi, sp. n., on Loranthus in Gold Coast, 110.

fasciata, Bogosiella; Prospaltella.

fasciatus. Dysdercus: Hercothrips: Oncopeltus.

Fatsia japonica, eradication of Pulvinaria pyriformis on, in California, 400.

fausta, Rhagoletis.

Felspar, micronised, mixed with grain against pests, 319.

femur-rubrum, Melanoplus.

fenestratus, Scymnus.

Fennel, Calocoris norvegicus on, in Italy, 383.

Fermate (see Ferric Dimethyldithiocarbamate).

Dimethyldithiocarbamate, fluence of sprays of, on Arthropods in apple orchards, 271, 392, 393; test of, against Kalotermes brevis, 233; use and compatibility of insecticides with, 131, 132, 165, 224, 350.

Ferric Oxide, reducing effect on plants of calcium arsenate in soil, 371; DDTimpregnated ore of, against Gnorimoschema operculella in potatoes, 217.

ferrisi, Tachardiella.

Ferrisia virgata, food-plants and relation of, to swollen-shoot of cacao in Gold Coast, 111; in U.S.A., 189, 315; on avocado, 189; ants associated with, 111; parasites of, 111, 315.

ferrugineus, Dacus (Strumeta).

ferus, Nabis.

Ficus, Euploea core corinna on, in New South Wales, 344.

Ficus carica (see Fig). ficus, Chrysomphalus.

Fig. pests of, in Australia, 108; new mite on, in Egypt, 411; Coccid on, in Persia, 386; Coccids on, in Punjab, 123; Lamiid on, in Tahiti, 250; pests of, in U.S.A., 14, 189; thiocyanoacetates on, 253. effect of

Figs, Dried (see Fruit, Dried).

Fiji, injurious Aphids of, 413; miscellaneous pests in, 266, 267, 390; insects and virus diseases in, 46, 266, 413; natural enemies and biological control of insects in, 266, 267, 390, 416; use of Liothrips urichi against Clidemia hirta in, 390; Cerambycid intercepted

in Syncarpia logs from Australia in, 267; bunchy-top of banana introduced into Australia from, 46.

filamentosus, Pseudococcus,

Filbert (see Corvlus).

fimetarius, Onychiurus, finitella, Tlascala,

Fiorinia theae, on Camellia and Citrus in U.S.A., 188, 189.

Firebrat (see *Thermobia domestica*). Fish, Dried, Coleoptera infesting, 367. Fish Oil, as an adhesive for sprays, 19, 372.

flabellipes, Geonemus.

Flame-throwers, of doubtful value against locusts, 133.

flammea, Panolis.

flava, Hoplocampa: Sipha. flaveola, Poëphilina (Poëphila). flavescens, Empoasca (Chlorita).

flavilatera, Tomaspis.

flavipennis, Aphrastobracon; Stomatodexia (Leskiopalpus).

flavus, Aphelinus; Leptomastix. Flax, pests of, in Australia, 74. 248: pests of, in Belgium, 291; pests of, in Eire, 208; pests of, in Italy, 268, 269; effect of benzene hexachloride on, 413.

Flea-beetle, Beet (see Chaetocnema tibialis).

Flea-beetle, Flax (see Aphthona euphor-

Flea-beetle, Oak (see Haltica quercetorum).

Flea-beetle, Potato (see Epitrix cucumeris and E. tuberis).

Flea-beetle, Striped (see Phyllotreta vittata).

Fleahopper, Garden (see bracteatus).

fletcherella, Coleophora (see C. cerasivorella).

floccifera, Pulvinaria.

floralis, Nemorilla; Plagionotus (Clytus). Florida, Citrus pests in, 13, 83; status of Gnorimoschema operculella in, 56; potentially injurious Tingid in, 23, 24; natural enemies and biological control of Coccids in, 83, 316; Coccinellid introduced into Porto Rico from, against Asterolecanium spp., 54.

Florida Red Scale (see Chrysomphalus ficus).

floridae, Bruchus (Acanthoscelides). floridensis, Epigrimyia (Siphophyto).

Flour, pests in, and their control, 16, 68, 69, 140, 282, 309, 310; suitability of types of, for insects, 265, 310; as carrier for DDT dust, 266; not protecting stored beans from Bruchus obtectus, 21.

Flour Beetle, Long-headed (see Latheticus

Flour Beetle, Red (see Tribolium castaneum).

Flowers, legislation against introduction of pests and diseases into U.S.A., Hawaii and Porto Rico with, 71, 209.

[difluorodiphenyltrichlor-Fluoro-DDT ethane], toxicity of, to insects, 274, 421; liberation of HC1 by, 275. fockeui, Vasates (Phyllocoptes).

Foeniculum vulgare var. dulce (see Fennel). Fog Generator, insecticides applied from, 87, 88, 147.

forbesi, Cremastus.

Forest Entomology, bibliography of, of Croatia, 104.

Forest Tent Caterpillar (see Malacosoma disstria).

Forests, pests of, in Austria, 128, 407; pests of, in Belgium, 277, 353; pests of, in Britain, 315; pests of, in Canada, 33-36, 78, 163, 207, 208, 239, 240, 255, 319, 320, 321, 334; Eulecanium corvli in, in Czechoslovakia, 65; pests of, in Germany, 75, 239, 407; pests of, in Holland, 112, 353; Hyphantria cunea in, in Hungary, 254; Scolytids in, in Italy, 195; Lymantria dispar in, in Jugoslavia, 115, 116, 117; Sirex in, in New Zealand, 244; Celosterna scabrator in, in Réunion, 317; pests of, in Switzerland, 273, 407, 408; pests of, in Tasmania, 70; pests of, in U.S.A., 14, 33, 36, 87, 92, 143, 144, 158, 220, 234, 243, 254, 301, 320, 333, 334, 336, 388; regulation of density of insect populations in, 352, 353.

Forficula auricularia, dusts against, in

California, 187, 399.

Formica fusca var. subsericea, predacious on Cydia pomonella in W. Virginia, 220; effect of sprays on, 220.

Formica pallidefulva schaufussi incerta, predacious on Cydia pomonella in W. Virginia, 220; effect of sprays on, 220.

Formica rufa, destroying Anthonomus pomorum in Britain, 315.

formicarius, Cylas; Thanasimus.

Formicococcus tafoensis, sp. n., foodplants of, in Gold Coast, 110.

Formosa, Xyleborus fornicatus on tea in, 251; new parasites of Lepidoptera in, 177, 208.

formosa, Encarsia.

formosanus, Coptotermes.

fornicatior, Xyleborus fornicatus.

fornicatus, Xyleborus.

Forsythia, Pseudocneorrhinus bifasciatus on, in New York, 301.

foveicollis, Aulacophora.

foveolatus, Lissorhoptrus.

Fowls, use of, against Pantomorus godmani, 345: unaffected by feeding on chlordane-poisoned grasshoppers, 227.

fragaefolii, Capitophorus.

fragariae, Capitophorus (Pentatrichopus); Macrosiphum; Tarsonemus (see T.

pallidus).

France, Cossus cossus in, 32; Locusta migratoria in, 237; Melolontha melolontha in, 280; orchard pests in, 66, 238, 277, 293, 296; books on pests in, 105, 405; parasites of Leptinotarsa decemlineata in, 349; Rodolia cardinalis introduced into Persia from, 387.

frangulae, auct., Aphis (Doralis) (see

A. gossypii).

Frankliniella, transmitting spotted wilt of tomato in Brazil, 76.

Frankliniella occidentalis, insecticides against, on onion in New Mexico, 126.

Frankliniella schultzei, method of reducing loss of tobacco from spotted wilt transmitted by, in Transvaal, 45.

Fraxinus (see Ash).

Freon-12 (see Dichlordifluoromethane). frit, Oscinella.

froggatti, Typhlocyba.

Froghopper, Demerara Sugar-cane (see Tomaspis flavilatera).

Froghopper, Sugar-cane (see Tomaspis saccharina).

Frogs, toxicity of insecticides to, 174, 347.

frugiperda, Laphygma.

Fruit, Dried, pests in, 265, 309, 310.

Fruit-fly, Mediterranean (see Ceratitis capitata).

Fruit-fly, Mexican (see Anastrepha ludens).

Fruit-tree Red Spider (see Paratetranychus pilosus).

frustrana, Rhyacionia.

frutetorum, Gilpinia (Diprion).

fulleri, Pantomorus (see P. godmani). Fuller's Earth, as carrier for dusts, 22;

sprays of micronised DDT on, 58. fulmeki, Polynema.

fulva, Aphaenogaster.

fulvovestitus, Anomocaulus.

fulvus, Desmoris.

fumiferana, Harmologa (Archips).

fumiferanae, Apanteles; Glypta; Phyto-

Fumigants, equipment for applying, 30, 105, 356; book on control of insects with, 297.

funebris, Bruchophagus (see B. gibbus). funestus, Lagochirus.

Fungi, Entomogenous, 24, 40, 83, 84, 119, 171, 172, 186, 220, 263, 303, 315, 319, 385, 386; utilisation of, 220, 303. Fungi, Injurious, 165, 192, 229, 230, 282, 320, 353, 361, 362, 363, 401; relations of insects to, 163, 192, 220, 221, 293, 387; mite associated with, 202.

Fungus, Green Muscardine (see Metarrhizium anisopliae).

Furniture, protection of, against termites, 169. Fusarium juruanum, infesting Chrysom-

phalus dictyospermi in Persia, 386. fusca, Busseola; Formica; Lachnosterna (Phyllophaga).

fuscedinella, Coleophora (Eupista).

fuscicornis, Podagrica.

fuscipennis, Dahlbominus (Microplectron).

futilalis, Pyrausta.

futilis, Lachnosterna (Phyllophaga). Fygran A, 319.

G

Galeruca semipullata (see Galerucella). Galeruca tanaceti, dusts against, on vegetables in Italy, 346.

Galerucella luteola, insecticides against, on elm in U.S.A., 36, 143, 234, 336.

Galerucella semipullata (Fig-leaf Beetle), Pentatomid predacious on, in New South Wales, 108.

Galerucella tenella, on strawberry in Belgium, 353.

Galerucella xanthomelaena (see G. luteola). gallicus, Polistes.

Gambia, migrations of Schistocerca gregaria in, 207.

gamma, Autographa (Plusia).

Gammexane, 237, 280, 281, 309, 369, 404. Gammexane Dust D.034, 133, 134.

Garden Fleahopper (see Halticus bracteatus).

Gardenia, Coccids on, 54, 101, 187; methyl-bromide fumigation of blooms of, 186, 187.

Garlic, mite on bulbs of, in U.S.A., 83. Gasterocerodes brasiliensis (see Eutinobothrus).

Gastrophysa, Lathromeris bruchocida not parasitising, 242.

gei, Macrosiphum. Geigy 33, 38, 134. Geigy 33 Liquid, 39.

Gelechia trialbamaculella (see Aroga). gelechiae, Bracon (Microbracon).

gelotopoeon, Thyreion.

gemellata, Rossi, Chrysomela (Chrysolina) (see C. quadrigemina).

gemmatalis, Anticarsia.

Genicide-DDT, 58.

genitalis, Hylemvia.

Geocoris, little affected by DDT in California, 91.

Geometra piniaria (see Bupalus).

Geonemus flabellipes, on vines in Spain, 198.

Georgia (U.S.A.), insects in shelled ground-nuts in, 25; Pantomorus and legislation against it in, 184, 398; Scapteriscus vicinus in, 93, 94; identity of parasite introduced against Pseudococcus boninsis in, 316.

georgiana, Tortrix (Archips).

germanica, Blattella. germanicus, Rhynchites.

Germany, forest pests in, 75, 239, 407; parasites and hyperparasites of Diprion pini in, 239; parasites of Apanteles glomeratus in, 406, 407.

Gesafid, 60, 61.

Gesarol, 62, 63, 112, 237, 239, 268, 269, 346, 347.

Gesarol A3, 101, 134.

Gesarol A5, 134.

Gesarol A20, 101, 102, 134.

Gesarol AK40, 58, 133, 325, 326.

Gesarol AK50, 324, 325, 326.

Gesarol M, 104. Gesarol SH5, 103.

ghesquierei, Lygus.

gibberosus, Triaspis (Sigalphus).

gibbus, Bruchophagus.

giffardi, Dirhinus; Tetrastichus.

giffardianus, Tetrastichus. gifuensis, Macrocentrus.

gigantorhynchus, Diptilomiopus (Epitri-

Gilbert Is., Papuana huebneri introduced into, 390.

Gilpinia frutetorum, spray against, on pine in U.S.A., 143.

Gilpinia hercyniae, disease, parasites and biological control of, on spruce in Canada, 207-208, 319, 320, 321; bionomics of, 321.

Gilpinia pallida, characters of excreta of, on pine in Germany, 76.

Gilpinia polytoma, characters of excreta of, on spruce in Germany, 75, 76. gilvifrons, Stethorus.

Gipsy Moth (see Lymantria dispar).

Gladiolus, Taeniothrips simplex on, in U.S.A., 234, 291, 331; selenium treatment of, against T. simplex, 291.

Gladiolus Thrips (see Taeniothrips simplex).

glaucinans, Pericyma.

Gleditsia triacanthos, favouring Acrosternum hilare in Virginia, 373.

66736

D 2

Gliricidia sepium, tests of toxicity of, to insects, 52.

glomeratus, Apanteles.

gloveri, Lepidosaphes (Mytilococcus).

Glue, in sprays, 199, 235; used in onion seed treatment with insecticides, 283. glycinae, Phalaenoides.

Glycinonitriles, low toxicity of, to Cydia pomonella, 93.

Glycyphagus destructor, infesting cheese in New Zealand, 416.

Glycyphagus domesticus, infesting cheese in New Zealand, 416.

Glypta fumiferanae, parasite of Harmologa fumiferana in Canada, 255, 320, 321; bionomics of, 255.

Glypta haesitator, parasite of Cydia nigricana in Britain, 263.

Gnathocerus cornutus, in dried cassava roots in Uganda, 367; tests with benzene hexachloride against, 235.

gnidiella, Cryptoblabes.

Gnorimoschema, sprays against, on tomato in Peru, 133.

Gnorimoschema operculella, in Argentina, 109; in Australia, 46, 69, 178, 216, 217, 249, 363; in Bermuda, 49; in California, 56, 186; question importance of, in U.S.A., 56; on field potatoes, 46, 49, 56, 178, 216, 217, 249; on stored potatoes, 46, 49, 56, 186, 217, 249; on tobacco, 69, 178; on tomato, 49. 363: parasites and biological control of, 49, 109, 216; insecticides against, 46, 69, 217, 249, 363; other measures against, 56, 186, 249; factor affecting rearing of Macrocentrus ancylivorus on, 18.

godmani, Pantomorus.

Gold Coast, study of Coccids and associated insects in connection with swollen-shoot of cacao in, 110; Dacus spp. in, 176.

Gold-tail Moth (see *Euproctis similis*).

Golden Rod (see Solidago).

Golf Greens (see Turf).

Gonatocerus, 407.

Gonipterus scutellatus (on Eucalyptus), in E. Africa, 106, 367, 405; in Mauritius, 125; differences in susceptibility of Eucalyptus spp. to, 106, 367; establishment of Anaphoidea nitens against, 106, 125, 405.

Gonocephalum simplex (see Dasus).

Gooseberry, Itame wauaria on, in Britain, 359; Nematus ribesii on, in U.S.A., 335. Gorse (see *Ulex europaeus*).

Gorse Seed Weevil (see Apion ulicis).

Goryphus nursei, bionomics of, parasitising Earias spp. in Punjab, 121.

Gossyparia spuria, insecticides against, on elm in U.S.A., 143, 234.

gossypiella, Platyedra.

gossypii, Aphis; Phenacoccus. Gossypium (see Cotton).

Gracilaria, on blueberry in New Jersey, 137.

Gracilaria azaleella, DDT against, on azalea in Switzerland and Belgium, 60, 292; bionomics of, 60.

Gracilaria theivora, new parasite of, in Formosa, 177.

gracilariae, Apanteles.

gracilis, Compsolacon; Dacnusa; Scym-

Graeffea crouani, on coconut in Fiji, 390. Grain Beetle, Flat (see Laemophloeus minutus).

Grain Beetle, Saw-toothed (see Oryzaephilus surinamensis).

Grain Borer, Lesser (see Rhizopertha dominica).

graminum, Pediculoides (Pediculopsis); Toxoptera.

Granadilla (see Passiflora quadrangularis). granaria, Calandra (Sitophilus).

granarium, Macrosiphum (see M. avenae).

granati, Tenuipalpus.

grandalis, Megastes.

grandicorne, Xanthogramma.

grandiosella, Diatraea.

grandis, Anthonomus. granicollis, Coelosternus.

Grape Berry Moth (see Polychrosis viteana).

Grape Leaf Skeletoniser, Western (see Harrisina brillians).

Grapefruit, pests of, in Eritrea, 108;
Dacus ferrugineus dorsalis infesting, in Hawaii, 249; tolerance of fruit of, to methyl bromide, 80.

Graphognathus (see Pantomorus).

Grapholitha molesta (see Cydia). Graphosoma lineatum, 375.

Grass, Para (see Brachiaria mutica).

Grass, Reed (see Miscanthus japonicus).

Grass, Sudan (see Sorghum sudanense). Grass, Timothy (see *Phleum pratense*).

Grasses, Laphygma spp. on, in E. Africa. 45, 426; Lachnosterna damaging, in N. America, 14, 164; Adoretus versutus on, in Fiji, 266; Persectania composita damaging, in New Zealand, 244; pests of, in Queensland, 307, 308; locusts and grasshoppers damaging, 48, 109, 227, 228, 294, 310, 311; cereal and sugar-cane pests on, 37, 49, 128, 147, 267, 304, 332, 387. (See Turf.)

Grasshopper, Differential (see Melanoplus differentialis).

Grasshopper, Red-legged (see Melanoplus femur-rubrum).

Grasshopper, Two-striped (see Melanoplus bivittatus).

Greece, Dacus oleae on olive in, 292.

Greenhouse Fumigation, 361.

Greenhouse Leaf-tier (see *Phlyctaenia rubigalis*).

Greenhouse Whitefly (see *Trialeurodes vaporariorum*).

Greenhouses, book on pests in, in England, 173.

greeni, Bracon (Microbracon).

gregaria, Schistocerca, grenadensis, Ipobracon.

Grey-banded Leaf-roller (see Eulia mariana).

grioti, Charips; Sagaritopsis (Sagaritis).

grossus, Ptyelus.

Groundnut Oil, dinitro-o-cyclohexylphenol in emulsion of, against *Quadraspidiotus perniciosus*, **296**; in formulae for insecticide wall washes, **139**.

Groundnuts, pests of, in tropical Africa, 250, 426; new mite on, in Egypt, 411; pests of, in Solomon Is., 390; pests of, in U.S.A., 20, 337, 338; Aphis laburni transmitting rosette disease of, 426; effect of insecticides on, 338.

Groundnuts (Stored), pests in, and their control, 25, 44, 426; development of *Ephestia elutella* on, 265; bibliography of pests of, 272.

Gryllotalpa africana, outbreak of, in Mauritius, 210.

Gryllotalpa gryllotalpa, damaging vegetables in Persia, 388.

Gryllotalpa unispina, damaging vegetables in Persia, 388.

gryllotalpa, Gryllotalpa.

Gryllulus (Gryllus) desertus, damaging vegetables in Persia, 388; measures against, damaging grape vines in Turkmenistan, 4.

Guarea cedrata, new Coccid on, in Gold Coast, 110.

Guava, Aphis gossypii on, in Fiji, 413; pests of, in Ivory Coast, 67, 389; pests of, in Punjab, 123.

Guayule (see Parthenium argentatum).

Guiana, British, Sacadodes pyralis on cotton in, 126; sugar-cane pests in, 302-305; beneficial insects and biological control in, 304, 305.

Gum Arabic, in DDT spray, 28.

Gum Tree, Red (see Eucalyptus rostrata). Gymnosoma, parasite of Eurygaster integriceps in Tadzhikistan, 241.

Gypsum, as carrier for dusts, 90, 112, 113, 114, 133, 140, 222, 223, 329; other uses of insecticides with, 289, 412. Gyron, 269, 346, 347.

H

Habrobracon (see Bracon).

Habrocytus, parasite of Apanteles glomeratus in Europe, 406.

Habrocytus eucerus, 406; hyperparasite of Diprion pini in Germany, 239.

Habrocytus phycidis, hosts of, in Quebec, 321.

Habrocytus poecilopus, parasite of Apanteles glomeratus in Europe, 406, 407; bionomics of, 406.

Habrolepis dalmanni, utilisation of, against Asterolecanium variolosum in Tasmania, 70.

Hackberry (see Celtis occidentalis). haemorrhoidalis, Heliothrips.

haesitator, Glypta.

Hairy Chinch Bug (see Blissus hirtus).

halli, Nilotaspis.

Halotydeus destructor, on flax and leguminous plants in W. Australia, 74, 244; damaging vegetables in New Zealand, 244, 309; measures against, 74, 309.

Haltica, on beans in Uganda, 367.

Haltica bimarginata, comparison of toxicity of benzene hexachloride and DDT to, 22.

Haltica quercetorum, difference in susceptibility to benzene hexachloride of adults and larvae of, 296.

Halticus bracteatus, food-plants of, in U.S.A., 187, 234, 394; eradication of, in California, 187; tests of insecticides against, 234, 394.

hampei, Stephanoderes (Hypothenemus).

Haplomyza imitans, food-plants and control of, in New South Wales, 73. hariolus, Phaeogenes.

Harlequin Cabbage Bug (see Murgantia histrionica).

Harmologa fumiferana, on Abies balsamea and spruce in Canada, 33, 34, 78, 207, 239, 240, 255, 319, 320; in U.S.A., 143; bionomics of, 320; parasites and biological control of, 78, 207, 240, 255, 320, 321; sprays against, 33, 34, 143.

Harpactor iracundus (see Rhinocoris).
Harpalus rufipes (pubescens), on straw-

berry in Belgium, 354.

Harrisina brillians, on vines in California,

185, 190, 397; bionomics of, 190; measures against, 185, 397.

Harvester Ants (see *Pogonomyrmex*).

Hawaii, Coptotermes formosanus in, 249; Dacus ferrugineus dorsalis in, 249; new mite on mango in, 80; quarantines and precautions against introduction of pests into, 71, 79, 105, 209; pests intercepted in quarantine in, 240; attempted introduction of Pseudococcobius terryi into other countries from, 316.

Hawthorn (see Crataegus).

haywardi, Campoplex; Timocratica.

Heat, treatment of timber with, against *Lyctus*, 1, 248; treatments of cottonseed with, against *Platyedra gossypiella*, 66, 105; other uses of, against pests, 114, 384. (See Solar Heat and Water, Hot.)

hebesana, Argyroploce (Endothenia).

hebetor, Bracon (Microbracon).

Hedera helix, Aphis hederae on, in Utah, 388.

hederae, Aphis.

heichinensis, Apanteles.

heldoranus, Cloanthanus.

helianthana, Suleima.

Helianthus (Sunflowers), Homoeosoma electellum on, in Cuba, 83, 84; pests of, in U.S.A., 84, 150, 151; strains of, resistant to Desmoris, 151.

Helianthus tuberosus (Jerusalem Artichoke), pests of, in U.S.A., 84, 150.

Helicopters, equipment and uses of, for applying sprays, 168, 260.

Heliopsis helianthoides var. pitcheriana, Homoeosoma electellum on, in Missouri, 84.

Heliopsis longipes, structure of insecticidal amide of, 25; formerly recorded as Erigeron affinis, 12, 25.

Heliothis armigera (obsoleta), in Australia, 69, 74, 177, 178, 248, 307, 362, 363; in U.S.A., 10, 12, 13, 14, 95, 126, 157, 158, 229, 230, 231, 232, 234, 370; on beans, 12, 13, 229, 230; on cabbage, 177, 178; on cotton, 231, 232; on flax, 74, 248; on maize, 12, 14, 95, 126, 157, 158, 177, 234, 307, 370; unusual injury to maize by, 157; on tobacco, 69, 178; on tomato, 10, 178, 362, 363; forecasting of infestations by, 10; sprays and dusts against, 10, 14, 22, 74, 178, 229, 230, 231, 232, 362, 363; treatment of maize ears against, 12, 13, 126, 158, 234, 370; DDT aerosol ineffective against, 95; calcium arsenate favouring infestation by, 231.

Heliothis virescens, insecticides against, on tobacco in S. Carolina, 372.

Heliothrips haemorrhoidalis, on tobacco in Australia, 178; on Cinchona in

Porto Rico, 53; on greenhouse plants in Switzerland, 282; insecticides against, 53, 282.

Hellula phidilealis, dusts against, on crucifers in Trinidad, 319.

Hellula undalis, on beet and cabbage in Australia, 177; insecticides against, 178.

Helomyia lateralis (parasite of Eurygaster integriceps), bionomics of, in Tadzhi-kistan, 241.

Helopeltis, dusts against, on tea in Travancore, 250, 251.

Helopeltis bergrothi, on cotton in Uganda, 366.

Helopeltis theivora, outbreak of, on nursery Eucalyptus in Madras, 251.

Helophorus, on vegetables in England and Holland, 173, 190.

helvolus, Metaphycus.

Hemerobius, predacious on Anuraphis devecta in Holland, 66.

Hemiberlesia howardi, food-plants of, in California, 399, 400.

hemipterus, Metamasius.

Hemisarcoptes malus, predacious on Lepidosaphes ulmi in Nova Scotia, 392; sulphur sprays destroying, 392.

Hemiteles ridibundus, parasite of Cydia nigricana in Britain, 263.

Hemp, Mirids on, in Uganda, 26. Hepialus behrensi, on apple in California,

heraclella, Aphis.

Hercothrips fasciatus, in California, 79, 399; fumigation of Citrus fruits for export to Hawaii against, 79; winter food-plants affecting infestation of beans by, 399.

Hercothrips marginipennis, on maize in California, 187.

Herculia intermedialis, on spruce in Ouebec, 321.

hercyniae, Gilpinia.

hesperidum, Coccus.

hesperus, Lygus.

Hessian Covers, to protect bananas from thrips, 307.

Hessian Fly (see Mayetiola destructor).

Heterocampa, spray against, on shade trees in U.S.A., 143.

Heterocordylus malinus, on apple in New York, 271.

Heteronychus sanctae-helenae, soil treatments against, damaging grasses and maize in New South Wales, 73, 138.

heteropneusticus, Coccophagus (see C. caridei).

Heteroscapiscus, n.n. for Heteroscapus, Brèth., 316.

Heteroscapus, Brèth. nec Faust (see Heteroscapiscus).

Hevea, Xyleborus fornicatus on, in Netherlands Indies, 251.

Hexabutyl Tetraphosphate, toxicity of, to Aphids, 228.

Hexachloryclohexane (see Benzene Hexachloride).

Hexachlorethane, treatment of rice-field water with, against Nymphula nymphaeata, 346.

Hexaethyl Tetraphosphate, against Aphids, 14, 228; ineffective against Onychiurus fimetarius, 305; against Psylla pyricola, 335; against Tetranychid mites, 154, 228, 246; toxicity of, to bees, 222; in dust, 222; in sprays, 154, 222, 228, 246, 335; duration of effectiveness of, 222, 228.

Hexapropyl Tetraphosphate, toxicity of, to Aphids, 228.

hibisci, Cerococcus.

Hibiscus calyphyllus, Dysdercus on, in Swaziland, 27.

Hibiscus cannabinus, Earias spp. on, in Punjab, 312.

Hibiscus esculentus, Earias spp. on, in Punjab, 312.

Hibiscus rosa-sinensis, Aphis gossypii on, in Fiji, 413.

Hibiscus tiliaceus, Adoretus versutus on, in Fiji, 266.

Hieroglyphus nigrorepletus, bionomics of, in United Provinces, 122.

hilare, Acrosternum.

hilaris, Aulacophora (Ceratia).

hippocoon, Papilio dardanus.

Hippodamia convergens (in U.S.A.), killed by benzene-hexachloride dust, 100; DDT apparently not harmful to, 147. hipposideros, Lyctus.

Hirsutella, infesting Leptopharsa machalana vinnula in Florida, 24.

hirtibasis, Arachnophaga.

hirtipennis, Epitrix.

hirtiventris, Lachnosterna.

hirtus, Blissus.

histrionica, Murgantia.

Hochberg-LaMer Generator, 87, 88. Holland, pests of currant in, 285, 408; forest pests in, 112, 270, 353; orchard pests in, 66, 111, 129, 130, 276, 285, 286; Vasates fockeui on greenhouse peach and plum in, 239; pests of vegetables in, 130, 190, 270, 283, 285, 286; beneficial insects in, 66; standards for insecticides and fungicides in, 295; tests of pyrethrum grown in, 270.

Holly, Prionus coriarius on, in England, 215.

Holocremnus incrassator, parasite of Neurotoma nemoralis in Poland, 255.

Homadaula albizziae, spray against, on Albizzia in U.S.A., 143.

Homala polita, infesting stored ground-

nuts in Senegal, 44. Homoeosoma electellum, on sunflower in Cuba, 83, 84; bionomics of, in U.S.A.,

84; natural enemies and attempted biological control of, 83, 84.

homoeosomae, Apanteles.

Homona coffearia, new parasites of, in Formosa, 177.

Homorocoryphus vicinus, damaging cereals in Nyasaland, 44.

Honduras, British, digests of plant pest legislation in, 105, 209.

Honey, used for feeding insects, 29, 106, 126, 222, 258, 328.

Honey Locust (see Gleditsia triacanthos). Hoplocampa brevis, dust against, on pear in Holland, 286.

Hoplocampa flava (on plum), in Britain, 359; sprays against, in Switzerland, 296.

Hoplocampa minuta (on plum), in Holland, 285, 286; in Switzerland, 296; sprays against, 285, 296.

Hoplocampa testudinea (Apple Sawfly), in Britain, 359; in Holland, 286; on Astrakhan apple in U.S.A., 335; sprays against, 286, 335.

Hops, pests of, in Britain, 287, 288, 358. Horismenus, hyperparasite of Toxoptera

graminum in Uruguay, 50.

Horismenus missouriensis, parasite of Bruchus floridae in Oklahoma, 9.

horni, Dicerca,

Horogenes cacoeciae (see Angitia).

hortuellus, Crambus.

House-fly (see Musca domestica).

howardi, Hemiberlesia.

Howardia biclavis, on Citrus in California, 189.

Hubam Clover (see Melilotus alba).

huebneri, Papuana.

Humbe tenuicornis, on Eleusine in Uganda, 367.

Humidity, relations of locusts to, 171; relations of other insects to, 8, 243, 251, 252, 266, 371, 378, 379, 391; effect of, on *Tyroglyphus farinae*, 417, 418; effects of, on insecticides, 289, 330, 378, 379.

humilis, Iridomyrmex.

Hungary, miscellaneous pests in, 61-64, 254; natural enemies of *Bruchus pisorum* in, 62.

hyalinata, Diaphania.

Hyalomya cuthbertsoni (see Alophora nasalis).

Hyalopterus arundinis (on stone fruits), bionomics of, in Italy, 191, 192; in Spain, 199; in Turkey, 352; sprays against, 192, 199, 352.

Hyblaea puera, on teak in Samoa, 267. Hyderabad, new gall-midge and associated insects on castor in, 110.

Hydnocera, attacking Chrysopa in California, 396.

Hydnocera pubescens, attacking Lepidoptera in U.S.A., 84.

Hydrangea, Diaspidiotus ancylus on, in California, 399; not injured by azobenzene, 284.

Hydrochloric Acid, for removing DDT residues from apples, 135, 136.

Hydrocyanic Acid Gas, against Aphids, 247, 361; against Coccids, 30, 65, 79, 80, 185, 189, 197, 292, 387; oil sprays not decreasing resistance of Aonidiella aurantii to, 160, 161; use of, against Epilachna varivestis, 397; thrips, 79, 385; against Trialeurodes fumigation of vaporariorum, 361; grain bags in barges with, 355-357; greenhouse fumigation with, 361; fumigation of nursery stock with, 30, 65, 247, 292; tent fumigation with, 185, 189, 197, 385, 387; and injury to plants, 361; effects of fumigating Citrus fruits with, 79, 80; methods of generating and applying, 30, 80, 185, 356; comparison of other fumigants with, 79, 292, 357. (See Calcium Cyanide.)

Hydrogen Chloride, liberation of, as explanation of insecticidal action of chlorine compounds, 274, 275.

Hydrogen-ion Concentration, effect of, on dinitro-o-cresol, 111, 274, 287.

Hydroxyapatite, 329.

2'-Hydroxy-2,4,4,4',7-pentamethylflavan, tests of, against insects and mites, 13, 341; and DDT, 13, 341.

Hylastes, species of, on conifers in Britain, 315.

Hylastes angustatus, 315.

Hylastes ater, 315.

Hylastes attenuatus, 315. Hylastes cunicularius, 315.

Hylastes palliatus, 315.

Hylastes rufipes, in Quebec, 163, 320; in U.S.A., 15, 103, 320; vector of Dutch elm disease, 163, 320; effect of phloem condition on infestation of elm logs by, 15; measures against, 103.

Hylemyia antiqua (on onion), in Canada, 89, 90, 156; in England, 173; in Holland, 270, 283; in N. Dakota, 388; relation of, to decay in onions, 283; on leek, 270; on shallot, 388; measures and experiments against, 89, 90, 156, 270, 388.

Hylemyia brassicae (on crucifers), in Belgium, 278; in England, 173; in U.S.A., 102, 222, 350; bionomics of, 222; measures and experiments against, 102, 222, 278, 350.

Hylemyia cilicrura, bionomics of, on vegetables in England, 173; on beans in Switzerland, 281; measures against,

174, 281.

Hylemyia genitalis, on wheat in Italy, 114; cultural measures against, 115.

Hylemyia trichodactyla, bionomics and control of, in England, 173, 174.

Hylesinus vestitus, bionomics of, on pistachio in Persia, 386.

Hylotrupes bajulus, experiments against, in timber in S. Africa, 107.

Hylurgopinus (see Hylastes).

Hylurgops (see Hylastes).

Hylurgus ligniperda, in felled pine in Italy, 195.

Hymenia recurvalis, on beet in Australia, 177; insecticides against, on spinach in Virginia, 331.

Hyoscyamus, experiments with Macrosiphum solanifolii and, 362; relation of Aphids to virus of, 423, 424.

Hypera brunneipennis, bionomics of, in Arizona, 55; liberation of Bathyplectes curculionis against, 55.

Hypera meles, on beans and Trifolium spp. in Ontario, 166.

Hypera nigrirostris, on clover in California, 187; bionomics and control of, in Italy, 115; on beans in Ontario, 166.

Hypera variabilis, sprays against, on lucerne in Spain, 203, 204.

hyperici, Agrilus; Chrysomela (Chrysolina).

Hypericum androsaemum, use of Chrysomela hyperici against, in New Zealand, 244.

Hypericum perforatum, use of insects against: in Australia, 216; in California, 188, 216; in New Zealand, 244.

Hyphantria cunea, introduction and biology of, in Hungary, 254; in U.S.A., 137, 143, 234, 254, 336; on fruit trees, 254, 336; on other plants, 137, 143, 254; insecticides against, 143, 234, 336.

Hyphantria textor, H. cunea confused with, in Hungary, 254.

Hyponomeuta, tests with DDT against, on willow in Turkey, 352.

Hyponomeuta padellus malinellus, sprays against, on apple in Holland and Turkey, 276, 352.

Hypothenemus hampei (see Stephanoderes). hypparchus, Pompostola.

Hyssopiscus, n.n. for Hyssopus, Gir., 316. Hyssopus, Gir. (nec Isopus, Montr.), new name proposed for, 316.

Hysteropterum sepulchralis, on Parthenium

argentatum in Mexico, 97.

T

Icerya purchasi (on Citrus), effect of insecticides on control of, by Rodolia cardinalis in California, 145, 160; introduction and "use of R. cardinalis against, in Persia, 387.

Idaho, Bruchus pisorum in, 12; Cydia molesta in, 185; Limonius californicus

in, 91.

Idechthis (see Nemeritis).

Idiocerus stali, sprays against, on pistachio

in Persia, 386.

Igepal CA Extra High Concentrate, 158. Illinois, greenhouse pests in, 290, 291; maize pests in, 157, 158; Melanoplus spp. in, 92, 225-228; orchard pests in, 95, 127, 322, 323; Epicauta spp. on potato in, 228; sunflower pests in, 84, 150; vine pests in, 127; beneficial insects in, 84, 127, 151.

imbricatus, Epicaerus.
imitans, Haplomyza.
impersonatella, Diatraea.
inaequalis, Coelophora.
Inareolata (see Angitia).
includens, Scymnus.
inconspicua, Sturmia.
incrassata, Phorocera.
incrassator, Holocremnus.

Incurvaria capitella, sprays against, on currant in Holland, 285.

India, Coleoptera infesting bamboos in, 119; pests of cereals and grasses in, 267, 427; cotton pests in, 120, 121, 310, 312; Schistocerca gregaria in, 122, 310-312; Xyleborus fornicatus on tea in, 251; pests of vegetables in, 123; beneficial insects in, 119, 427. (See also under the various Provinces.)

Indiana, apple pests in, 13.

indicum, Syntomosphyrum (Melittobia); Tribolium.

indicus, Paratetranychus.

inferens, Sesamia.

infirmus, Lygus.

inflata, Pimpla.

Inga, susceptibility of species of, to borers in Porto Rico, 300. inopinata, Melanaspis (Aonidiella).

inornatus, Cryptus.

Inositol, question of competition of benzene hexachloride with, in cell metabolism, 175, 215, 277.

Insect Cuticle, permeability of, by insecticides, 174, 175; toxicity of dusts increased by carrier abrasive to, 330.

Insect Specimens, Dermestid damaging,

37; preparation of, 104, 348.

Insecticides, tests of organic compounds as potential, 13, 14, 87, 93, 274, 275, 403, 421; study on plants as sources of, 9, 51, 52, 298, 299; review of literature on, obtained from plants, 308; mode and mechanism of action of, on insects, 174, 175, 215, 274, 275, 277, 330, 369; need for standardisation of tests of, 292, 295; analysis of factorial series of tests of, 381; catalogue of, 297; standards for, in Holland, 295.

Insects, species of, eradicated in various countries, 219; methods of estimating populations of, 310; distance dispersion of, 8; book on attractants and repellents for, 298; surveys of literature on chemotropism and diapause in, 32; micro-respirometer for, 272; mode and mechanism of insecticides on, 174, 175, 215, 274, 275, 277, 330, 369; lists of popular names of, in various countries,

32, 72. insignis, Closterocerus.

insulana, Earias.

insularis, Ptychodes trilineatus.

integriceps, Eurygaster.

intermedialis, Herculia.

intermedius, Dysdercus.

interpunctella, Plodia.

interrupta, Actia; Melasoma (Chrysomela) lapponica.

inversa, Lachnosterna (Phyllophaga).

Iowa, Gnorimoschema operculella of little importance in, 56; Homoeosoma electellum in, 84.

Ipem, 199.

Ipobracon, introduction of, against Diatraea saccharalis in Porto Rico, 54.

Ipobracon grenadensis, parasite of Diatraea in Br. Guiana, 304.

Ips erosus, in felled pine in Italy, 195.

Ips typographus, bionomics of, on spruce in central Europe, 407; measures against, 408.

iracundus, Rhynocoris (Harpactor).

Iran (see Persia).

Iridomyrmex detectus, dusts applied to nests of, in Australia, 218, 305.

Iridomyrmex humilis, measures against, associated with Coccids on Citrus in W. Australia, 74.

Iridomyrmex rufoniger, associated with Anuraphis persicae-niger in New South Wales. 247.

Iris, Agasphaerops nigra not feeding on, 142.

Iris germanica, Argyroploce hebesana on, in U.S.A., 84.

Ischiodon scutellare (see Xanthogramma). Isobutyl-2, 6, 8-decatrienoamide, affinin composed of, 25.

Isophorone, DDT aerosol containing, 230.

Isopropyl Alcohol, used in preparation of Coccids, 104.

Isopus, Montr., 316.

Italian Jasmine (see Jasminum humile). italicus, Calliptamus; Perilampus.

Italy, pests of cereals in, 37, 114, 346, 347, 383; pests of clover and lucerne in, 30, 115; Halticids on flax in, 268; orchard pests in, 30, 64, 191, 192, 194, 253, 269; Prays oleellus on olive in, 141, 193; Cydia pomonella on walnut in, 194; pine Scolytids in, 195; pests of vegetables in, 30, 193, 253, 254, 346, 383; Athalia rosae on cruciferous oil crops in, 192; pests of stored grain in, 30, 38; new mite in condensed milk, etc., in, 114; miscellaneous pests in, 253, 254; list of insect pests in, 141; book on pests and diseases of crops in, 355; beneficial insects in, 31, 37, 65, 191, 192, 193, 195, 269; failure to establish Antheraea pernyi in, 67.

Itame wauaria, bionomics of, in Britain, 359.

Itoplectis conquisitor, hosts of, in Quebec, 320, 321.

Ivory Palm (see Metroxylon vitiensis). Ivy (see Hedera helix).

J

Jack Pine (see Pinus banksiana).

Jamaica, digest of plant-quarantine restrictions in, 105.

Japan, Pseudocneorrhinus bifasciatus in, 301; Pseudococcus comstocki in, 316, 410; introduction of beneficial insects into other countries from, 77, 316, 410. Japanese Beetle (see Popillia japonica).

Japanese Privet (see Ligustrum japonicum)

Japanese Privet (see Ligustrum japonicum). japonica, Popillia.

Jarring, to dislodge insects from trees, 75, 151, 235, 314, 373.

Jasminum humile, experiment with Parlatoria oleae and, in California, 342. javanus, Plaesius.

Jersey, Leptinotarsa decemlineata in, 259.

Jerusalem Artichoke (see Helianthus tuberosus).

jocosatrix, Bombotelia.

johnstoni, Elasmus.

juglandis, Chromaphis (Callipterus).

Juglans regia (see Walnut).

Jugoslavia, Lymantria dispar and associated insects in, 115–117; bibliography of entomology of, 104.

Juniper (Juniperus), eradication of Dichomeris marginella on, in California, 187; pests of, in Poland, 257, 258; new Torymid on, in Rumania, 267.

juniperina, Chlorochroa. Juniperus (see Juniper).

Juniperus bermudiana (Bermuda Cedar), pests of, in Bermuda, 48, 49.

Jute, Tetranychus on, in Bengal, 427.

K

kahawaluokalani, Myzocallis.

Kale, pests of, in England, 132, 173; pests of, in U.S.A., 159, 332, 333.

Kalotermes brevis (in Porto Rico), toxicity of Mammea extract to, 52; treatments of timber against, 169, 232, 233, 300; list of timbers resistant to, 169.

Kalotermes brouni, tests with timber preservatives against, in New Zealand, 245, 414, 415.

Kaolin, ineffective against insects in flour, 141; as carrier for dusts, 138, 140, 218, 340, 341; unsatisfactory carrier for dinitro-o-cyclohexylphenol, 288; benzene-hexachloride spray powders containing, 13, 199; grades of, 340, 341.

Kapok (see Ceiba pentandra).

kargalike, Euproctis (Porthesia).

Keiferia lycopersicella (Tomato Pinworm), in greenhouses in Ontario, 167.

Kenya Colony, cereal pests in, 403, 405; Trypetid on cucurbits in, 176; pests of coffee in, 316, 317, 404; Gonipterus scutellatus on Eucalyptus spp. in, 106, 405; new Mirid in, 26; Schistocerca gregaria in, 123-124, 366, 375, 403; termites in, 404; beneficial insects and biological control in, 27, 106, 404, 405; pests intercepted in quarantine in, 405; introduction of Anaphoidea nitens into Mauritius from, 125.

kenyae, Pseudococcus.

Kerosene, use of, to destroy *Pentalonia* nigronervosa, 46; sump oil in, as spray against locusts, 48; alone or in mixtures against Coleoptera in logs or timber, 103, 248, 414, 415, 416; uses

of DDT in, 22, 35, 36, 88, 89, 96, 103, 109, 210, 336, 337, 342, 364, 404; as solvent or spray carrier for other insecticides, 52, 155, 168, 341, 364, 395, 396; soybean phosphatides in, as adhesive for lead arsenate, 372.

Kerosene Emulsion (see Oil Emulsions). Khaya senegalensis, Cerambycid in, in

Senegal, 250.

Kieselguhr, unsatisfactory carrier for dinitro-o-cyclohexylphenol dust, 288. kiowa, Trachyrhachys.

kivuensis, Anagyrus. Klamath Weed (see Hypericum perforatum).

Kochia, 216.

koehleri, Copidosoma.

Kohlrabi, pests of, in Britain, 132: Helophorus on, in Holland, 190; pests of. in U.S.A., 159.

Korea, new parasite of Pseudococcus comstocki in. 316.

Koster's Curse (see Clidemia hirta).

kraussi, Dociostaurus.

Kryocide, 21. Kryocide D-50, 21.

kuehniella, Ephestia.

kunoense, Eulecanium (Lecanium).

kuntzei, Megastigmus. kuriri, Tyndarichus.

L

laburni, Aphis.

lacca (Lac Insect), natural Laccifer enemies and control of predator of, in Bihar, 120.

lachesis, Campsomeris.

Lachnosterna, outbreak of, in Ontario, 163, 164; investigations on, in Quebec, 167; bionomics of, in Texas, 14. Lachnosterna anxia, in Canada, 164, 167. Lachnosterna fusca, in Ontario, 164. Lachnosterna futilis, in Ontario, 164. Lachnosterna hirtiventris, in Texas, 14, 15. Lachnosterna inversa, in Ontario, 164. Lachnosterna rubiginosa, in Texas, 14, 15. Lachnosterna rugosa, in Ontario, 164. Lachnosterna submucida, in Texas, 14. Lachnosterna torta, in Texas, 14. Lacon murinus, in Hungary, 64.

lactella, Endrosis.

Lactic Acid, survival of Tyroglyphus nadinus in, 114.

Lactuca (see Lettuce).

Lactuca scariola, destruction of, against Hercothrips fasciatus in California, 399. Laemophloeus (in stored products), study on primary infestation of bulk wheat by, in Britain, 418; in Canada, 418; type of bag affecting infestation by, in U.S.A., 25; in Uganda, 367; experiments with DDT against, 19, 28.

Laemophloeus minutus, 19, 418; Australia, 182; fumigants against, in stored wheat, 20, 182.

Laemostenus terricola (see Pristonychus). laeta, Polynema.

laevigata, Atta (Neoatta).

laevinucha, Trichomalus.

Lagerstroemia, Aphid on, in California, 400.

Lagochirus funestus, factors affecting utilisation of, against Opuntia in S. Africa, 171, 172.

Lagria villosa, on lima and tepary beans in Uganda, 367.

lambens, Sarcophaga.

lambertianae, Conophthorus.

lanigera, Oregma.

lanigerum, Eriosoma.

Lantana camara, establishment of Teleonemia scrupulosa against, in Australia, 181.

Lantana Leaf Bug (see Teleonemia scrupulosa).

Laphygma eridania, tests of insecticides against, in U.S.A., 13, 14, 22, 23.

Laphygma exempta, in E. Africa, 45, 426; bionomics and control of, in Queensland, 307; on graminaceous plants, 45, 307, 426; on leguminous crops, 426; parasites and disease of, 45, 307.

Laphygma exigua, plants attacked by, in

E. Africa, 426.

Laphygma frugiperda, on sugar-cane in Br. Guiana, 304; on tomato in Peru, 134; on maize and grasses in U.S.A., 95, 96, 157, 158, 332, 333; measures and experiments against, 51, 52, 95, 96, 134, 158, 332, 333.

laphygmae, Euplectrus.

lapidarius, Biscirus. lapponica, Melasoma (Chrysomela).

Larch, Lymantria monacha on. Austria, 128; Coleophora laricella on, in Canada, 208.

Lard, stored beans treated with, against Bruchus obtectus, 76; experiment with Tyroglyphus nadinus and, 114.

laricella, Coleophora.

laricinellae, Chrysocharis.

Lasioderma serricorne, in Senegal, 44; in U.S.A., 25, 85, 86; factors affecting infestation of shelled groundnuts by, 25, 44; processes affecting control of, in stored tobacco, 85, 86.

Lasiophthicus pyrastri, overwintering of, in Switzerland, 273.

Lasiophthicus seleniticus, overwintering of, in Switzerland, 273.

Laspeyresia (see Cydia).

lata, Phytolyma.

lataniae, Cerataphis.

lateralis, Helomyia (Anantha); Phytomyza.

Latheticus oryzae (in stored cereals), in Switzerland, 38, 39; in Victoria, 182; interaction between temperature and moisture in bulk wheat and, 182; experiments against, 20, 38, 39.

Lathromeris bruchocida, sp. n., parasite of Bruchids in U.S.S.R., 242; bionomics and suggested use of, 242, 243.

Lathyrus spp., Cydia nigricana on, in Britain, 263.

latro, Lixus.

Latrodectus mactans, destroying grass-hoppers in Chile, 109.

latus, Anychus.

Latvia, Rhopalosiphoninus ribesina in, 408.

Laurel, Rhynchites coeruleus on, in Britain, 358.

Laurus nobilis, Chrysomphalus dictyospermi on, in Persia, 386.

Lauryl Isoquinolinium Bromide, ineffective against *Psylla pyricola*, 335.

Lauryl Thiocyanate, 380. (See Thiocyanates.)

Lavatera, Podagrica spp. on, in Rumania, 41.

Lawns (see Turf).

Lead, residues of, from lead-arsenate dusts on apples, 59.

Lead Arsenate, of little value against Pantomorus godmani on Citrus, 345; against pests of crucifers, 41, 123, 133, 218, 319, 385; against maize pests, 332, 370; against orchard pests, 13, 19, 24, 57, 58, 59, 92, 93, 98, 100, 101, 126, 127, 132, 149, 150, 157, 179, 180, 218, 219, 225, 245, 246, 247, 256, 257, 258, 271, 279, 296, 323, 324, 339, 372, 373, 401; strain of Cydia pomonella resistant to, 323; not favouring Tetranychus pacificus, 149; against Prays oleellus on olive, 141, 194; against pests of ornamental plants, 142, 143, 344; against pests of pecan, 151, 152; increasing infestation of pecan by Aphids, 152; against pests of solanaceous plants, 134, 165, 196, 259, 362, 363, 371, 372; against vine pests, 127, 198; against pests on other plants, 137, 319, 383; in bait, 372; dusting with, 57, 58, 141, 218, 319, 332, 362, 363, 370; oil-impregnated dusts of, 58; formulae for, in sprays, 40, 58, 59, 98, 123, 126, 127, 133, 137, 141, 142, 149, 150, 151, 165, 179, 180, 194, 218, 225, 245, 246, 256, 257, 258, 279, 339, 362, 371, 372, 373, 401; units applying concentrated sprays of, 36; adhesives and spreaders 257, 372; and DN-111, 101, 372, 373; lime in sprays containing, 59, 123, 133, 151, 180, 372; and lime-sulphur, 137, 373; and nicotine, 58, 59, 100, 152, 219, 225, 257, 325; and oils, 58, 59, 98, 126, 148, 149, 150, 151, 179, 180, 218, 225, 245, 246, 257, 372; sunflowerseed oil emulsion compatible with, 354; and phenothiazine, 59; and sugar, 296; and sulphur, 57, 59, 165, 362; and Toxaphene, 224, 225; and zinc sulphate, 151, 180, 373; and injury to plants, 151, 152, 247, 332, 373, 401; residues of, on apples, 58, 59, 180, 257; DDT compared with, 35, 59, 165, 179, 218, 245, 246, 323; other insecticides compared with, 14, 19, 24, 59, 93, 245.

Lead Fluoride, repellency of wood treated with, to *Kalotermes brevis*, **232.** Lead Oxide, reducing effect on plants of

calcium arsenate in soil, 371.

Lead Pentachlorphenate, repellency of wood treated with, to *Kalotermes brevis*, 233.

Leaf-cutting Ants (see *Atta*).

Leafhopper, Potato (see *Empoasca fabae*). Leafroller, Grey-banded (see *Eulia mariana*).

Leafroller, Red-banded (see *Eulia velu-tinana*).

Lebia cyanocephala, predacious on Entomoscelis adonidis in Rumania, 40.

Lecaniodiaspis, on Parthenium argentatum in Mexico, 97.

Lecanium, Burm. (see Coccus).

Lecanium, auct. (see Eulecanium).

lecheana, Tortrix (Cacoecia).

Leconte's Sawfly (see Neodiprion lecontei). lecontei, Neodiprion.

lectularius, Cimex.

Leek, pests of, in Britain, 173, 362; seed treatment of, against *Hylemyia antiqua* in Holland, 270.

lefroyi, Bracon.

leguminis, Cydia (Laspeyresia).

Leis conformis, predacious on Anuraphis persicae-niger in Australia, 247; attempted introduction of, into Egypt, 410; bionomics of, 411.

Lema melanopa, bionomics of, on cereals

and grasses in Italy, 37, 38.

Lema trilineata, benzene hexachloride dust against, on potato in New Hampshire, 234.

Lemon, pests of, in Australia, 181, 248; Coccids on, in Spain, 197; pests of, in U.S.A., 89, 187, 188; persistence of DDT spray deposits on, 89; tolerance of fruit of, to methyl bromide, 80. lepelleyi, Pseudococcus.

Lepidiota caudata, in pastures in

Queensland, 308.

Lepidoglyphus destructor (see Glycyphagus). Lepidosaphes, on mango in Egypt, 409.

Lepidosaphes beckii (on Citrus), in Australia, 177; in California, 187, 188, 189; in Persia, 387; in Peru, 134; in Spain, 196, 197; in Trinidad, 319; bionomics of, 188, 189, 197; Chilocorus bipustulatus predacious on, 197; measures and experiments against, 134, 187, 189, 197.

Lepidosaphes gloveri (on Citrus), measures to prevent introduction of, into California from Mexico, 79; of little importance in Persia, 387; bienomics and control of, in Spain, 196, 197; in

Trinidad, 319.

Lepidosaphes machili, methyl bromide against, on orchid in California, 187.

Lepidosaphes newsteadi, on Juniperus bermudiana in Bermuda, 45.

Lepidosaphes pistaciae, on Pistacia in Persia, 386.

Lepidosaphes ulmi (on apple), in California, 400; effect of sprays on natural enemies and abundance of, in Nova Scotia, 392.

Lepisma saccharina, in houses in Belgium, 292; types of textiles attacked by, 324; tests and uses of DDT against, 292, 324, 378, 379; influence of heat and humidity on toxicity of dust insecticides to, 378, 379.

Leptinotarsa decemlineata (Potato Beetle), in Belgium, 291; in Czechoslovakia, 278; in France, 296, 349; in Jersey, 259; in New Hampshire, 234; Ontario, 165; in Poland, 254, 384; in Spain, 196, 199, 293; in Switzerland, 297; experiments with Polish varieties of potato and, 42, 43; bionomics and physiology of, 196, 349, 375; natural enemies of, 196, 349; insecticides against, 165, 196, 199, 234, 259, 278, 291, 293, 296, 297, 349; other measures against, 259, 297.

Leptocoris lurida, habits of, in New South Wales, 108; DDT against, 108.

leptogramella, Calamotropha.

Leptomastix, new parasite of, in Uganda, 109.

Leptomastix flavus, factors affecting value of, against mealybugs in Palestine, 410. Leptopharsa machalana vinnula, bionomics of, in Florida, 23.

Leptospermum, pests of, in California, 187.

Leptothrips mali, predacious on Cydia pomonella in W. Virginia, 219; effect of sprays on, 219.

Leskiomima tenera, parasite of Homoeosoma electellum in U.S.A., 84.

Leskiopalpus flavipennis (see Stomatodexia).

Lespedeza, Melanoplus infesting, in Illinois, 228; effect of border planting of, on insects in Virginia, 374.

Lethane, 396.

Lethane 60, 17.

Lethane A-70, 327.

Lethane B-71, 102, 126, 184, 218, 327, 394, 395.

Lethane B-72, 101, 165, 325, 326, 394.

Lettuce, pests of, in Australia, 74, 177; Agallia albidula on, in Brazil, 77; pests of, in Britain, 173, 361, 362; Macrosteles divisus transmitting aster yellows virus to, in U.S.A., 350, 351; unsuitable food for Limonius californicus, 91; experiments with Macrosiphum solanifolii and, 361, 362; question of DDT dust deposits on, 81; effect of HCN on, 361; selenium absorbed from soil by, 290; Velsicol causing injury to, 421; DDT mixed with stored seed of, against insects, 19.

Lettuce, Prickly (see Lactuca scariola). Leucaena glauca, Ferrisia virgata on, in Gold Coast, 111.

Leucaspis pistaciae, parasites of, Pistacia in Persia, 386.

leucaspis, Anisoplia.

Leucinodes elegantalis, on tomato in Brazil, 76.

leucoloma, Pantomorus.

leucophaearia, Erannis. leucophthalmus, Philaenus.

Leucoptera scitella, bionomics and control of, on apple in Italy, 269; on Pistacia in Persia, 386.

leucopterus, Blissus.

leucotreta, Argyroploce.

leveri, Erythroneura.

Liberia, Trypetid on cucurbits in, 176.

Libya, Diptera on Capsicum in, 72; Schistocerca gregaria in, 206, 207; lists of pests in, 43.

licoides, Castnia.

licus, auct., Castnia (see C. licoides). Light, reactions of Ephestia elutella to. 213, 214; reducing toxicity of insecticides, 168, 299.

ligniperda, Hylurgus. lignosellus, Elasmopalpus. ligustici, Otiorrhynchus.

Ligustrum, Pseudocneorrhinus bifasciatus on, in New Jersey, 301.

Ligustrum japonicum, Parlatoria oleae on, in California, 16.

Lilioceris lilii, on Lilium regale in Quebec, 104.

Lily (*Lilium* spp.), *Lilioceris lilii* on, in Quebec, 104; pests of, in U.S.A., 142, 187; thiocyanoacetates causing injury to, 253.

Lima Beans (*Phaseolus lunatus*), new Jassid transmitting leaf-crinkle of, in Fiji, 266; Coleoptera on, in Uganda, 367; Coleoptera on, in U.S.A., 224, 301, 370, 396; insecticides and injury to, 224, 225; effects of seed treatments on germination of, 331.

limacina, Caliroa.

limatulus, Pseudaphycus.

Limba (see Terminalia superba).

Lime (Citrus), treatment of fruits of, against Coccids, 79, 80.

Lime, as a repellent, 32; as a carrier for dusts, 52, 59, 218, 222, 223, 272, 329, 332, 395; addition of, to arsenical sprays, 41, 59, 123, 133, 151, 180, 372, 373, 404; addition of, to other sprays (see also Bordeaux mixture), 165, 334; uses of mixtures of soy-bean flour and, in sprays, 127, 354; wall washes of DDT and, 139; reducing effectiveness of DDT, 166, 218; calcium and magnesium types of, 329; formulae containing, 41, 52, 59, 123, 133, 139, 151, 165, 180, 222, 223, 332, 334, 372, 373, 404.

Lime-sulphur, against Anthonomus piri, against Coccids, 181, destroying natural enemies of Lepidosaphes ulmi, 392; against Collembola, 305; against mites, 82, 181, 249, formulae for, 46, 137, 249, 344; 278; formulae containing, 46, 137, 373; and DDT, 46, 47, 359; risk of damage to plants by dinitro compound following, 288; and lead arsenate, 137, 373; sunflower-seed oil emulsion compatible with, 354; oil-compatible fungicides compared with, 131, 132.

Limestone, top-dressings of selenium salts in, 289.

Limnerium crassifemur (see Eulimneria). Limonius californicus (in U.S.A.), effect of plants on growth of, 91; D-D mixture against, 187.

lineaticollis, Antestia.

lineatopunctata, Melasoma.

lineatum, Graphosoma; Trypodendron (Xyloterus).

lineatus, Agriotes; Sitona.

Linseed Oil, in mixtures for treating timber against pests, 107, 414.

Liothrips oleae, fumigation against, on olive in Spain, 385.

Liothrips urichi, utilisation of, against Clidemia hirta in Fiji, 390.

Liquidambar, discovery and eradication of Eriococcus borealis on, in California, 400.

liriodendri, Toumeyella.

Lissonota parva, parasite of Dioryctria reniculella in Quebec, 321.

Lissopimpla semipunctata, parasite of Laphygma exempta in Queensland, 307.

Lissorhoptrus bosqi, L. foveolatus compared with, 8.

Lissorhoptrus foveolatus, sp. n., on rice in Brazil, 8.

Lissorhoptrus oryzae, L. foveolatus compared with, 8.

Lissorhoptrus simplex, L. foveolatus compared with, 8.

Listroderes obliquus (in California), plants attacked by, 187; introduction of parasites from S. America against, 97. Lithobius, predacious on Prionus coriarius

in Britain, 215.

Lithocolletis blancardella, on apple in Massachusetts, 100; parasites of, 101.

Lithosia deplana (see Eilema).

Litoprosopus coachella, on Washingtonia palms in Mexico and California, 400; damaging clothing, etc., in houses, 400.

littoralis, Epiplagiops; Thisoecetrus.

litura, Prodenia. Litus, **407.**

Lixophaga diatraeae, failure to establish, against Diatraea saccharalis in Br. Guiana, 304.

Lixophaga variabilis, hosts of, in U.S.A., 84.

Lixus latro, bionomics and control of, on crucifers in Eritrea, 384.

Lizards, destroying insects, 24, 172. Lochites, Först. (see Lochitisca).

Lochitisca, n.n. for Lochites, Först., 316. Locust, Desert (see Schistocerca gre-

garia).
Locust, Red (see Nomadacris septemfasciata).

Locust Trees (see Robinia).

Locusta migratoria, tests of insecticides against, in France, 237, 238.

Locusts and Grasshoppers, in Africa, 44, 123–125, 169–171, 205, 206, 294, 366, 375, 403, 412, 426; in S. America, 108, 133, 134; in Arabia, 124; in Australia, 47, 48, 74, 216, 217, 305, 344; in Canada, 99, 166, 341; in France, 237; in India, 310–312; in Mexico, 97; in

Persia, 387, 388; in Spain and Portugal, 202, 205; in U.S.A., 13, 14, 92, 99, 157, 186, 225-228, 234, 236, 237, 333, 398; breeding centres of, 47, 170, 171, 202; seasonal breeding of, 123-125. 169-171, 206, 207, 311; migrations of, 48, 124-125, 133, 169, 170, 171, 205, 206, 207, 310-311, 403; climatic factors affecting, 109, 123–125, 170, 205, 311, 375, 398; measurement of body temperature of, 375; water exchange between air and, 375; birds destroying, 109, 312; fungi infesting, 171, 186; insect enemies of, 171, 344; destroying, 109; measures and experiments against (see also under Baits), 13, 14, 48, 92, 99, 109, 133, 134, 157, 186, 216, 217, 225–228, 234, 236, 237, 238, 294, 305, 341, 342, 403, 412, 413; cage for field sampling of, 166.

loftini, Chilo.

Loganberry, Anthonomus rubi on, in Belgium, 278; Rhynchites germanicus on, in Britain, 358.

Lonchaea aurea (splendida), on Capsicum annuum in Libya, 72.

Lonchocarpus (see Cubé).

Long-headed Flour Beetle (see Latheticus oryzae).

longicaudata, Ctenolepisma. longicornis, Paratrechina (Prenolepis).

longior, Tyrophagus.

longipennis, Scirtothrips.

longispinus, Pseudococcus (see P. adonidum).

longissima, Brontispa.

Longitarsus, on crucifers in England, 132. Longitarsus parvulus (on flax), in Eire, 208; bionomics and control of, in Italy, 268, 269.

Lophyrus pallidus (see Gilpinia).

Lophyrus pini (see Diprion).

Lophyrus polytomus (see Gilpinia).

Lophyrus rufus (see Neodiprion sertifer). Loquat (see Eriobotrya japonica).

loranthi, Farinococcus.

Loranthus, new Coccid on, in Gold Coast, 110.

lotellus, Nothris.

Lotus corniculatus, pests of, in Czechoslovakia, 278; pests of, in New York, 71; Bruchophagus in seeds of, in Sweden, 408.

Louisiana, miscellaneous pests in, 12, 84, 159, 316; beneficial insects in, 84, 316.

lounsburyi, Aspidiotiphagus.

Loxostege commixtalis, parasite of, in U.S.A., 29.

Loxostege similalis, killed by benzene hexachloride on lucerne in Oklahoma, 100.

Loxostege sticticalis, parasites and disease of, in N. America, 28, 29; on lucerne in Spain, 293; tests of insecticides against, 14, 293.

loxostegei, Meteorus.

Loxotropa tritoma, parasite of Psila rosae in Britain, 417.

Lucerne, grasshoppers damaging, in Chile, 109; pests of, in Hungary, 63; Hypera nigrirostris on, in Italy, 115; Nomophila noctuella on, in Rumania, 41; pests of, in Spain, 203, 293; pests of, in U.S.A., 24, 55, 81, 90, 99, 100, 160, 226, 228, 236, 394, 421; Lygus spp. causing seed loss of, 90; unfavourable food for Limonius californicus, 91; bees pollinating, 10, 11.

ludens, Anastrepha.

lulongaensis, Dacus armatus.

lunata, Euproctis. lunigera, Selenephera.

lurida, Leptocoris. luteola, Galerucella.

lybica, Empoasca.

Lychnis alba, Hemiptera on, in Virginia, 334.

Lycium halimifolium, Paratrioza cockerelli on, in U.S.A., 144.

lycopersicella, Keiferia.

lycopersici, Phyllocoptes.

Lyctus, lists of Australian timbers susceptible and immune to, 248, 344; measures and experiments against, 1, 107, 248, 295.

Lyctus africanus, 1.

Lyctus brunneus (in timber), in S. Africa, 1, 107; in Australia, 248, 344; in Belgian Congo, 295; Clerid predacious

Lyctus discedens, in timber in Queensland, 248; factors affecting infestation by,

248.

Lyctus hipposideros, 1.

Lyda abietis (see Cephaleia).

Lyda stellata (see Acantholyda pinivora). Lydella nigripes and virilis (see Blondelia). Lygaeonematus abietinus (pini) (see Pristiphora).

Lygidea mendax (on apple), in Nova Scotia, 392; in U.S.A., 127, 128, 271, 335; bionomics of, 271, 272; insecticides against, 272, 335; market losses caused by, 127, 128.

Lygocerus, parasite of Clausenia purpurea in Japan, 410.

Lygus, effect of species of, on seed production of beet, lucerne and guayule in U.S.A., 11, 90, 94; revision of simonyi groups of, in E. Africa, 26.

Lygus arboreus, sp. n., food-plants of, in Uganda, 26.

Lygus campestris, dusts against, on celery in Massachusetts, 102.

Lygus coffeae, on coffee in Uganda, 26. Lygus elisus, dusts against, on beet and lucerne in U.S.A., 11, 90.

Lygus ghesquierei, on coffee in Uganda, 26.

Lygus hesperus, dusts against, on lucerne and guayule in U.S.A., 11, 90, 94.

Lygus infirmus, sp. n., food-plants of, in Uganda, 26.

Lygus nairobiensis, food-plants of, in Uganda, 26.

Lygus oblineatus, in Mexico, 97; in U.S.A. 11, 90, 156, 235, 334, 373; on field crops, 11, 12, 13, 14, 90, 97, 156; causing deformation of peaches, 235, 334, 373; on weeds, 334; measures and experiments against, 11, 12, 13, 14, 90, 235, 334, 373.

Lygus palus, sp. n., food-plants of, in Uganda, 26.

Lygus ricini, sp. n., on castor, etc., in Uganda, **26.**

Lygus simonyi, L. vosseleri confused with. in Uganda, 26.

Lygus virens, sp. n., food-plants of, in Uganda, 26.

Lygus vosseleri, on cotton in Uganda, 26, 366.

Lymantria dispar, on oak in Jugoslavia, 115-117; on cork oak in Portugal. 285; on forest and shade trees in U.S.A., 36, 87, 143, 336; on cranberry in Massachusetts, 100; on fruit trees, 117; parasite of, 116; outbreak areas of, 117; cold-resistance and destruction of eggs of, 115, 116; tests and uses of DDT against, 36, 87, 100, 143, 285, 336.

Lymantria monacha (on conifers), outbreaks and distribution of, in Austria, 128, 129; in Germany, 76; disease of, 129; measures against, 129; identification of excreta of, 76.

M

Mabra russoi, sp. n., bionomics of, on sugar-cane in Dominican Republic, 29. machalana, Leptopharsa.

machili, Lepidosaphes.

Macrocentrus, parasite of Spodoptera mauritia in Tonga I., 267.

Macrocentrus ancylivorus, in U.S.A., 56, 84, 178, 185, 258, 259, 370, 397; causes of failure to establish, in Victoria, 178; value and use of, against Cydia molesta, 56, 178, 185, 370, 397; other hosts of, 84, 178; factor affecting rearing of, on Gnorimoschema operculella, 18; effect of DDT on, 56, 258, 259.

Macrocentrus delicatus, parasite of Cydia molesta in S. Carolina, 370.

Macrocentrus gifuensis, introduced parasite of Pyrausta nubilalis in Canada, 77, 391; bionomics of, 77, 78, 391; strains of, 77; technique of rearing, 77, 78.

Macrocentrus peroneae, parasite of Tortrix packardiana in Quebec, 321.

Macrodactylus subspinosus, tests DDT against, in Massachusetts, 101. Macrosiphum, on radish in Fiji, 413.

Macrosiphum avenae, dust against, on oats in Arkansas, 25; flights of, in England, 196.

Macrosiphum dirhodum, flights of, in England, 196.

Macrosiphum fragariae, flights of, in England, 196.

Macrosiphum gei, M. solanifolii (q.v.) considered identical with, 361.

Macrosiphum granarium (see M. avenae). Macrosiphum onobrychis (pisi), flights of, in England, 196; measures and experiments against, in U.S.A., 9, 12, 14, 153, 229, 325, 326-328, 421; on lucerne, 421; on peas, 12, 14, 153, 229, 325, 326-328; effect of insecticides on production of young by, 421.

Macrosiphum rosae (on rose), in Italy, 254; in Switzerland, 282; in U.S.A., 335; insecticides against, 254, 282, 355.

Macrosiphum solani (Myzus pseudosolani), HCN against, on greenhouse plants in Britain, 361; Aphid resembling, causing leaf curl of Capsicum in Bulgaria, 5; transmitting tobacco rosette viruses, 422.

Macrosiphum solanifolii, in Australia, 219, 366; in Brazil, 7; in Britain, 196, 361, 362; in U.S.A., 234, 332; parasites of, in Uruguay, 50; flights of, 196; bionomics and colour forms of, on greenhouse plants, 361, 362; on potato, 219, 234, 332, 362; relation of, to virus diseases of tobacco and potato, 7, 8, 422; transmitting virus of passion-fruit mosaic, 366; insecticides against, 234, 332; considered identical with M. gei, 361.

Macrosteles divisus, relation of, to virus disease of aster, potato and lettuce in U.S.A., 230, 231, 350; DDT against, 230, 231, 350, 351.

mactans, Latrodectus.

maculatus, Bruchus (Callosobruchus); Dermestes.

maculipennis, Dichroplus; Plutella. maculitarsus, Apanteles. maculosa, Nemorilla floralis. Madagascar Beans (see Lima Beans).

madens, Tribolium.

maderae, Calosoma.

Madras, Helopeltis theivora on nursery Eucalyptus in, 251. (See Travancore.) Magdalis spp., on elms attacked by Ceratostomella ulmi in Quebec, 320.

Magnesite, against pests of stored products, 217, 249; as carrier for dust.

218.

Magnesium Oxide, mixed with stored seed against insect pests, 12; affecting calcium arsenate in soil, 371; contents of, in calcium and magnesium limes, 329.

Magnolia, Toumeyella liriodendri on, in California, 400; Chrysomphalus dictyospermi on, in Persia, 386.

maidis, Aphis (Rhopalosiphum); Peregrinus.

Maine, Bruchus obtectus in stored beans in, 21; tests of resistance of potato varieties to leaf-roll virus in, 209.

Maize, pests of, in Australia, 138, 178, 307; pests of, in Canada, 78, 163, 164, Aphis maidis on, in Fiji, 413; Hieroglyphus nigrorepletus on, in India, 122; Busseola fusca on, in Kenya, 405; pests of, in U.S.A., 12, 13, 81, 91, 95, 96, 102, 126, 128, 136, 147, 148, 157, 158, 187, 228, 234, 332, 370, 374, 401; unusual injury to, by Heliothis armigera, 157; effect of growth of, on establishment of Pyrausta nubilalis, treatment of ears of, against pests, 13, 126, 158, 370; effect of insecticides on, 148, 149, 228, 234, 236, 332, 338; planting of, with rape against Entomoscelis adonidis, 41.

Maize (Stored), pests in, and their control, 2, 19, 52, 68, 85, 210, 367, 369; sorption of fumigants by, 85; effects of insecticide treatments on germination of, 3,

19, 331.

Maize Meal, in baits, 53, 350, 372.

major, Penthaleus.

malacellus, Crambus.

Malacosoma, concentrated sprays against, in U.S.A., 36.

Malacosoma americana, DDT against, on

shade trees in U.S.A., 143.

Malacosoma disstria, in Minnesota, 243; in New Brunswick, 208; factors affecting diapause and hatching of eggs of, 243, 244.

Malacosoma neustria, DDT against, in

Turkey, 352.

Maldive Is., new mite parasitising Orthoptera in, 427.

malella, Nepticula.

mali, Aphelinus; Eriophyes pyri; Leptothrips.

malifoliae, auct., Yezabura (see Anuraphis roseus).

maligna, Empoasca.

malinellus, Hyponomeuta padellus; Heterocordylus; Pseudaphycus.

malivagrans, Vasates (see V. schlechtendali).

malus, Hemisarcoptes,

Malva spp., Earias insulana on, in Punjab, 312; Podagrica spp. on, in Rumania, 41.

malvae, Podagrica.

Malvastrum tricuspidatum, Earias spp. on, in Puniab, 312.

Mamestra brassicae, tests of thiocyanoacetates against, on cabbage in Italy, 253.

Mammea americana (Mamey), experiments with preparations of, as insecticides, 51, 52.

Mandevilla, Euploea core corinna on, in New South Wales, 344.

Manganese Oxide, not affecting calcium arsenate in soil, 371.

Mangifera indica (see Mango).

mangiferae, Aceria; Eriophyes; Oxypleurites.

Mango, pests of, in Egypt, 409, 411; pests of, in Hawaii, 80, 249; pests of, in India, 110, 123; Anastrepha ludens on, in Mexico, 20; A. mombinprae-optans on, in Porto Rico, 54; new Mirids on, in Uganda, 26; smoke from burning leaves of, repellent to A. ludens, 20.

Manihot utilissima (see Cassava).

Manitoba, Hylemyia antiqua on onion in, 89, 156.

Maori Mite (see Phyllocoptruta oleivorus). Maple, pests of, in Quebec, 319.

Maple, Soft (see Acer glabrum).

Marasmia venilialis (on rice, etc.), in Fiji, 267, 390; in India, 267; parasites of, 267; sprays against, 267.

margaritosa, Peridroma (see P. saucia). marginata, Pennisetia.

marginella, Dichomeris.

marginipennis, Hercothrips.

mariana, Eulia.

maritimus, Pseudococcus.

maroccanus, Dociostaurus.

marshalli, Chaetonodexodes.

martinii, Neodiscodes.

Maryland, insects attacking fruit trees in, 152, 157, 339; Lepidoptera on maize in, 95, 96; Macrosiphum onobrychis on peas in, 325; Mezium americanum in, 98; new parasite of Phenacoccus in, 315.

Massachusetts, Armadillidium spp. in greenhouses in, 159; miscellaneous pests in, 100-103, 336; beneficial insects in, 28, 101.

Mastrus carpocapsae, introduced into New Zealand against Cydia pomonella,

maulei, Crypturgus (see C. pusillus). mauritanicus, Tenebroides (Trogosita).

mauritia, Spodoptera.

Mauritius, Gonipterus scutellatus on Eucalyptus in, 125; miscellaneous pests in, 209, 210; natural enemies and biological control of insects in, 125, 209, 210; proposed biological control of Cordia macrostachya in, 424; Celosterna scabrator not occurring in, 317. maxillosum, Psalidium.

Mayetiola destructor, on wheat in Ohio,

136.

MB 320, 131.

Mealybug, Baker's (see Pseudococcus maritimus).

Meat, determination of DDT in, 393; (dried), pests in, in Queensland, 309, 310.

Medicago falcata, Bruchophagus in seeds of, in Sweden, 408.

Medicago sativa (see Lucerne).

Mediterranean Fruit Fly (see *Ceratitis* capitata).

Megachile, pollinating lucerne in Utah,

Megacyllene robiniae, spray against, on Robinia in U.S.A., 144.

megalopygus, Dysdercus (see D. cingu-

Megaselia rufipes, parasite of Leptinotarsa decemlineata in France, 349; bionomics of, 349.

Megaselia scalaris, parasite of Desmeocraera varia in Ivory Coast, 390; bionomics of, 390.

Megastes grandalis, on sweet potato in Trinidad, 319.

Megastigmus kuntzei, sp. n., bionomics of, on juniper in Poland and Rumania, 267, 268.

Megastigmus pistaciae, bionomics of, on Pistacia in Persia, 386.

Megastigmus wachtli, M. kuntzei compared with, 267.

Meigenia mutabilis, parasite of Entomoscelis adonidis in Rumania, 40.

Melanagromyza (see Agromyza).

Melanaspis, characters of, 401.

Melanaspis inoptinata, on Pistacia and pear in Persia, 386.

Melanaspis obscura, elimination of, in California, 189.

melanocephala, Diabrotica.

melanocephalus, Exochomus. melanogaster, Drosophila.

melanopa, Lema.

Melanoplus, in New Hampshire, 234; baits against, 157, 398; other uses and tests of insecticides against, 92, 99, 157, 226-228, 234, 236, 237, 341, 342. Melanoplus bivittatus, in U.S.A., 99, 236.

Melanoplus differentialis, in U.S.A., 92, 99, 157, 226-228, 236, 237; defoliating

apple, 157.

Melanoplus femur-rubrum, in U.S.A., 92, 157, 226-228, 236, 237; defoliating apple, 157.

Melanoplus mexicanus devastator,

California, 398.

Melanoplus mexicanus mexicanus, in Canada, 99, 341; in U.S.A., 99, 236. melanostictus, Catantops.

Melanotus rufipes, predacious on Prionus coriarius in Britain, 215.

Melasoma lapponica var. interrupta, spray against, on willow in U.S.A., 143.

Melasoma lineatopunctata, dust against, on willow in New Hampshire, 234.

Melcha ornatipennis, parasite of Scirpophaga nivella in India and Burma, 119; bionomics of, 119.

meles, Hypera.

Melia azadirachta, Anychus orientalis on,

in Egypt, 412.

Melilotus (Sweet Clover), weevils on, in U.S.A., 55, 81, 187; experiment with Limonius californicus and, 91; doubtful relation to, of bugs causing cat-facing of peaches, 334; question of DDT residues on, 81.

Melilotus alba, Hypera brunneipennis developing on variety of, 55.

Melilotus indica, weevils on, in U.S.A., 55, 187.

Melilotus officinalis, Hypera brunneipennis developing on, 55.

Melittia cucurbitae (in U.S.A.), susceptibility of Cucurbita spp. to, 101; DDT analogue against, 14.

Melittobia indica (see Syntomosphyrum).

mellitor, Bracon (Microbracon).

Melolontha melolontha, benzene hexachloride against, in Switzerland and France, 279, 280.

Melon, Empoasca terrae-reginae on, in W. Australia, 74; Diaphania on, in Peru, 134; pests of, in U.S.A., 101, 225, 331, 398; Eutettix tenellus transmitting curly-top to, 398; insecticides and injury to, 134, 224, 225, 331; benzene hexachloride not affecting flavour of, 235; DDT mixed with stored seed of, against insects, 19.

Melon Aphis (see Aphis gossypii).

mendax, Lygidea.

meracus, Pseudaphycus.

mercator, Oryzaephilus surinamensis.

Mercuric Chloride, toxicity of, to *Agriotes*, 286, 287; solutions of, against *Hylemyia* spp., 90, 102, 283; toxic to bacteria causing onion-bulb decay, 283.

Mercurous Chloride, tests and uses of, against *Hylemyia* spp., 90, 102, 174, 222, 223, 270, 282; ineffective against *Psila rosae*, 377; toxic to bacteria causing onion-bulb decay, 283; effect of, on germination and growth of onion and leek, 90, 270.

Mercury Fluoride, repellency of compounds of, to Kalotermes brevis, 232,

233.

meridionalis, Otiorrhynchus (Brachy-rhinus).

meritorius, Pseudaphycus.

Mescinia peruella, dusts ineffective against, on cotton in Peru, 134.

Mesochorus perniciosus, parasite of Meteorus loxostegei in U.S.A., 29.

Mesoleius, parasite of sawflies in Italy, 31, 193.

Mesoleius tenthredinis, parasite of Gilpinia hercyniae in Quebec, 320.

Mesopeltita, n.n. (Mesopeltis, Masi), 316. Metagonistylum minense (parasite of Diatraea saccharalis), established in Br. Guiana, 304; introduction of, into Porto Rico, 54.

Metamasius hemipterus, on sugar-cane in

Br. Guiana, 304.

Metaphycus helvolus (parasite of Saissetia oleae), unlikely to become established

in S. Australia, 216.

Metarrhizium anisopliae, introduction of, against Tomaspis flavilatera in Br. Guiana, 303; infesting T. saccharina in Trinidad, 303, 319; infesting Homoeosoma electellum in U.S.A., 84.

Metatetranychus ulmi, auct. (see Para-

tetranychus pilosus).

Meteorus loxostegei, bionomics and distribution of, in N. America, 28, 29; rearing of, 29.

Meteorus narangae, sp. n., parasite of Naranga aenescens in Formosa, 208.

Meteorus trachynotus, parasite of Tortrix packardiana in Quebec, 321.

β-Methallyl Chloride, mixtures of other fumigants with, 85.

Methoxy-DDT (Dimethoxydiphenyltrichlorethane), against Coleoptera, 14, 230; against Lepidoptera, 14, 230, 341, 403; ineffective against Lygus oblineatus, 14; against Melanoplus spp., 237; toxicity of, to bees, 222; insecticidal action not related to liberation of HC1 by, 274, 275; dusting with, 14, 230, 237, 403; spraying with, 341; mixed with stored wheat, 14.

Methyl Bromide, tests and use of, against pests on plants and planting material, 79, 187, 292, 296, 308, 353, 397; and injury to plants, 187, 400; fumigation of fruits and flowers with, 79, 80, 186, 187; against pests in other stored products, 105, 182, 217, 245, 355; sorption of, by wheat, 39; residues of, in flour, 39; fumigation of grain bags in barges with, 355–357; fumigation of beans and harvesting equipment with, against *Epilachna varivestis*, 396; equipment for using, 105, 308, 356.

Methyl Chloride, use of, for producing

aerosols, 95, 230, 325, 327.

Methyl Formate, unsuitable against pests in stored grain, 30.

Methyl Isopropenyl Ketone, test of, as a fumigant, 85.

Methyl Pentachlorphenyl Ether, 224.

Methylated Naphthalene, as a solvent for DDT, 33. (See Velsicol AR-60.)

Methylated Spirit, against Pseudococcus kenyae, 404.

Metopolophium dirhodum (see Macrosiphum).

Metroxylon vitiensis, Cerataphis lataniae on, in Fiji, 413.

Mexican Bean Beetle (see Epilachna varivestis).

Mexican Fruitfly (see Anastrepha ludens). mexicanus, Melanoplus; Pityophthorus.

Mexico, Anastrepha spp. in, 20, 148, 221; pests of guayule in, 96, 97; plant-quarantine restrictions in, 105; spread of pests into U.S.A. from, 146, 147, 400, 401; measures to prevent introduction of Coccids into U.S.A. on limes from, 79; new insects from, found in quarantine in U.S.A., 316, 401.

Mezium americanum, DDT against, in bakery in Maryland, 98.

Mica, as a carrier for dusts, 329; pyrophyllite resembling, 341.

Mice, destroying *Ephestia elutella*, **212**. Michigan, pests and disease of vegetables in, **350**.

Microbracon (see Bracon).

Microcerotermes parvulus, attacking groundnut pods in Senegal, 250.

Microgaster ecdytolophae, parasite of Walshia amorphella in Oklahoma, 9.

Microphanurus, parasite of Antestia in Uganda, 367.

Microplectron fuscipenne Dahl-(see bominus).

Microtermes parvulus (see Microcerotermes).

migratoria, Locusta.

miliaris, Asterolecanium.

Milk, colorimetric determination of DDT in, 393; (condensed), new mite infesting, 114.
Milkweed Bug, Large (see Oncopeltus

fasciatus).

Millettia pachycarpa, insecticidal action and constituents of, 298, 299.

Millettia pulchra, insecticidal properties of, 299.

Mimosa bracaatinga, Xyleborus fornicatus on, in Ceylon, 251.

Mimusops spp., Desmeocraera varia on, in S. Africa, 389.

minense, Metagonistylum.

Mineola vaccinii, dusts against, on cranberry in Massachusetts, 100.

minimum, Monomorium.

Minnesota, Malacosoma disstria in, 243. Minthea rugicollis, in Belgian Congo, 1. minuta, Hoplocampa; Womersleyia.

minutulus, Scymnus. minutum, Trichogramma.

minutus, Dinoderus; Laemophloeus; Orius; Rhynchites.

Miscanthus japonicus, Oregma lanigera on, in Fiji, 413.

Mississippi, Mirids on cotton in, 156; maize pests in, 157, 158; legislation against Pantomorus in, 184.

Missouri, Cydia molesta in, 186; insects on Helianthus in, 84, 150.

missouriensis, Horismenus.

Mites, review of N. American Eriophyid, 82, 83; list of, recorded on cotton, 388; technique for testing acaricides on, 343; attacking insects, 62, 70, 172, 212, 392, 427; attacking other mites, 82, 343; natural enemies of, 82, 282, 288, 343, 393, 426, 427; classification and new species of, 80, 83, 114, 409, 411, 412, 427.

mixta, Orgyia.

Mocis repanda, on sugar-cane in Br. Guiana, 304.

Molasses, in baits, 62, 74, 132, 158, 261, 294, 307; in mopping mixture against Anthonomus grandis, 370; in sprays, 90, 123, 404

molefaciens, Pogonomyrmex barbatus.

molesta, Cydia (Grapholitha, Laspeyresia); Solenopsis.

molitor, Tenebrio.

mombinpraeoptans, Anastrepha.

Momordica, Dacus bivittatus on, in W. Africa, 176.

monacha, Corythaica; Lymantria.

Monarthropalpus buxi, DDT against, on box in U.S.A., 143.

Monellia (see Myzocallis).

Monochamus scutellatus, bionomics of, in Quebec, 320.

Monodontomerus obsoletus, parasite and hyperparasite of Diprion pini Germany, 239.

Monomorium minimum, effect of sprays on, destroying Cydia pomonella in W. Virginia, 220.

Montana, parasites and hyperparasites of Loxostege sticticalis in, 29.

Montezuma speciosissima, 52. montrouzieri, Cryptolaemus. monuste, Ascia (Pieris).

mori, Bombyx.

Mormoniella vitripennis, hyperparasite of Diprion pini in Germany, 239; effects of temperature on, 427.

Morocco, Schistocerca gregaria in, 205, 206, 207; digest of plant-quarantine restrictions in, 105.

Mottle Umber (see Erannis defoliaria). Mountain Ash (see Sorbus aucuparia).

Mulberry, Hyphantria cunea on, in Hungary, 254; Pseudococcus filamentosus on, in Punjab, 123.

Mullein (see Verbascum thapsus).

multistriatus, Scolytus. mundus, Pseudaphycus.

Murgantia histrionica (Harlequin Cabbage Bug), tests of insecticides against, in U.S.A., 13, 333.

murinus, Lacon.

Musca domestica, tests of insecticides on, 223, 224, 328, 377; used for tests of DDT in milk and butter, 379.

muscae, Aphaereta. Muscardine Fungus, Green (see Metarr-

hizium anisopliae).

Muscina stabulans, parasite of Leptinotarsa decemlineata in France, 349. musculus, Anthonomus.

Mushroom Flies (see Sciara).

Mushrooms, Sciara on, in Britain, 284; pests of, in California, 187; pests of, in New South Wales, 365; effect of benzene hexachloride on, 365.

Mustard, Aphid on, in Arkansas, 25; Athalia rosae on, in Italy, 192.

mutabilis, Meigenia.

mycophagus, Caloglyphus.

Myelophilus, revision of Palaearctic species of, 32.

Myelophilus piniperda var. rubripennis, bionomics of, in *Pinus pinea* in Italy,

Myiocnema, hosts and systematic position of, 109.

myosotidis, Nematus (Pteronidea). Myrobalans (see Terminalia).

myrsusalis, Rhodoneura. mytilaspidis, Aphytis.

Mytilococcus (see Lepidosaphes).

Myzocallis, lead arsenate increasing infestation of pecan by, in Texas, 152. Myzocallis annulata, biological control of, on oak in Tasmania, 70.

Myzocallis kahawaluokalani, on Lagers-

troemia in California, 400.

Myzocallis ulmifolii, spray against, on elm in U.S.A., 143.

Myzus cerasi, dust against, on cherry in New Hampshire, 234.

Myzus circumflexus, on Cinchona in Porto Rico, 53.

Myzus persicae, in Australia, 219, 366; in Brazil, 7, 8; in Britain, 196; in Bulgaria, 4, 5; in Czechoslovakia, 355; in Fiji, 413; in U.S.A., 24, 79, 92, 234; in Uruguay, 50; flights of, 196; natural enemies of, 50, 93; transmitting beet mosaic, 355; virus transmitted by, in *Passiflora edulis*, 366; on peach, 5, 92; transmitting peach mosaic, 79, 93; on potato, 219, 234; relation of, to virus diseases of solanaceous plants, 4, 5, 7, 8, 422; on radish, 413; on spinach, 24; measures and experiments against, 5, 24, 92, 228, 234.

Myzus porosus, toxicity to, of esters of

phosphorus acids, 228.

Myzus pseudosolani (see Macrosiphum solani).

Myzus solani (see Macrosiphum).

N

Nabis ferus, little affected by DDT in California, 91; predacious on Entomoscelis adonidis in Rumania, 40.

Nacoleia diemenalis, spray against, on beans in Fiji, 267.

nadinus, Tyroglyphus.

naevana, Rhopobota.

nairobiensis, Lygus.

nana, Trichogrammatoidea.

Naphtha, as a solvent for DDT, 28, 46, 47, 69, 178, 217, 218, 219, 263; types of, as solvents for pentachlorphenol, 107.

Naphthalene, soil treatment with, against Coleoptera, 137; tests of treatments with, against Hylemvia spp., 174, 283; against Onychiurus fimetarius, in spray against Tetranychus, 103; and synthesis and naphthylamine, 283; insecticidal action of derivatives of, 275, 276.

Naphthalene, Chlorinated, 275: ment of timber with, against Anobium and Kalotermes brouni, 245, 414, 415. β -Naphthol, thiocyanoacetic ester of, as

an insecticide, 253, 254.

Naphthylamine, experiments with, against Hylemyia antiqua, 283; and naphthalene, 283.

napi, Ceuthorrhynchus.

Naranga aenescens, new parasite of, in Formosa, 208.

narangae, Meteorus.

Narcissus, pests of, in Britain, 32; Agasphaerops nigra not feeding on, 142. nasalis, Alophora; Cermatulus.

nasturtii, Contarinia.

Nasturtium, Aphids on, in U.S.A., 335, 421.

Nasutitermes (see Eutermes).

Nebraska, Lepidoptera on maize in, 157; Melanoplus spp. in, 236; pests of solanaceous plants in, 56, 144.

nebulosa, Cassida.

Necrobia rufipes, in salt fish in Uganda,

Nectarine, Rhynchites coeruleus on, in Britain, 358; Cydia molesta on, in France, 238; pests of, in New South Wales, 247, 306.

Nectria diploa, infesting Chrysomphalus ficus in Florida, 83.

negligens, Acritocera.

Nematodes, infesting Psila rosae Britain, 417.

Nematus myosotidis, bionomics of, on clovers in Italy, 30.

Nematus ribesii, sprays against, on gooseberry in U.S.A., 335.

Nemeritis canescens, parasite of Ephestia elutella in Britain, 212.

Nemeritis pastranai, sp. n., parasite of Rhyacionia buoliana in Argentina, 109. nemoralis, Neurotoma.

Nemorilla floralis, parasite of Homoeosoma electellum in Cuba, 83; tests of toxicity of DDT to, in Ohio, 258, 259.

Nemorilla floralis var. maculosa, parasite of Harmologa fumiferana in Quebec, 321.

Nemorilla pyste, parasite of Tortrix packardiana in Quebec, 321.

nemorum, Anthocoris; Phyllotreta.

nenuphar, Conotrachelus.

Neoatta (see Atta).

Neocopidosoma, Blanch. (see Noblanchardia).

Neocopidosoma, Ishii, 316.

Neodiprion lecontei, insecticides against, on pine in U.S.A., 143, 234.

Neodiprion sertifer, characters of excreta of, on pine in Germany, 76.

Neodiscodes martinii, parasite of Coccids in Gold Coast, 111.

Neolygus communis novascotiensis, on apple in Nova Scotia, 392.

Neopristomerus appalachianus var. dorsocastaneus, Meteorus loxostegei destroying, in Loxostege sticticalis, 29.

Nephrolepis exaltata, insecticides injuring, 282.

Nepticula malella, bionomics of, on apple in Britain, 359.

Nepticula promissa, on Pistacia in Persia, 396.

nerii, Aphis.

Nerium (Oleander), Euploea core corinna on, in New South Wales, 344; Saissetia oleae on, in Persia, 385.

Netherlands Indies, *Xyleborus fornicatus* in, **251**; new hyperparasite of *Pseudococcus lepelleyi* in, **109**.

Neurocolpus nubilus, on cotton and Croton capitatus in U.S.A., 156.

Neurotoma nemoralis, bionomics of, on stone fruits in Poland, 254, 255. neustria, Malacosoma.

New Brunswick, pests of forest and shade trees in, 35, 207, 208, 334; potato Aphids in, 209; Pyrausta nubilalis in, 77, 167; beneficial insects and biological control in, 77, 207, 208.

New Caledonia, Aulacophora similis on cucurbits in, 267.

New Hampshire, miscellaneous pests in, 234.

New Hebrides, Brontispa longissima on coconut in, 390; Aulacophora similis on cucurbits in, 267, 390.

New Jersey, pests of blueberry and cranberry in, 137; Cydia molesta in, 152, 341; pests in nurseries in, 136, 152; miscellaneous pests in, 150, 301; beneficial insects in, 302, 315.

New Mexico, Meteorus loxostegei in, 28; miscellaneous pests in, 126.

New South Wales, fruit pests in, 46, 108, 181, 247, 248, 306, 344, 345, 365; grasshoppers in, 48, 216, 305, 344; Heteronychus sanctae-helenae in, 73, 138; Iridomyrmex spp. in, 247, 305; Lyctus brunneus in, 344; miscellaneous pests in, 248, 305, 344; pests of mushrooms in, 365; pests of stored products in, 46, 68, 217; vegetable pests in, 46, 73, 216, 217, 344, 345; insects and virus diseases of plants in, 46, 344, 365; natural enemies and biological control of insects in, 108, 216, 247, 344; biological control of weeds in, 181, 216.

New York, pests of *Lotus corniculatus* in, 71; miscellaneous pests in, 301, 421;

orchard pests in, 56-59, 127, 271, 324, 334, 338; *Erythroneura* on vines in, 255.

New Zealand, cheese mites in, 244, 416; miscellaneous pests in, 244, 309, 427; Anobium punctatum and Kalotermes brouni in timber in, 414-416; natural enemies and biological control of Arthropods in, 244, 427; biological control of noxious plants in, 244; Pteromalus puparum introduced into Tasmania from, 70.

Newfoundland, digest of plant-quarantine restrictions in, 105.

newsteadi, Lepidosaphes.

Newsteadia wacri, sp. n., food-plants of, in Gold Coast, 110.

ni, Trichoplusia.

Nicandra physaloides, Aphids transmitting tobacco rosette to, 422.

Nicotiana, greenhouse fumigation against pests of, in Britain, 361; differences in susceptibility of species of, to tobacco virus, 7.

Nicotiana glutinosa, used in experiments with Plusia gamma and tobacco mosaic,

Nicotiana tabacum (see Tobacco).

Nicotine, against Aphids, 13, 98, 199, 228, 314, 326, 327, 332, 371; varying effectiveness of, against Jassids and Mirids, 176, 383, 386, 395; not increasing effect of oil on Parlatoria oleae, 17; against Psylla pyricola, 334, 335; against Coleoptera, 62, 204, 279; effect of temperature on toxicity of, to Tribolium castaneum, 380; against Contarinia nasturtii, 286; against Cydia pomonella, 57, 58, 59, 93, 100, 126, 127, 219, 225, 258, 323; against other Lepidoptera, 102, 402, 409; against mites, 13, 149, 239; against sawflies, 279, 286; against Scirtothrips signipennis, 306; applied by bloweratomisers, 36; dusting with, 204, 306, 326, 327, 332, 371, 395, 402; carriers for, 395, 402; oil-impregnated dusts containing, 57, 59, 326; fumigation with vaporised, 314; fixed compounds of, 57, 58, 59, 98, 100, 102, 126, 127, 149, 258, 323, 335, 395, 402; formulae containing, 17, 58, 59, 98, 102, 126, 127, 149, 150, 199, 239, 258, 306, 326, 327, 332, 334, 371, 395, 402, 409; and arsenicals, 59, 100, 225, 371; and benzene hexachloride, 326, 327; and DDT, 57, 58, 98, 127, 150, 306, 371, 395; and oil emulsions, 98, 126, 149, 225, 258; and soap, 62, 199, 286; optical isomers of, 297, 298; other insecticides compared with, 13, 93, 228, 297, 298, 332, 402.

Nicotine Sulphate, against Aphids, 46, 50, 52, 102, 152, 180, 192, 218, 247, 334; against Jassids, 127, 325, 326; against other Hemiptera, 75, 272, 334, 335; against Apion damryi, 31; against Collembola, 305; against Cydia pomonella, 58, 59, 257, 323; against other Lepidoptera, 4, 52, 159, 218, 269; against Diptera, 46, 47, 73, 248; ineffective against grasshoppers, 157; ineffective against grasshoppers, 157; ineffective against mites, 246, 309; against thrips, 53, 102; in bait-sprays, 46, 47, 248; dusting with, 218, 272; formulae containing, 46, 50, 52, 53, 59, 75, 159, 180, 246, 248, 257, 272, 325, 326; and DDT, 58, 218; and lead arsenate, 58, 59, 152, 257, 325; and oil emulsions, 46, 47, 53, 58, 59, 73, 159, 180, 257, 272, 309, 325, 326, 334; and soap, 4, 50, 52, 75, 102, 218, 247, 305, 334; Toxaphene compatible with, 224.

Nigeria, cacao mealybug in, 110; Dacus spp. in, 176; Schistocerca gregaria in, 206, 207; parasite of Dysdercus in, 27.

nigeriensis, Alophora (see A. nasalis).

nigra, Agasphaerops.

nigricana, Ĉydia (Laspeyresia). nigricans, Pityophthorus.

nigricans, Pityophthorus, nigricornis, Oecanthus.

nigripes, Blondelia (Lydella); Phyllotreta; Pleurotropis.

nigrirostris, Hypera (Phytonomus).

nigritulus, Cybocephalus.

nigriventris, Anagrus armatus.

nigrofasciatus, Dysdercus.

nigronervosa, Pentalonia.

nigrorepletus, Hieroglyphus.

Nigrospora oryzae, associated with Pediculoides graminum on wheat in Spain,

202.
Nilotaspis halli, on almond, peach and related plants in California, 156, 185, 398; Nitidulid predacious on, 156; measures against, 185.

nitens, Anaphoidea.

nitidalis, Diaphania.

nitidipennis, Cordylomera.

Nitriles, 12, 20, 38, 62, 87, 93.

nivella, Scirpophaga.

njalensis, Pseudococcus.

Noblanchardia, n.n. for Neocopidosoma, Blanch., 316.

noctilio, Sirex.

noctuella, Nomophila.

Nomadacris septemfasciata, outbreak of, in Africa, 169-171, 294; breeding and migrations of, 170, 171; uses of benzene hexachloride against, 294.

Nomia, pollinating lucerne in Utah, 10,

Nomophila noctuella, bionomics and control of, on clover and lucerne in Rumania, 41.

Nornicotine, tests of toxicity of, to Arthropods and mammals, 13, 93, 297, 298; optical isomers of, 297.

norvegicus, Calocoris.

Notamorphota timocraticae, gen. et sp. n., parasite of Timocratica haywardi in Argentina, 109.

Nothris lotellus, insecticides against, on

lucerne in Spain, 293.

Nova Scotia, orchard pests in, 270, 271, 392, 393; failure to establish European parasite of *Pyrausta nubilalis* in, 77.

novascotiensis, Neolygus communis.

Novius cardinalis (see Rodolia).

nubilalis, Pyrausta.

nubilus, Neurocolpus.

numidicus, Crypturgus.

Nun Moth (see Lymantria monacha).

nuperus, Bracon (Microbracon).

nursei, Goryphus.

Nutmegs, Stored, bibliography of pests of, 272.

Nyasaland, Aphids and virus diseases of plants in, 422, 426; miscellaneous pests in, 44, 45, 176; beneficial insects in, 27, 44, 45; plant pest legislation in, 32, 105.

Nygmia phaeorrhoea (see Euproctis chrys-

orrhoea). nymphaeae, Rhopalosiphum.

nymphaeata, Nymphula.

Nymphalis antiopa (Spiny Elm Caterpillar), benzene hexachloride against, in New Hampshire, 234; bionomics and control of, on rice in Lombardy, 346, 347.

Nymphula nymphaeata, bionomics and control of, on rice in Lombardy,

346, 347.

Nysius vinitor, insecticides ineffective against, on potato in W. Australia, 74.

0

Oak, Cosmia spp. on, in Belgium, 277; insects on, in Britain, 215, 358, 359; Antheraea pernyi reared on, in Italy, 67; Lymantria dispar on, in Jugoslavia, 115, 116, 117; pests of, in Tasmania, 70; pests of, in U.S.A., 14, 301, 302, 400.

Oak, Cork (see *Quercus ilex* var. *suber*). Oak Aphis (see *Myzocallis annulata*). Oak Flea-beetle (see *Haltica quercetorum*). Oak-wood Flour, as carrier for dinitro dust, **288**.

Oats, Oscinella frit on, in Belgium, 286; Lema melanopa on, in Italy, 37; Lachnosterna damaging, in Ontario, 164; Aphids on, in U.S.A. and Uruguay, 25, 49; experiments with wireworms in relation to, in U.S.A. and Britain, 91, 183, 381, 382; effect of benzene hexachloride on germination and growth of, 13.

Oats (Stored), DDT mixed with, against pests, 19; effect of, as food for

Ephestia elutella, 265. oblineatus, Lygus.

obliquus, Listroderes.

obliterata, Angitia (Inareolata).

oblongus, Phyllobius.

obovatus, Brevipalpus (Tenuipalpus).

obscura, Melanaspis (Chrysomphalus).

obscurus, Agriotes.

obsoleta, Heliothis (see H. armigera).

obsoletus, Monodontomerus.

obsoletus, auct., Acanthoscelides (see Bruchus obtectus).

obtectus, Bruchus (Acanthoscelides).

occidentalis, Copa; Frankliniella; Pogonomyrmex.

ocellana, Spilonota.

ocellaris, Dinoderus.

ochraceipennis, Trimerotropis.

ochropa, Arge.

1, 2, 4, 5, 6, 7, 8, 8—Octachlor—4, 7—methano—3a, 4, 7, 7a—tetrahydroindane (see Chlordane).

Odynerus tigris, bionomics of, destroying Lepidopterous larvae in Ontario, 167.

Oecanthus nigricornis, dust against, on tobacco in S. Carolina, 372.

Oechalia consocialis, predacious on Lepidopterous larvae in New South Wales, 108.

Oedaleonotus enigma, weather factors affecting, in California, 398.

Oedaleus, on Eleusine in Uganda, 367.

Oedaleus decorus, crops damaged by, in Persia, 387, 388.

Oeonistis quadra, characters of excreta of, on conifers in Germany, 76.

oestracea, Trichopsidea.

Ohio, cereal pests in, 136; greenhouse pests in, 289, 290; *Cydia molesta* on peach in, 23; vegetable pests in, 12, 13; beneficial insects in, 23, 258.

Oil Emulsions, against Aphids, 130, 192, 247, 276, 287, 334; against Coccids, 17, 79, 160, 185, 189, 197, 198, 282, 293, 385, 387; not decreasing resistance of *Aonidiella aurantii* to HCN, 161; against other Hemiptera, 272, 282, 284, 287, 335; against Coleoptera, 276, 278, 293; against Diptera,

46, 73, 90, 284; against Cydia pomonella, 179, 180, 218, 225, 323; against other Lepidoptera, 159, 273; ineffective other Lepidoptera, 159, 273; ineffective against locusts, 133; against mites, 49, 59, 82, 101, 129, 130, 131, 155, 180, 218, 247, 258, 276, 282, 287, 309, 360, 361, 393; against thrips, 53, 282; Coniopteryx tineiformis not affected by, 359; tests of ovicidal action of, 273; types of oil for, 17, 58, 79, 81, 90, 98, 101, 126, 127, 131, 133, 149, 155, 158 101, 126, 127, 131, 133, 149, 155, 158, 159, 160, 187, 189, 197, 225, 227, 258, 272, 273, 278, 282, 284, 287, 293, 331; emulsifiers for, other than soap, 160, 272; formulae for, 17, 19, 49, 53, 59, 79, 90, 98, 101, 126, 127, 131, 133, 158, 159, 160, 179, 180, 185, 187, 189, 197, 218, 225, 227, 247, 254, 256, 257, 258, 272, 276, 278, 284, 287, 293, 309, 335, 360; formulae containing, 19, 46, 53, 59, 90, 98, 126, 131, 149, 150, 151, 155, 158, 159, 160, 179, 180, 187, 189, 218, 225, 227, 245, 254, 256, 257, 258, 285, 287, 293, 309, 331, 335, 361; and calcium arsenate, 42; and calcium chloride, 361; and copper compounds, 42, 90, 131, 293, 332; and cryolite, 19, 149, 150, 151; solutions of DDT in oil for, 35, 131, 150, 284, 285, 309, 337; mixtures of DDT and 58, 69, 81, 98, 126, 127, 150, 158, 179, 180, 187, 189, 225, 256, 257, 258, 276, 293, 323, 327; reducing toxicity of DDT to Cydia pomonella, 256; and dinitro compounds, 130, 272, 273, 274, 276, 294, 3274 276, 284, 287; and other synthetic organic insecticides, 13, 149, 155, 224, 225, 227, 228, 254, 293, 321, 331, 335; and ferric dimethyldithiocarbamate, 131; and lead arsenate, 58, 59, 98, 126, 149, 150, 151, 179, 180, 218, 225, 245, 246, 257; and nicotine, 46, 53, 58, 59, 73, 98, 126, 149, 150, 159, 180, 225, 258, 272, 309, 334; and rotenone insecticides, 17, 160, 189, 282, 335; and sodium chloride, 361; and sulphur, 225; and tetramethylthiuramdisulphide, 131; and zinc fluoarsenate, 180; and injury to plants, 58, 127, 132, 152, 179, 180, 218, 327, 361; risk of damage to plants by dinitro compound following, 288; treatment of packed limes with, 79.

Oils, spraying with, against locusts, 48; treatment of rice-fields with, against Nymphula nymphaeata, 346, 347; effect of temperature on toxicity of, to Tribolium castaneum, 380; tests on Parlatoria oleae of measured amounts of, with and without insecticides, 16, 17; in baits, 333; dusts impregnated

with, 57, 58, 59, 145, 229, 326, 327, 370; as solvents or supplements for DDT sprays or aerosols, 34, 36, 37, 87, 144, 147, 230, 327, 341, 342, 380, 398; as solvents for other insecticides in sprays, 99, 147, 224, 341, 342, 398; duration of toxicity of films of pyrethrum in, 168; ants' nests treated with chlordane in, and supplementary materials, treatment of maize ears with, against Noctuids, 126, 158, 370; in mixtures for treating logs and timber against insects, 35, 103, 107, 414, 415.

Oils, Vegetable, in mixtures for treating timber against pests, 107, 414, 415; emulsified solution of polychlorcyclane sulphide in, 278; repellency of derivatives of, to Kalotermes brevis, (See Groundnut, Soybean and

Sunflower-seed Oils.)

Oklahoma, pests of Amorpha and their parasites in, 9; Mirids on cotton in, 183, 184; miscellaneous pests in, 94, 99, 100, 154; Rhyacionia frustrana in, 158, 388.

oleae, Dacus; Liothrips; Parlatoria; Saissetia.

Oleander (see Nerium).

oleellus, Pravs.

Oleic Acid, uses of, in sprays, 98, 149. oleivorus, Phyllocoptruta.

Oligonychus ulmi, auct. (see Paratetranychus pilosus).

Oligota, predacious on Tetranychus carpini in Switzerland, 282.

Olive, 385; Dacus oleae on, in Greece, 292; Prays oleellus on, in Italy, 141, 193, 194; Chrysomphalus dictyospermi on, in Persia, 386; Teleonemia australis on, in S. Rhodesia, 74, 75; pests of, in Spain, 201, 385; Coccids on, in U.S.A., 16, 187, 188, 342, 399, 400; not attacked by Quadraspidiotus perniciosus, 64; bait-sprays injuring, 75.

Olive Moth (see Prays oleellus).

Olive Scale (see Parlatoria oleae).

omissa, Euphorocera.

Omorgus phthorimaeae, attempted introduction of, into Australia against Gnorimoschema operculella, 216.

Oncoba echinata, Poëphilina flaveola on, in Ivory Coast, 67.

Oncopeltus fasciatus, tests of toxicity of insecticides to, 13, 22.

Onion, Hylemyia spp. on, in Britain, 173; H. antiqua on, in Canada, 89, 156; new mite on, in Egypt, 411; H. antiqua on, in Holland, 270, 283; pests of, in U.S.A., 12, 83, 102, 126, 234, 388;

unfavourable food for Limonius californicus, 91; effects of insecticide treatments on, 90, 102, 156, 234, 270, 283; selenium absorbed from soil by, 290.

Onion Fly (see Hylemyia antiqua). Onion Thrips (see *Thrips tabaci*),

Onobrychis sativa, Bruchus unicolor on, in U.S.S.R., 242.

onobrychis, Macrosiphum (Acyrthosiphon). Ontario, pests of forest and shade trees in, 33, 34, 35, 78, 104, 240, 334; Lachnosterna spp. in, 163, 164; orchard pests in, 83, 167; Pyrausta nubilalis in, 77, 163; pests of vegetables in, 164, 165, 166, 167; beneficial insects and biological control in, 77, 78, 167, 240. ontario, Apechthis (Ephialtes).

Onychiurus ambulans, benzene hexachloride against, on mushrooms in

New South Wales, 365.

Onychiurus fimetarius, insecticides against, in New South Wales, 305.

oophagus, Tetrastichus.

Ootheca spp., on lima and tepary beans in Uganda, 367. opaca, Eleodes.

Opatrum sabulosum, on beet in Hungary,

operculella, Gnorimoschema.

Operophtera brumata (on fruit trees), in Belgium, 277; in Britain, 359; in Holland, 276; in Switzerland, 273; bionomics of, 277; tests of sprays against, 276.

Ophiola vaccinii, dusts against, on cran-

berry in Massachusetts, 100.

Opuntia, utilisation and value of insects against, in S. Africa, 171, 172, 389; control of Dactylopius opuntiae on spineless species of, 389.

Opuntia monterey, 389. opuntiae, Dactylopius.

Orange, pests of, in Australia, 74, 181; insects causing fruit-fall of, in Eritrea, 108; pests of, in Ivory Coast, 67, 68; Coccids on, in Persia, 387; Lepidosaphes beckii on, in Spain, 197; Coccids on, in U.S.A., 79, 83, 89, 145; persistence of DDT deposits on, 89; (fruit), new mite infesting, in Italy, 114; effect of methyl bromide fumigation on, 79, 80.

Orange, Mandarin, insects causing fruitfall of, in Eritrea, 108; Biprorulus bibax

on, in Queensland, 248. orbicularis, Otiorrhynchus.

orchidarum, Tenuipalpus. orchidii, Anaphothrips.

Orchids, Lepidosaphes machili on, California, 187.

Oregma lanigera, on Miscanthus japonicus in Fiji, 413.

Oregon, Cydia molesta in, 185; lily weevil in, 142; Scolytids in elm logs in, 15; Diabrotica duodecimpunctata intercepted in Bermuda from, 49.

Orgyia antiqua, identification of excreta of, on conifers in Germany, 76.

Orgyia mixta, on Aleurites in Nyasaland, 44.

orichalcea, Plusia (Phytometra).

Oriental Fruit Moth (see Cydia molesta). orientalis, Anychus.

Orius minutus, predacious on Tetranychus carpini in Switzerland, 282.

ornatipennis, Melcha. ornatus, Reuteroscopus.

ornithogalli, Prodenia.

Orosius argentatus (in Australia), foodplants of, 177, 178; measures against, transmitting diseases of tobacco and tomato, 219, 344.

Orthezia, on Parthenium argentatum in Mexico, 97.

Orthezia praelonga, on Citrus in Trinidad, 319.

Orthodichlorbenzene, in mixtures for treating logs and timber against pests, 103, 414; in DDT spray formula, 337.

Orthohydroxy - α -(1-piperidyl) - α -tolunitrile, toxicity of, to *Cydia pomonella*, **93.**

Orthol-D, 256.

Ortho - para - dichlorbenzophenone, 421.

oryzae, Calandra (Sitophilus); Latheticus; Lissorhoptrus.

Oryzaephilus surinamensis, in Australia, 182, 309, 310; in Switzerland, 39; in U.S.A., 25, 85; in stored cereals, 20, 39, 85, 182, 310; in other products, 25, 310; tests of DDT and benzene hexachloride on, 19, 39, 235, 236; fumigants against, 20, 85, 182; type of bag affecting infestation by, 25.

Oryzaephilus surinamensis var. mercator, infesting stored groundnuts in Senegal, 44.

Oscinella frit, DDT ineffective against, on oats in Belgium, 286.

Otiorrhynchus, on vines in Spain, 198.

Otiorrhynchus ligustici, on beet and lucerne in Hungary, 63.

Otiorrhynchus meridionalis, on Leptospermum in California, 187.

Otiorrhynchus orbicularis, on beet and lucerne in Hungary, 63.

Otiorrhynchus singularis, on raspberry in Britain, 358.

Otiorrhynchus sulcatus, on strawberry and hops in Britain, 358; test of D-D mixture against, in California, 187; not affected by DDT dust, 358.

oudemansi, Brevipalpus (Tenuipalpus).

ovata, Brachymeria. ovulorum, Entedon.

Oxalic Acid, survival of Tyroglyphus nadinus in, 114.

Oxides, survey of types of, as dust diluents, 329.

Oxypleurites mangiferae, sp. n., on mango in Hawaii, 80.

Oyster-shell Scale (see Lepidosaphes ulmi).

P

P-40, 289.

Pachymerus cassiae, parasite and control of, in stored groundnuts in Senegal, 44.

Pachyneuron siphonophorae, hyperparasite of Toxoptera graminum in Uruguay, 50.

Pachyrhina (see Pales).

Pachyrhizus, relation of constituents to insecticidal activity of seeds of, 351; colour test for evaluating, 351.

Pachyrhizus ahipa, toxicity of ground seed of, to Epilachna varivestis, 351.

Pachyrhizus erosus, 300; tests of preparations of, as insecticide, 51, 52, 103, 299, 351.

Pachyrhizus palmatilobus, toxicity of ground seed of, to insects, **51**, **52**.

Pachyrhizus strigosus, toxicity of ground seed of, to Epilachna varivestis, 351.

Pachyrhizus tuberosus, tests of preparations of seeds of, as insecticides, 300, 351.

Pachyzancla (see Psara).

pacificus, Tetranychus.

packardiana, Tortrix.

padellus, Hyponomeuta.

padi, auct., Rhopalosiphum (see R. prunifoliae).

Palaearctic Region, keys to bark-beetles of, 31, 32, 348.

Paleacrita vernata, disease of, on maple in Quebec, 319.

Pales, in tea nursery in Ceylon, 368; experiments with benzene hexachloride against, 369.

Palestine, pests of *Citrus* in, 409, 410, 412; *Empoasca lybica* in, 175, 176; natural enemies and biological control of mealybugs in, 410; Coccids introduced into Persia from, 387.

palliatus, Hylastes (Hylurgops); Tany-

mecus.

pallida, Gilpinia (Lophyrus)

pallidefulva, Formica.

pallidus, Coranus; Dysdercus; Encoptolophus.

palmerae, Periphyllus.

palmerstoni, Aulacophora (Rhaphidopalpa). Palms, book on insects associated with, 177. (See Coconut, Metroxylon and Washingtonia.)

palus, Lygus.

Panama, Anastrepha serpentina on Citrus in, 221.

pandanae, Diathetes.

Pandanus spp., Coleoptera on, in Fiji, 266, 390.

pandurus, Spilostethus.

Panicum, pests of, in Queensland, 48. 307.

Panolis flammea, identification of excreta of, on pine in Germany, 75, 76.

Panscopus sulcirostris (see Agasphaerops nigra).

Pantomorus (Graphognathus), in U.S.A., 13, 14, 184; legislation against, 184;

insecticides against, 13, 14. Pantomorus godmani (fulleri), on ornamental shrubs in Egypt, 411; bionomics and control of, on Citrus, etc., in

New South Wales, 344, 345. leucoloma, precautions **Pantomorus** against importation of, into California

from Georgia, 398.

Papaver rhoeas, food-plant of Calocoris norvegicus in Italy, 383.

Papaya, Trypetid on, in S. Africa, 176.

Paper Tubes, to protect bananas from thrips. 306.

Papilio dardanus f. hippocoon, bionomics of, on orange in Ivory Coast, 67.

Papilio demodocus (on Citrus), in Ivory Coast, 67; parasite of, in Mauritius, 210.

Papuana huebneri, food-plants of, in Pacific Is., 390.

Para Grass (see Brachiaria mutica).

Para-aminoazobenzene, tests of dust of, against *Protoparce* spp., 87.

Para-aminoazobenzene Hydrochloride, tests of dust of, against Protoparce spp., 87.

bis(Parabromomethylphenyl)trichlorethane, 421.

bis(Parabromophenyl)trichlorethane (see Bromo-DDT).

Parachloracetophenone, 421.

Parachlor- α -dichloracetophenone, 421. bis(Parachlorphenyl) Acetic Acid, excretion of, by mammals fed on DDT, 162. bis(Parachlorphenyl)dichlorethane (see DDD).

bis(Parachlorphenyl)dichlorethylene, degradation product of DDT, 140.

bis(Parachlorphenyl)sulphone, 421.

bis(Parachlorphenyl)trichlorethane (see DDT).

Parachlor-a-trichloracetophenone, 421. Paracomacris, on Eleusine in Uganda,

Paradichlorbenzene. against Aegeria exitiosa, 321, 322; against Onychiurus fimetarius, 305; in kerosene, treatment of timber with, against Coleoptera, 248, 414, 415, 416; test of, as a contact insecticide, 421.

bis(Parafluorophenyl)trichlorethane (see Fluoro-DDT).

Paraguay, Dysdercus in, 142.

parallela, Tiphia.

parallelocollis, Crypturgus pusillus.

bis(Paramethoxyphenyl)trichlorethane (see Methoxy-DDT).

bis (Paramethylphenyl) trichlorethane.

paranensis, Schistocerca.

Para-para-dichlorbenzophenone, 421.

Parascalecide, against Aegeria exitiosa, 321, 322.

Parasites, of insect pests, catalogue of, 348.

Paratetranychus citri, tests of organic compounds against, on Citrus in U.S.A., 13, 155.

Paratetranychus indicus, on sorghum in India, 427; on apple in New Zealand, 427: Coccinellids predacious on, 427.

Paratetranychus pilosus (on fruit trees), in Britain, 131, 287, 288, 359; Czechoslovakia, 288; in Holland, 129, 130, 276, 285; in Nova Scotia, 271, 392, 393; in Switzerland, 296; Tasmania, 246; in U S.A., 13, 57, 59, 101, 127, 257, 335, 374; bionomics and rearing of, 129, 374; natural enemies of, 288, 359; measures and experiments against, 13, 59, 101, 129, 130, 131, 287, 288, 335, 393; effect of DDT on populations of, 57, 59, 101, 127, 132, 246, 257, 276, 296; other sprays increasing infestation by, 271, 393.

Paratetranychus terminalis, sp. n., foodplants of, in Egypt, 411.

Paratetranychus ununguis, measures against, on Juniperus bermudiana in Bermuda, 49.

Paratheresia claripalpis (diatraeae) (see Theresia).

Paratoluenesulphonic Acid, low toxicity of esters of, to Cydia pomonella, 93.

Paratolylglycinonitrile, toxicity of, to Cydia pomonella and plants, 93.

Paratrechina longicornis, tests of Mammea americana against, 52.

Paratrioza cockerelli, bionomics of, in North Platte Valley, 144; in Colorado, 230, 231; net necrosis of potato not a symptom of yellows caused by, 230, 231; insecticides against, 230, 231.

Paris Green, in baits, 53, 96, 109, 333, 369; in dust, 184; in sprays, 41, 354, 404; DDT less toxic to man than, 162.

parkeri, Parthenoleskia; Porizon.

Parlatoria oleae, on various plants in California, 16, 185, 342, 398; measures and experiments against, 16, 17, 342, 398; possible control of, by killing males, 342, 343.

Parlatoria pergandei (on Citrus), measures to prevent introduction of, into California from Mexico, 79; in

Persia, 387.

Parlatoria pittospori, on Leptospermum in California, 187.

III Camoina, 167.

Parlatoria zizyphus, on Citrus in Persia, 387.

Parthenium argentatum, Lygus hesperus affecting seed production of, in California, 94; pests of, in Mexico, 96, 97.

Parthenoleskia parkeri, introduction of, against Diatraea saccharalis in Porto

Rico, 54.

parumstriata, Capnodis.

parva, Lissonota.

parvulus, Longitarsus; Microcerotermes (Microtermes).

Paspalum, Diatraea spp. on, in Br. Guiana, 304; Lepidoptera damaging, in Queensland, 307, 308.

Passiflora edulis, Aphids and mosaic of, in New South Wales, 365, 366.

Passiflora quadrangularis, Trypetid on, in Tanganyika, 176.

Passion Fruit (see *Passiflora edulis*).

Pastor roseus, destroying locusts in Sind, 312.

pastranai, Nemeritis (Idechthis). patruelis, Dieuches.

pauperculus, Stethorus. pauxillus, Rhynchites.

Pea, pests of, in Australia, 73, 74; Sitona lineatus on, in Belgium, 291; pests of, in Britain, 173, 261-264; pests of, in Hungary, 61, 63; Halotydeus destructor on, in New Zealand, 309; Cydia nigricana on, in Sweden, 409; Bruchus pisorum on, in U.S.S.R. 242, 243; pests of, in U.S.A., 14, 91, 153, 160, 229, 325, 326-328; variations and time of sowing of, affecting infestation by C. nigricana, 263; not attacked by Hypera brunneipennis, 55; risks and effects of insecticide treatments on, 81, 153, 154, 234, 421;

(harvested), fumigation of, against Bruchus pisorum, 62; effect of dried, on development of Ephestia elutella, 265; DDT mixed with seed of, against insects, 19.

Pea Aphis (see Macrosiphum onobrychis).

Pea Beetle (see Bruchus pisorum).
Pea Moth (see Cydia nigricana).
Pea Weevil (see Sitona lineatus).

Peach, Ceresa bubalus on, in Albania, 37; pests of, in Australia, 108, 178, 247; Myzus persicae on, in Bulgaria, 5; pests of, in Britain, 288, 358; Cydia molesta on, in Canada, 167; molesta on, in France, 238; Vasates fockeui on, in greenhouses in Holland, 239; pests of, in Italy, 64, 191, 192, Anastrepha serpentina on, in Mexico, 221; Spilostethus pandurus on, in Punjab, 123; pests of, in U.S.A., 23, 56, 79, 83, 92, 94, 127, 152, 153, 184, 189, 225, 235, 321, 334, 340, 341, 370, 373; Myzus persicae transmitting mosaic of, 79, 93; insects causing deformation of fruits of, 235, 334, 373; insecticides and injury to, 93, 224, 253.

Peach (Fruit), new mite feeding on, 114;

DDT residues on, 135.

Peach Aphis, Black (see Anuraphis persicae-niger).

Peach Moth, Oriental (see Cydia molesta).

Peanuts (see Groundnuts).

Pear, pests of, in Australia, 179, 245, 246, 247; pests of, in Britain, 358; new mite on, in Egypt, 412; pests of, in France, 238, 277; pests of, in Holland, 285, 286; pests of, in Italy, 64, 194, 253; Pseudococcus comstocki on, in Japan, 410; Melanaspis inopinata on, in Persia, 386; pests of, in U.S.A., 80, 334, 399; not attacked by Leucoptera scitella, 269; thiocyanoacetates causing scorching of, 253.

Pear (Fruit), new mite feeding on, 114; tolerance for DDT residues on, in

U.S.A., 162, 209.

Pear Bud Mite, a form of *Eriophyes* pyri, 82.

Pear Leaf Blister Mite (see Eriophyes

Pear Psylla (see Psylla pyricola).

Pear Rust Mite (see Epitrimerus pyri). Pecan, pests of, in U.S.A., 14, 151, 152, 399.

Pecan Weevil (see Curculio caryae). pectinicornis, Cladius.

pectoralis, Wlk., Dacus (African Species) (see D. bivittatus cucumarius).

pectoralis, Wlk., Dacus (Oriental Species).

Pediculoides graminum, associated with Nigrospora oryzae on wheat in Spain, 202, 203.

Pediculoides ventricosus, attacking Bruchus pisorum in Hungary, 62.

Pediculopsis (see Pediculoides).

Pediculus ventricosus (see Pediculoides). pellionella, Tinea.

Pelopidas borbonica, on rice in Mauritius, 210.

penitalis, Pyrausta.

Pennisetia marginata, on raspberry, boysenberry and blackberry in California, 399.

Pennisetum typhoides, Schistocerca gregaria damaging, in Sind, 310, 311.

Pennsylvania, miscellaneous pests in, 18, 301, 325, 336.

Pentabromophenol, repellency of, to

Kalotermes brevis, 233.

Pentachlorphenol, 275; treatments of timber with solutions containing, against insects, 1, 107, 169, 245, 414, 415; and derivatives, tests of toxicity of, to Arthropods and plants, 223, 224; question of toxicity of, to man, 107, 108.

Pentalonia nigronervosa, transmitting bunchy-top virus of banana in Australia and Pacific Is., 46, 267, 413; campaign against, in New South Wales, 46.

Pentatrichopus (see Capitophorus).

Penthaleus major, in New Zealand, 309; Halotydeus destructor compared with, 309.

Pepper, Black, use of ground, against Bruchus obtectus in stored beans, 21.

Peppers (see Capsicum).

Peregrinus maidis, dusts against, on maize in Virginia, 332.

Perforadix sacchari, measures against, on sugar-cane in Porto Rico, 53.

pergandei, Corythucha; Parlatoria.

Pericyma glaucinans, on Poinciana regia in Mauritius, 210.

Peridroma saucia (margaritosa), toxicity of benzene hexachloride and DDT to, 22.

Perilampus, parasite of Apanteles glomeratus in Germany, 407.

Perilampus italicus, parasite of Athalia rosae in Italy, 193.

Perilampus similis, hosts of, in U.S.A., 29, 84.

Periphyllus palmerae, sp. n., on Acer glabrum in Utah, 388.

Periplaneta americana, use of, for insecticide tests, 52, 175, 378.

periusalis, Psara.

perniciosus, Mesochorus; Quadraspidiotus (Aonidiella, Aspidiotus). pernyi, Antheraea.

peroneae, Macrocentrus.

Persectania composita (ewingi), damaging pastures in New Zealand, 244.

Persia, digest of plant pest legislation in, 209; miscellaneous pests in, 385-387; natural enemies and biological control of Coccids in, 385, 386, 387.

persicae, Myzus.

persicae-niger, Anuraphis.

Persimmon, Coccids on, in California, 189, 399.

persuasoria, Rhyssa.

Peru, miscellaneous pests in, 133, 134; new Coccid on avocado in, 401; beneficial insects in, 133, 134; digest of plant-quarantine restrictions in, 105. peruella, Mescinia.

Petrobia cepae, sp. n., on onion in Egypt,

411.

Petrol, experiments with types of, as grain fumigants, 1-3; treatment of timber with, against *Lyctus*, 1.

Petroleum Ether, as a solvent, 51.

Phaeogenes hariolus, parasite of Harmologa fumiferana in Canada, 207.

phaeorrhoea, Nygmia (see Euproctis chrysorrhoea).

Phalaena chrysorrhoea (see Euproctis). Phalaena similis (see Euproctis).

Phalaenoides glycinae, Pentatomids predacious on, on vines in New South Wales, 108.

phalerata, Campsomeris.

phaseoli, Agromyza (Melanagromyza).

Phaseolus aconitifolius, 310.

Phaseolus acutifolius (Tepary Bean), Coleoptera on, in Uganda, 367.

Phaseolus lunatus (see Lima Beans).

Phaseolus vulgaris (see Beans).

Pheidole punctulata, measures against, associated with Coccids in Kenya, 404.
Phenacoccus, new parasite of, on Andropogon in Maryland, 315.

Phenacoccus gossypii, laboratory host of

Pseudaphycus mundus, 316.

Phenobremia aphidimyza, bionomics of, predacious on Aphis gossypii in Italy, 193.

Phenol, in DDT spray formula, 28; production of insecticides by chlorination of, 275; thiocyanoacetic ester of, as an insecticide, 253, 254; used in preparation of Coccids, 104.

Phenothiazine, ineffective against grass-hoppers, 341, 342; against orchard pests, 58, 59, 278, 335; tests of, on Leptinotarsa decemlineata, 296; low toxicity of, to bees, 222; not preventing onion-bulb decay, 283; in oil-impregnated dusts, 58, 59; preventing

scorching by lead arsenate in mixed dust, 59; spraying with, 278, 335, 341, 342.

Phenylmercuric Chloride, repellent to Kalotermes brevis, 233.

1-Phenyl-1-p-chlorphenyl-2, 2, chlorethane (4-Chlordiphenyltrichlorethane), 275, 421.

phidilealis, Hellula.

Philaenus leucophthalmus (in U.S.A.), on cereals, 136; damage to other crops by, 160.

Phleum pratense, 374; Lachnosterna damaging, in Ontario, 164.

Phlox, Tetranychus bimaculatus on, in U.S.A., 335.

Phlyctaenia rubigalis, tests of benzene hexachloride and DDT against, U.S.A., 22, 234.

Phlyctaenia tertialis, parasite

U.S.A., 29.

Phlyctaenodes sticticalis (see Loxostege). Phoracantha senio, intercepted in Fiji in logs from Australia, 267.

Phorocera incrassata (parasite of Harmologa fumiferana), in British Columbia, 78; introduction of, into eastern Canada, 78.

Phosphine, unsatisfactory for fumigation of nursery stock against Quadraspidiotus

perniciosus, 292.

Phosphorus Acids, toxicity of esters of, to Aphids, 228. (See Hexaethyl Tetraphosphate.)

Phragmites communis, Hyalopterus arundinis on, in Italy, 191, 192.
Phthalonitrile, tests of dusts of, against

Protoparce spp., 87.

Phthorimaea (see Gnorimoschema).

phthorimaeae, Chelonus; Omorgus.

phycidis, Habrocytus.

Phyllobius, on fruit trees in Britain, 358. Phyllobius oblongus, on strawberry in Belgium, 292, 354; spray against, 292. Phyllocoptes fockeui (see Vasates).

Phyllocoptes lycopersici, dusts and sprays against, on tomato in Queensland, 362, 363.

Phyllocoptes schlechtendali (see Vasates).

Phyllocoptruta oleivorus (on Citrus), in Australia, 181, 249; in U.S.A., 13, 83, 187; not affected by high temperature, 181; acaricides against, 13, 249; DDT increasing infestation by,

Phyllophaga (see Lachnosterna).

Phyllotreta, measures against, on crucifers in Spain, 204.

Phyllotreta aerea, on crucifers in Ontario, 167.

Phyllotreta atra (on crucifers), in England, 132; bionomics of, in Spain, 204.

Phyllotreta cruciferae (on crucifers), in Britain, 132, 377; in Spain, 204; in United Provinces, 122; transmitting new virus disease of turnip, 377; bionomics and control of, 122, 204.

Phyllotreta diademata, on crucifers in England, 132.

Phyllotreta nemorum (on crucifers), England, 132; bionomics of, Spain, 204.

Phyllotreta nigripes, bionomics of, crucifers in Spain, 204.

Phyllotreta undulata (on crucifers), 132; bionomics of, England, Spain, 204.

Phyllotreta variipennis, bionomics of, on crucifers in Spain, 204.

Phyllotreta vittata, dusts against, on crucifers in U.S.A., 234, 421.

Phyllotreta vittula, transmitting new virus disease of turnip in Britain, 377; bionomics of, in Spain, 204.

Phylloxera vitifoliae (on vines), outbreak of leaf form of, in California, 398; in Switzerland, 296; measures against, 296, 398.

Physalis, Paratrioza cockerelli on, in U.S.A., 144.

Physonota alutacea, proposed introduc-tion of, into Mauritius from Trinidad against Cordia macrostachya, 424.

Phytodietus annulatus, parasite of Tortrix

packardiana in Quebec, 321.

Phytodietus fumiferanae (parasite of Harmologa fumiferanae), introduction of, into eastern Canada from British Columbia, 78; bionomics and rearing

Phytolyma lata, bionomics of, on Chlorophora excelsa in Ivory Coast, 68.

Phytometra (see Plusia).

Phytomyza lateralis, benzene hexachloride against, in carrot in Switzerland, 281.

Phytomyza rufipes, benzene hexachloride against, in cabbage in Switzerland, 281. Phytonomus (see Hypera).

Phytophthora infestans, sprays affecting infestation of potato by, in Ontario.

Phytoptus avellanae, on Corylus in N. America, 83.

Picea (see Spruce).

Picea glauca, Harmologa fumiferana on, in Ontario, 33, 240.

Picea mariana, pests of, in Canada, 33, 320.

piceae, Chermes (Adelges). piceus, Attagenus.

picipes, Camponotus (Tanaemyrmex); Rhabdopterus; Stethorus.

picta, Bagrada (see B. cruciferarum).

Pieris brassicae, parasite and hyperparasites of, in Europe, 406, 407; tests of DDT dust against, in India, 123.

Pieris monuste (see Ascia).

Pieris rapae (on crucifers), in Australia, 70, 74, 215, 216, 218; in China, 299; in New Zealand, 244; in U.S.A., 159, 234, 333, 421; parasites and biological control of, 70, 215, 216, 244; insecticides against, 22, 23, 74, 218, 234, 299, 333, 421.

Piesma quadratum, transmitting beet crinkle in Czechoslovakia, 355.

Pigmeophorus americanus, 365.

Pigs, effects of feeding, on apples sprayed with lead arsenate, 71, 72.

pilosella, Campsomeris.

Pimento (see Capsicum).

Pimpla inflata, parasite of Camptylochila aemula in Quebec, 321.

Pimpla pomorum (see Ephialtes).

pinastri, Sphinx.

Pine, Lymantria monacha on, in Austria, 128; pests of, in Britain, 215, 315; pests of, in Canada, 320, 334; pests of, in Germany, 75, 76; Scolytids in, in Italy, 195; Sirex noctilio on, in New Zealand, 244; pests of, in U.S.A., 143, 144, 158, 220, 301, 333, 334; timber of, used in tests of preservatives against insects, 107, 249, 414.

Pine, Jack (see *Pinus banksiana*). Pine, Sugar (see *Pinus lambertiana*). Pine, White (see *Pinus strobus*).

Pine Oil, in mixtures for treating timber against pests, 414; in formula for DDT dip, 146; Wetsit (q.v.) containing, 69.

Pine Tip Moth (see *Rhyacionia frustrana*). Pineapple, *Anomocaulus fulvovestitus* on, in Fiji, **266**; mealybugs inter-

cepted on, in U.S.A., 315. Pineus (see Chermes).

pini, Dendrolimus; Diprion (Lophyrus); Lygaeonematus (see Pristiphora abietina).

piniarius, Bupalus (Geometra). pinifoliae, Chermes (Pineus).

piniperda, Myelophilus (Blastophagus).

pinivora, Acantholyda. Pinus (see Pine).

Pinus banksiana, Monochamus scutellatus in fire-damaged, in Quebec, 320; pests of, in U.S.A., 220.

Pinus echinata, Rhyacionia frustrana on, in Oklahoma, 158.

Pinus lambertiana, twig injury to, by Conophthorus lambertianae in California, 301.

Pinus pinea, Scolytids on, in Italy, 195. Pinus radiata, used in tests of timber preservatives against insects, 414.

Pinus resinosa, pests of, in U.S.A., 220. Pinus strobus, Chermes causing twig injury to, in N. Carolina and Canada, 333, 334.

Piper nigrum (see Pepper, Black).

Piperonyl Cyclohexenone or Cyclohexanone, as synergist with pyrethrum, 222, 230, 331, 335, 395; with other insecticides, 99, 394; toxicity of, to Halticus bracteatus, 394.

Pirenisca, n.n. (Pirene, Hal.), 316.

piri, Anthonomus.

pirifoliae, Epitrimerus (see E. pyri).

Piripiri (see Acaena).

piriventris, Atta (Neoatta) sexdens. pisi, Macrosiphum (see M. onobrychis).

pisorum, Bruchus.

Pissodes strobi, spray against, on pine in U.S.A., 144.

Pistacia spp. (Pistachio), pests of, in Persia, 385, 386.

pistaciae, Lepidosaphes; Leucaspis; Megastigmus; Pulvinaria.

Pithecolobium dulce, Celosterna scabrator on, in Réunion, 317.

pittospori, Parlatoria.

Pittosporum tobira, Asterolecanium arabidis on, in California, 187.

Pityophthorus, species of, on Parthenium argentatum in Mexico, 97.

Plaesius javanus, failure to establish, against Cosmopolites sordidus in Uganda, 367.

Plagiodera versicolora, on willow in Canada, 104.

Plagionotus floralis, bionomics of, on lucerne in Hungary, 63.

Plant Pest Legislation, in Nyasaland, 32, 105; in U.S.A. and territories, 71, 105, 184, 209; summaries of, in other countries, 105, 184, 209.

Plant Protection, bibliography of literature of, in 1937–39, 72.

Plants, study of, as sources of insecticides, 9, 51, 52, 298, 299; review of literature on insecticides obtained from, 308.

platanoides, Drepanosiphum.

Platanus occidentalis, Lachnosterna submucida on, in Texas, 14.

platensis, Aphidius.

Platyedra gossypiella, factors favouring increase of, in Porto Rico, 300; parasites of, in Punjab, 120, 121; long-cycle larvae of, not occurring in Somalia, 67; in Trinidad, 126; in

Uganda, 366, 367; domestic quarantine against, in U.S.A., 105; not found in California, 398; treatments of cottonseed against, 66, 105.

Platylactista azteca, on Parthenium argen-

tatum in Mexico, 97.

Platyparea poeciloptera (Asparagus Fly), in Sweden, 408.

Platypodids, bibliography of, 72.

Platypus, susceptibility of Inga spp. to, in Porto Rico, 300.

Plectops currei, parasite of Dioryctria reniculella in Quebec, 321.

Plesiocoris rugicollis, sprays against, on apple in Britain, 284.

plesiorapae, Diaeretus.

pleurostigma, Ceuthorrhynchus.

Pleurotropis, parasite of Leucoptera scitella in Italy, 269.

Pleurotropis nigripes, hyperparasite of Laphygma exempta in Nyasaland, 45. Plodia interpunctella, in Trinidad, 319;

dusts against, in stored seeds, 19, 319. Plum, pests of, in Britain, 358, 359; Eulecanium corni on, in Bulgaria, 354; mites on, in Canada, 83; pests of, in Czechoslovakia, 65, 289; new mite on, in Egypt, 412; pests of, in Holland, 239, 285, 286; pests of, in Italy, 64, 192; Anuraphis persicae-niger on, in New South Wales, 247; pests of, in Poland, 254, 255, 354; Aphids on, in Spain, 199; Hoplocampa spp. on, in Switzerland, 296; pests of, in U.S.A., 14, 83, 92, 399, 400; not injured by azobenzene, 284; methyl-bromide fumigation of fruits of, 186.

Plum, Beach (see Prunus maritima).

Plum Curculio (see Conotrachelus nenuphar).

Plusia argentifera, insecticides against, on vegetables in Australia, 74, 363.

Plusia brassicae (see Trichoplusia ni).

Plusia chalcites, on tobacco in Queensland, 69.

Plusia gamma (see Autographa).

Plusia orichalcea, on tobacco in Uganda, 367.

Plutella maculipennis (on crucifers), in Australia, 74, 177, 178, 218; in Belgium, 278, 292; in Holland, 286; in Louisiana, 159; introduced parasites of, in New Zealand, 244; in Porto Rico, 52; in Trinidad, 319; insecticides against, 51, 52, 74, 178, 218, 244, 278, 286, 292. pluto, Xylophanes.

Podagrica fuscicornis, on malvaceous

plants in Rumania, 41.

Podagrica malvae, bionomics of, on malvaceous plants in Rumania, 41.

podana, Tortrix (Cacoecia).

Podocarpus spp., used in tests of timber preservatives against insects, 414, 415.

poeciloptera, Platyparea. poecilopus, Habrocytus.

Poeciloscytus basalis (see Polymerus).

Poëphilina (Poëphila) flaveola, bionomics of, on Oncoba echinata in Ivory Coast, 67, 68.

Pogonomyrmex, killed by benzene hexachloride dust in Oklahoma, 100.

Pogonomyrmex barbatus var. molefaciens, measures against, in fields and gardens in Oklahoma, 94.

Pogonomyrmex occidentalis, lizards and

birds destroying, in Utah, 24.

Poinciana regia, Pericyma glaucinans on, in Mauritius, 210; use of wood of, in tests of repellents against termites, 233. Poinsettia (see Euphorbia pulcherrima).

Poland, introduced Bruchids in, 385; pests of juniper in, 267, 268; orchard pests in, 254, 255, 354; spread of Leptinotarsa decemlineata into, 254, 384; susceptibility to L. decemlineata of varieties of potato grown in, 42, 43; natural enemies of pests in, 255, 268; book on plant protection in, 117.

poligraphus, Polygraphus.

Polistes canadensis, predacious on Schematiza cordiae in Trinidad, 425. Polistes gallicus, predacious on Leptinotarsa decemlineata in Spain, 196.

polita, Homala.

politiventris, Bracon (Microbracon).

Pollination, of lucerne, bees causing, 10. Pollinia pollini, on olive in California, 187, 399.

Polychlorcyclane Sulphide, in sprays against *Anthonomus* spp., 278, 293; toxicity of, to bees, 237.

Polychrosis botrana (on vines), in Switzerland, 296; bionomics of, in Turkmenistan, 3, 4; insecticides against, 4, 296.

Polychrosis viteana, sprays against, on

vines in Illinois, 127.

Polydrusus cervinus, on fruit-trees, etc., in Britain, 358.

Polydrusus sericeus, on cherry in Britain, 358.

Polygraphus poligraphus, on spruce in Switzerland, 407.

Polyhedral Disease, of Lymantria monacha, 129.

Polymerus (Poeciloscytus) basalis, on Parthenium argentatum in Mexico, 97.

Polynema fulmeki, sp. n., parasite of Quadraspidiotus perniciosus in Austria, 64.

Polynema laeta, redescription of, 64. polytoma, Gilpinia (Lophyrus).

Pomegranate, new mite on, in Egypt, 412. pometaria, Alsophila,

pomi, Aphis (Doralis); Seius.

pomonella, Cydia (Carpocapsa, Laspeyresia); Rhagoletis.

pomorum, Anthonomus; **Ephialtes**

(Pimpla).

Pompostola hypparchus, bionomics of, on Ceiba pentandra in Ivory Coast. 67.

Popillia japonica (in U.S.A.), quarantine regulations against, 209; prevention of spread of, by land transport and aircraft, 209, 336; tests and uses of insecticides against adults of, 36, 101, 209, 234, 333, 335, 336, 337, 339; on shade and orchard trees, 36, 335, 336, 339; defoliating soya beans, 18; not found in California, 398.

Poplar, Cosmia trapezina on, in Belgium, 277; Hyphantria cunea on, in U.S.A.,

254.

Poppy (see Papaver rhoeas).

Popular Names, of insects, lists of, 32, 72. Population Studies, of insects, 5, 210, 352. populi, Byctiscus.

Porizon spp., attempted introduction of, into California from S. America against Listroderes obliquus, 97.

porosus, Myzus.

Porthesia (see Euproctis).

Porthetria (see Lymantria).

Porto Rican Mole Cricket (see Scapteriscus vicinus).

Porto Rico, Kalotermes brevis in, 52, 169, 232, 300; miscellaneous pests in, 51-55, 300; beneficial insects introduced into, 53, 54, 316; tests of plants as sources of insecticides in, 51, 52; plant quarantines in, 71, 105, 209; pests intercepted in quarantine in, 240.

Portugal, Lymantria dispar on cork oak in, 285; migration of Schistocerca gregaria into, 205.

postfasciatus, Euscepes.

Potamogeton natans, insects infesting, in Italy, 347.

Potassium Arsenate, tests of toxicity of, to Agriotes, 287.

Potassium Dichromate, Celcure (q.v.)containing, 414.

Potassium Hydroxide, in DDT spray formula, 28; survival of Tyroglyphus nadinus in solutions of, 114.

Potato, pests of, in Australia, 46, 74, 178, 216, 217, 249; Leptinotarsa decemlineata on, in Belgium, 291; Gnorimoschema operculella on, Bermuda, 49; Agallia albidula on, in Brazil, 77; pests of, in Canada, 164, 165, 167; grasshoppers damaging, in Chile, 109; Aphids on, in England, 195, 362; L. decemlineata on, in France, 349; Calocoris norvegicus on, in Italy, 383; L. decemlineata on, in Jersey, 259; Halotydeus destructor on. in New Zealand, 309; Empoasca lybica on, in Palestine, 176; L. decemlineata on, in Poland, 254, 384; L. decemlineata on, in Spain, 196: L. decemlineata on, in Switzerland, 297; pests of, in U.S.A., 56, 102, 144, 209, 225, 228, 230, 231, 234, 328, 332, 421; insects and virus diseases of, 5, 7, 8, 230, 231; resistance of varieties of, to Aphids and leaf-roll, 209; experiments with colour forms of Macrosiphum solanifolii and, 362; experiments with L. decemlineata and Polish varieties of, 42, 43; effects of insecticides on, 224, 225, 234, 236, 280; (tubers), Gnorimoschema operculella in, 46, 49, 56, 217, 249; factors affecting rearing Macrocentrus ancylivorus on G. operculella in, 18; quarantine treatment of, with DDT against *Popillia japonica* in U.S.A., 209; pieces of, as traps for wireworms, 183.

Potato Aphis (see Macrosiphum solani-

folii).

Potato Beetle, Colorado (see Leptinotarsa decemlineata).

Potato Beetle, Three-lined (see Lema trilineata),

Potato Flea-beetles (see Epitrix cucumeris and E. tuberis).

Potato Late Blight (see Phytophthora infestans).

Potato Leafhopper (see *Empoasca fabae*). Potato Psyllid (see Paratrioza cockerelli). Potato Tuber Moth (see Gnorimoschema operculella)

Potentilla, differences in susceptibility of species of, to colonisation by Capitophorus spp. and infection with strawberry crinkle in Britain, 313.

potentillae, Capitophorus (Pentatrichopus). Powdery Mildew (see Erysiphe polygoni).

praelonga, Orthezia. praetiosa, Bryobia. praeusta, Tetrops.

Prangos pabularia, constituents and value of, as a source of insecticides, 118.

Prays oleellus, bionomics and control of, on olive in Italy, 141, 193, 194.

Prenolepis longicornis (see Paratrechina). Prickly Lettuce (see *Lactuca scariola*).

Prickly Pear (see Opuntia).

Principe, Dacus bivittatus in, 176.

Prionomitus, parasite of Psylla in Persia, 386.

Prionus coriarius, bionomics of, in Britain, 214, 215.

Pristiphora abietina, identification excreta of, on spruce in Germany, 75, 76; effectiveness of locally grown pyrethrum against, in Holland, 270.

Pristomerus appalachianus (see Neopristo-

merus).

Pristomerus euryptychiae, parasite of Walshia amorphella in Oklahoma, 9.

Pristomerus testaceicollis, parasite of Eublemma amabilis in Bihar, 120.

Pristonychus terricola, damaging stored apples in Britain, 358.

Privet (see Ligustrum).

Probits, for expressing toxicity of insecticides, 381.

Proceras sacchariphagus, parasites of, on sugar-cane in Mauritius, 210.

processionea, Thaumetopoea.

Prodenia eridania (see Laphygma).

Prodenia litura, on tobacco in Queensland, 69.

Prodenia ornithogalli, dusts against, on tomato in Utah, 10.

promissa, Nepticula.

Prontaspis citri (see Unaspis).

Prophanurus alecto (see Telenomus). Propyl Pentachlorphenyl Ether, 224.

Propylene Dichloride, fumigation tests with, against insects in stored maize,

Propylene Laurate, and oil, dusts impregnated with, 229.

Prosayleus dispar, bionomics of, in New South Wales, 73.

Prospaltella, parasite of Quadraspidiotus perniciosus in Italy, 65.

Prospaltella fasciata, parasite of Quadraspidiotus perniciosus in Italy, 191.

proteus, Urbanus.

Protoparce quinquemaculata, tests insecticides against, on tobacco N. Carolina, 87.

Protoparce sexta, tests of insecticides against, on tobacco in U.S.A., 87, 371, 372.

Protopulvinaria (see Pulvinaria).

Protostrophus falsus, sp. n., on sweet potato in S. Africa, 388.

protuberans, Dendrosoter.

proxima, Athalia,

pruinosum, Eulecanium (Lecanium).

Prune, Eulecanium kunoense on, California, 399.

prunifoliae, Rhopalosiphum.

prunorum, Diptilomiopus (see D. gigantor-

Prunus, Hyalopterus arundinis on, in Italy, 192. (See Cherry, Peach, Plum, etc.)

Prunus emarginata, Rhagoletis spp. on, in California, 185, 186.

Prunus laurocerasus, Cydia molesta on, in France, 239.

Prunus maritima, Paratetranychus pilosus on, in Massachusetts, 101.

Psalidium maxillosum, on castor and beet in Hungary, 62, 63.

Psallus seriatus, on cotton, etc., in U.S.A., 156, 183, 184; bionomics of, 183; dusts against, 184.

Psammodes similis, soil treatments against, in tobacco fields in S. Rhodesia, 137.

Psara bipunctalis, comparison of toxicity of organic insecticides to, 13, 22, 23.

Psara periusalis, on tomato in Bermuda,

Pseudaonidia trilobitiformis, on Citrus in Nyasaland, 44.

Pseudaphycus, key to species of, 315.

Pseudaphycus abstrusus, sp. n., parasite of Pseudococcus comstocki in Virginia,

Pseudaphycus alveolatifrons, sp. n., parasite of Pseudococcus comstocki in New Jersey, 315.

Pseudaphycus angustifrons, sp. n., parasite of pineapple mealybugs intercepted in U.S.A. from Cuba, 315.

Pseudaphycus limatulus, sp. n., parasite of Phenacoccus in Maryland, 315.

Pseudaphycus malinus, sp. n. (parasite of Pseudococcus comstocki), in Japan and Korea, 315; established in U.S.A., 315.

Pseudaphycus meracus, sp. n., parasite of Ferrisia virgata in Virginia, 315.

Pseudaphycus meritorius, sp. n., parasite of Ferrisia virgata in Virginia, 315.

Pseudaphycus mundus, sp. n., hosts and distribution of, confused with Pseudococcobius terryi, 316.

pseudobrassicae, Rhopalosiphum (Aphis). Pseudocneorrhinus bifasciatus, in Japan, 301; food-plants and distribution of, in U.S.A., 301.

Pseudocneorrhinus setosus, P. bifasciatus recorded as, in U.S.A., 301.

Pseudococcobius terryi, parasite of Pseudococcus boninsis in Hawaii, 316; records of introduction of, into U.S.A. and Porto Rico due to confusion with Pseudaphycus mundus, 316.

Pseudococcus, new hyperparasite of, in Uganda, 109.

Pseudococcus adonidum, effect of DDT on populations and natural enemies of, on Citrus in California, 145; laboratory host of Pseudaphycus mundus, 316.

Pseudococcus boninsis, confusion of parasite of, on sugar-cane in Hawaii and U.S.A., 316.

Pseudococcus brevipes, on groundnuts in E. Africa, 426; probable parasite of, on pineapple in Cuba, 315.

Pseudococcus calceolariae, on sugar-cane

in Br. Guiana, 304.

Pseudococcus citri, food-plants and distribution of, in W. Africa, 110, 111; ants associated with, 110; parasite of, 111; cacao viruses transmitted by, 110.

Pseudococcus comstocki, in Japan and Korea, 316, 410; in U.S.A., 315, 316, 399, 410; on apple and pear, 399, 410; parasites and biological control of, 315, 316, 410; doubtful identity, natural enemies and biological control of mealybugs recorded as, on Citrus in Palestine, 409, 410.

Pseudococcus filamentosus, food-plants of,

in Punjab, 123.

Pseudococcus kenyae, factors affecting biological control of, on coffee in Kenya, 404.

Pseudococcus lepelleyi, new hyperparasite

of, in Java, 109.

Pseudococcus longispinus (see P. adonidum).

Pseudococcus maritimus, other mealybugs possibly confused with, on fruit trees in California, 399.

Pseudococcus njalensis, transmitting swollen-shoot of cacao in Gold Coast, 110; ants associated with, 110; natural enemies and list of food-plants of, 110.

Pseudococcus sacchari (see Trionymus). Pseudococcus virgatus (see Ferrisia). pseudomagnoliarum, Coccus.

Pseudomonas, infecting cherry in Britain,

Pseudosarcophaga affinis (see Agria).
pseudosolani, Aulacorthum (Myzus) (see
Macrosiphum solani).

Pseudotsuga taxifolia, Chermes cooleyi on, in Belgium, 353.

psidii, Pulvinaria.

Psidium, Dacus ferrugineus dorsalis in fruits of, in Hawaii, 249. (See Guava.)

Psila rosae (Carrot Fly), in Britain, 377, 417; bionomics of, 417; measures against, 377; morphology of immature stages of, 240.

Psilogaster, Blanch. (see Psilogastrellus). Psilogaster, R.L., 316.

Psilogastrellus, n.n. for Psilogaster,

Blanch., 316. *Psylla*, parasite of, on *Pistacia* in Persia, 386.

Psylla pyricola (on pear), Prangos pabularia ineffective against, in Kirghizia, 118; sprays against, in U.S.A., 36, 334, 335; not found in California, 398.

Psylla, Pear (see Psylla pyricola).

Psyllid, Potato (see Paratrioza cockerelli). Psylliodes chrysocephala (on crucifers), bionomics of, in Spain, 204; in Switzerland, 281; measures against, 204, 281.

Pteris cretica, insecticide injuring, 282. Pteromalus, parasite of Apion damryi in

Italy, 31.

Pteromalus puparum, utilisation and value of, against Pieris rapae in Australia, 70, 216.

Pteronidea (see Nematus).

pterostichus, Thisoecetrinus.
Ptinus tectus, in stored wheat in Britain,

212. Ptychodes trilineatus, 250.

Ptychodes trilineatus insularis, in fig trees in Tahiti, 250.

Ptyelus grossus, on castor in Uganda, 367. pubescens, Agonoscelis; Harpalus (see H. rufipes); Hydnocera.

puera, Hyblaea.

Pueraria yunnanensis, insecticidal properties of, 299.

pugnax, Solubea.

Pulvinaria floccifera, on Citrus in Persia, 387.

Pulvinaria pistaciae, on Pistacia in Persia, 386.

Pulvinaria psidii, on fruit trees in California, 189.

Pulvinaria pyriformis, discovery and eradication of, on Fatsia japonica in California, 400.

Pumice, as a diluent for dusts, 329.

pumila, Agathis (Bassus).

Pumpkin, not attacked by grasshoppers in Chile, 109; Aphis gossypii on, in Italy, 193; susceptibility of, to Melittia cucurbitae in Massachusetts, 101; effect of benzene hexachloride on, 23.

punctatissima, Coptosoma. punctatum, Anobium. punctiferalis, Dichocrocis. punctillum, Stethorus (Scymnus). punctiventris, Cleonus. punctulata, Cicindela; Pheidole.

Punjab, cotton insects in, 120, 312; miscellaneous pests in, 122, 123; beneficial insects in, 120, 121.

puparum, Pteromalus. purchasi, Icerya. purpurea, Clausenia. pusillus, Crypturgus. pustulans, Asterolecanium.

Puto yuccae, on Parthenium argentatum in Mexico, 96.

putrescentiae, Tyrophagus. pyralina, Cosmia (Calymnia). Pyrameis cardui (see Vanessa). pyrastri, Lasiophthicus.

Pyrausta futilalis, parasite of, in U.S.A.,

29.

Pyrausta nubilalis (on maize), in Canada, 77, 163, 167, 391; in U.S.A., 14, 22, 29, 81, 95, 102, 128, 401–403; factors affecting fecundity of, 251, 252; effect of varieties and stage of growth of maize on, 163, 402, 403; parasites and biological control of, 29, 77, 78, 391; tests and uses of insecticides against, 14, 22, 81, 95, 102, 400–403; cultural measures against, 163.

Pyrausta penitalis, parasite of, in U.S.A.,

29.

Pyrax, as carrier for dusts, 19, 229.

Pyrax ABB, 224, 233, 332.

Pyrethrins, content of, in pyrethrum dusts, sprays and aerosols, 11, 14, 102, 112, 147, 168, 272, 278, 295, 300, 331, 335, 336, 364, 394, 395; ester similar to, in mamey seed, 51.

Pyrethrum (including extracts), against Aphids, 300, 335; against Jassids, 147, 335, 398; against Mirids, 11, 102, 335, 394; against other Hemiptera, 75, 295, 335; against Coleoptera, 41, 230, 235, 278, 335, 336, 337, 345; against Lepidoptera, 126, 168, 230, 331; and oil, treatment of maize ears with, against Heliothis, 126; against Lepismids, 364, 379; against mites, 335; against Rhagoletis pomonella, against sawflies, 335; 112, 335; factors influencing effectiveness of oil films of, against warehouse insects, 168; in aerosols, 147, 331, 336, 337; dusting with, 11, 75, 102, 112, 230, 272, 295, 331, 379; carriers for, 112, 295, 379; and benzene hexachloride, 99, 235; and DDT, 336, 337; and DDT analogue, 14; and dinitro-o-cyclohexylphenol, 394; vaporised or atomised sprays of oil containing, 36, 147, 364, 398; and rotenone insecticides, 300, 335; and sulphur, 11, 102; and thiocyanate, 398; addition of other agents to, 222, 230, 331, 335, 380, 381, 395; effects of humidity, light and temperature on toxicity of, 168, 379, 380; review of properties and uses of, 297; quality of, obtained from Dutchgrown plants, 270.

Pyrethrum cinerariaefolium (see Chrysan-

themum).

Pyrgodera armata, crops damaged by, in Persia, 387, 388.

pyri, Brevipalpus; Epitrimerus; Eriophyes.

pyricola, Psylla.

Pyridine, and soap, spraying with, against Contarinia nasturtii, 286.

pyriformis, Pulvinaria (Protopulvinaria). Pyroderces rileyi, insecticides against, in stored maize in S. Carolina, 369.

Pyrophyllite, spray suspensions of DDT in, 57, 58, 98, 143, 150, 235, 246; as a carrier for dusts, 12, 25, 47, 69, 74, 87, 91, 94, 96, 126, 138, 143, 149, 158, 178, 217, 218, 224, 229, 233, 236, 237, 329, 332, 338, 371, 379, 402, 420; electrostatic charges of dusts of, 393; evaluation of, 340.

Pyrus malus var. astracanica, pests of, in U.S.A., 335.

pyste, Nemorilla.

Q

quadra, Oeonistis.

Quadraspidiotus perniciosus, in Algeria, 293; in Austria, 292; in France, 296; in Hungary, 64; in Italy, 64, 191; bionomics of, 64; natural enemies of, 64, 65, 191; measures and experiments against, on fruit trees and nursery stock, 65, 292, 293, 296.

quadratum, Piesma. quadridens, Ceuthorrhynchus. quadridentata, Ascogaster. quadrigemina, Chrysomela. quadripustulata, Brochymena. quadripustulatus, Exochomus. quadrisetus, Eriophyes.

Quarantine, pests intercepted in: in Bermuda, 49; in Fiji, 267; in Kenya, 405; in Hawaii and Porto Rico, 240; in U.S.A., 147, 233, 240, 315, 316, 400, 401.

Quartz, electrostatic charges of dusts of, 393.

Quassia Spray, against Hyalopterus arundinis, 192.

Quebec, pests of forest and shade trees in, 34, 35, 78, 104, 163, 240, 319, 320, 321, 334; Dutch elm disease in, 163, 320; miscellaneous pests in, 77, 104, 167, 168; beneficial insects and biological control in, 77, 78, 240, 320, 321.

Queensland, pests and disease of banana in, 46, 308; other fruit pests in, 180, 248, 249; Chortoicetes terminifera in, 47, 48; Lepidoptera damaging cereals and grasses in, 307, 308; pests of solanaceous plants in, 69, 216, 249, 362, 363; pests of stored products in, 68, 249, 309, 310, 363; Lyctus spp. in

timber in, 248; Ctenolepisma spp. in buildings in, 363, 364; parasites and biological control of insects in, 216, establishment of Teleonemia scrupulosa against Lantana in, 181.

auercetorum, Haltica. Quercus (see Oak).

Quercus ilex var. suber, Lymantria dispar on, in Portugal, 285.

quercus, Dendrotettix.

Quicklime, wall washes of insecticides and, 139.

Quince, Quadraspidiotus perniciosus on, in Italy, 64; pests of, in U.S.A., 152, 399.

quinquemaculata, Protoparce. quinquepunctatus, Tychius.

R

Radish, pests of, in Britain. 132 : Phyllotreta aerea on, in Canada, 167 Lixus latro on, in Eritrea, 384; Aphids on, in Fiji, 413; Empoasca lybica on, in Palestine, 176; Phyllotreta cruciferae on, in United Provinces, 123; Hylemyia brassicae on, in U.S.A., 350; tainted by benzene hexachloride, 280.

Rainfall, relation of locusts and grasshoppers to, 123, 124, 125, 170, 171, 206, 311, 398; relation of other insects

to, 27, 172, 222.

Raisins (see Fruit, Dried).

Ramburiella turcomana, crops damaged by, in Persia, 387, 388.

Rana esculenta, toxicity of insecticides to, 174; tadpoles of, little affected by rice-field treatment with DDT, 347.

rapae, Pieris.

Rape, Athalia rosae on, in Italy, 192; Entomoscelis adonidis on, in Rumania, 40; pests of, in Switzerland, 281; scorched by tar distillates, 41.

rapidus, Adelphocoris.

rapo, Tetrastichus.

Raspberry, pests of, in Belgium, 278, 279; Otiorrhynchus singularis on, in Britain, 358; pests of, in U.S.A., 399. Raspberry Root Borer (see Pennisetia marginata).

Rattlepod (see Crotalaria mucronata).

ratzburgiana, Zeiraphera. rectangulata, Chloroclystis.

recurvalis, Hymenia.

Recurvaria, parasite of, on spruce in Quebec, 321.

Red-banded Leafroller (see Eulia velu-

Red-legged Grasshopper (see Melanoplus femur-rubrum).

Reed Grass (see Miscanthus japonicus). Remigia repanda (see Mocis). reniculella, Dioryctria,

Rentokil, 414.

repanda, Coccinella; Mocis (Remigia).

Resin, in sprays, 131, 139. reticulana, Tortrix (Capua).

Réunion, Celosterna scabrator in, 317.

Reuteroscopus spp., bionomics of on cotton in Oklahoma, 183, 184.

Reviews:—Alkan (B.), Agricultural Entomology, 3; Barnes (H. F.), Gall Midges of economic Importance (vol. III), 308; Brues (C. T.), Insects and Human Welfare (revd. edn.), 427; Cerný (J.) & Drachovská-Simanová (M.), Beet Phytopathology, Chevalier (A.), Les caféiers du globe (fasc. III), 405; Dabrowski (Z.), Kamiński (E.) & Ruszkowski (J)., Ed., Manual of Plant Protection, 117; Dethier (V. G.), Chemical Insect Attractants and Repellents, 298; Fennah (R. G.), The Insect Pests of Food-crops in the Lesser Antilles, 71; Frear (D. E. H.), A Catalogue of Insecticides and Fungicides (vol. I), 297; Gabotto (L.), The Control of Diseases and Insect Pests of Crops (6th edn.), 355; Herklots (G. A. C.), Vegetable Cultivation in Hong Kong, 3; Langford (G. S.), Ed., Entoma, a Directory of Insect and Plant Pest Control (7th edn.), 347; Lepesme (P.) and others, Les insectes des palmiers, 177; Lepigre (A. L.), Technique de la désinsectisation, 297; Lhoste (J.), Ce qu'il faut savoir des maladies des plantes cultivées et de leurs ennemis, 105; Miles (H. W.) & Miles (M.), Insect Pests of Glasshouse Crops, 173; Noirot (C.) & Alliot (H.), La lutte contre les termites, 317; Oxley (T. A.), The Scientific Principles of Grain Storage, 405; Peterson (A.), A Manual entomological Equipment Methods (5th edn.), 9; Piédrola Gil (G.), New Insecticides and Repellents, 60; Quidet (P.), Les parasites animaux du tabac en France, 405; Smith (K. M.), Plant Viruses, 312; Smith (K. M.), A Textbook of agricultural Entomology (2nd edn.), 357; Snyder (T. E.), Our Enemy the Termite (revd. edn.), 317; Thompson (W. R.), Ed., A Catalogue of the Parasites and Predators of Insect Pests, Zimmerman (O. Y.) & Lavine (I.), DDT-Killer of Killers, 9.

Rhabdepyris, parasite of Tribolium spp.

in Senegal, 44.

Rhabdopterus picipes, soil fumigation against, on blueberry in New Jersey, 137.

Rhagium bifasciatum, predacious Prionus coriarius in Britain, 215.

Rhagoletis cingulata, distribution and association of, with Prunus emarginata in California, 185, 186.

Rhagoletis fausta, distribution and association of, with Prunus emarginata in

California, 185.

Rhagoletis pomonella (Apple Maggot), insecticides against, in U.S.A., 59, 100, 324, 335; on astrakhan apple, 335.

Rhagoletis suavis completa, on peach in California, 399.

rhamni, Aphis.

Rhamnus crenatus, insecticidal properties of, 299.

Rhaphidopalpa (see Aulacophora).

Rhizaspidiotus dearnessi, on Parthenium argentatum in Mexico, 97.

Rhizoglyphus echinopus, on sweet-potato

tubers in Fiji, 266.

Rhizopertha dominica (in stored cereals), in Australia, 182, 217, 364, 365; in Switzerland, 38, 39; factors affecting infestation by, in bulk wheat, 182, 364, measures and experiments against, 19, 20, 28, 38, 39, 139, 182, 217.

Rhodesia, Southern, Trypetid on cucurbits in, 176; Teleonemia australis on olive in, 74; pests and diseases of olive in, 74; pests and diseases of tobacco in, 137, 422.

Rhododendron (see Azalea).

Rhododendron molle, insecticidal action and constituents of, 298, 299.

Rhodoneura myrsusalis, bionomics of, on Acras zapota in Ivory Coast, 67.

Rhoecocoris sulciventris, on Citrus in New South Wales, 108.

Rhopalosiphoninus ribesina, distribution of, on currant, 408.

Rhopalosiphum crataegellum, flights of, in England, 196.

Rhopalosiphum maidis (see Aphis).

Rhopalosiphum nymphaeae, on Eichhornia in Fiji, 413; sprays against, on plum in Spain, 199.

Rhopalosiphum prunifoliae (padi, auct.), flights of, in England, 196.

Rhopalosiphum pseudobrassicae, experiment with Leis conformis and, in Egypt, 411; on radish in Fiji, 413; dusts against, on crucifers in U.S.A.,

Rhopobota naevana, dust against, on cranberry in Massachusetts, 100.

Rhothane (see DDD).

Rhyacionia buoliana, new parasite of, in Argentina, 109.

Rhyacionia frustrana (Pine Tip Moth), bionomics and control of, in Oklahoma, 158, 388.

Rhynchites, characters distinguishing Byctiscus from, 198.

Rhynchites aeneovirens (see R. minutus).

Rhynchites aequatus, bionomics of, on apple in Britain, 358.

Rhynchites coerulus, bionomics of, on fruit trees, etc., in Britain, 358.

Rhynchites germanicus, bionomics and control of, on small fruits in Belgium and Britain, 279, 358.

Rhynchites minutus, in Belgium, 353; in Britain, 358; on strawberry, 353; not feeding on strawberry, 358.

Rhynchites pauxillus, bionomics and control of, on apple in Belgium, 279.

Rhynchospermum, Euploea core corinna on, in New South Wales, 344.

Rhynocoris iracundus, predacious on Leptinotarsa decemlineata in Spain,

Rhyssa persuasoria, parasite of Sirex noctilio in New Zealand, 244.

ribesii, Nematus (Pteronidea). ribesina, Rhopalosiphoninus.

ribi-gillettei, Aphis. ribis, Cecidophyes.

Rice, new Curculionid on, in Brazil, 8; Chilo loftini on, in California, 147; in

Fiji, 267, 390, 413; Marasmia venilialis on, in India, 267; Nymphula nymphaeata on, in Italy, 346, 347; pests of, in Mauritius, 210; Sesamia on, in Solomon Is., 390; tests of effect of tests of effect of benzene hexachloride on, 13, 413.

Rice (Stored), observations on Ephestia elutella in, 265.

Rice Stinkbug (see Solubea pugnax).

Rice Weevil (see Calandra orvzae). ricini, Asphondylia; Lygus.

Ricinus communis (see Castor).

ridibundus, Hemiteles.

rileyi, Pyroderces.

Rio de Oro, Schistocerca gregaria in, 206. Robinia, Eulecanium coryli on, in Czechoslovakia, 65; pests of, in U.S.A., 143, 144.

Robinia pseudacacia, Etiella zinckenella on, in Hungary, 63; relation of insects to, in Virginia, 373, 374.

robiniae, Megacyllene (Cyllene).

Rodolia cardinalis (predacious on Icerya purchasi), effect of insecticides on, in California, 145, 160; introduction and use of, in Persia, 387.

Roduniella, on Eleusine in Uganda, 367. Rogas testaceus, bionomics of, parasitising

Earias spp. in Punjab, 121.

rosaceana, Tortrix (Archips). rosae, Athalia; Macrosiphum; Psila; Typhlocyba.

rosae, auct., Arge (see A. ochropa).

Rose, Anthonomus rubi on, in Belgium, 278; pests of, in Britain, 288, 313; Adoretus versutus on, in Fiji, 266, 390; pests of, in Italy, 253, 254; Chrysomphalus dictyospermi on, in Persia, 386; pests of, in greenhouses in Switzerland. 282; pests of, in U.S.A., 14, 103, 154, 335; insecticides and injury to, 103, 284.

Rose Chafer (see Macrodactylus subspinosus).

roseus, Anuraphis.

Rotenoids, and pyrethrins, tests of proprietary preparation of, on Brevicoryne brassicae, 300.

Rotenone, against Aphids, 24, 154, 192, 218, 229, 282, 335, 371; against Coleoptera, 14, 41, 62, 101, 225, 229, 230, 330, 335, 369, 370, 396; effect of, on Rodolia cardinalis, 160; ineffective against Empoasca lybica, 176; against Lepidoptera, 229, 230, 282, 369; against Mirids, 102, 272, 335, 394, 395; against other pests, 282, 335; tests of toxicity of, to bees, 222; comparison of action of, on insects and vertebrates, 174, 175; in aerosols, 230; in dusts, 14, 24, 41, 62, 101, 102, 154, 218, 225, 229, 230, 282, 330, 370, 371, 394, 395, 396 : oil-impregnated dusts containing, 229; factors affecting efficiency of dusts containing, 330; in sprays, 160, 192, 225, 272, 282, 335; applied from aircraft, 396; applied by bloweratomisers, 36; mixed with stored seeds, 369; formulae containing, 102, 154, 218, 225, 229, 230, 282, 371, 395; and benzene hexachloride, 154, 229; and calcium arsenate, 371; and copper carbonate, 282; and DDT, 154, 218, 229, 230, 371, 395; and Lethane B-71, 395; and oil emulsions, 160, 282; and P-C-H, 394; and sulphur, 24, 229, 230; review of properties and uses of, 297; not present in Amorpha frutiosa, 9; in Millettia pachycarpa, 298. (See Cubé, Derris and Pachyrhizus.)

Rothane (see DDD). rotundatus, Chrotogonus. Rowan (see Sorbus aucuparia). Rubber (see Hevea). rubi, Anthonomus. rubicunda, Anisota. rubigalis, Phlyctaenia. rubiginosa, Lachnosterna (Phyllophaga). rubripennis, Myelophilus (Blastophagus) piniperda.

Rucide, 179, 217, 218.

Rudbeckia fulgida, Homoeosoma electellum on, in Iowa, 84.

rufa, Formica; Solenopsis geminata.

ruficollis, Dysdercus.

ruficoma, Bucculatrix.

ruficornis, Andrector (Cerotoma).

ruficrus, Apanteles.

rufipes, Harpalus; Hylastes (Hylurgopinus); Megaselia; Melanotus; Necrobia; Phytomyza.

rufoniger, Iridomyrmex.

ruforum, Achrysocharella.

rufus, Lophyrus (see Neodiprion sertifer). rugicollis, Minthea; Plesiocoris,

rugosa, Lachnosterna (Phyllophaga).

rugulosum, Eulecanium. rugulosus, Scolytus.

Rumania, miscellaneous pests in, 40, 41; new Torymid on juniper in, 267. rumicis, auct., Aphis (see A. fabae).

Russian Thistle (see Salsola).

russoi, Mabra. Ryanex, 402.

Ryania, tests of preparations of,

insecticides, 12, 13, 335, 402. Rye, effect of wireworms on yield of, in Britain, 381; Calocoris norvegicus on, in Italy, 383; pests of, in U.S.A., 96,

183; Aphid on, in Uruguay, 49; as cover crop for tobacco against wireworms, 183.

S

Sabacide, 325, 326.

Sabadilla, against Coleoptera, 370; against Lepidoptera, 100, 331; against Mirids, 11, 91, 184, 235, 395; against other Hemiptera, 100, 134, 235, 325, 326, 335; in dust barrier against Blissus leucopterus, 154; toxicity of, to bees, 222; dusting with, 11, 91, 100, 134, 184, 222, 331, 370, 395; in sprays, 235, 325, 335; formulae containing, 11, 91, 100, 184, 222, 235, 325, 335; with Bordeaux mixture and oil, 325; reducing yield of cotton, 184.

Sabal bermudiana, Coccid on, in Bermuda, 49.

sabalis, Comstockiella.

sabulosum, Opatrum.

saccharalis, Diatraea.

sacchari, Perforadix; Trionymus (Pseudococcus).

Saccharin, Anastatus disparis unable to survive on, 117.

saccharina, Lepisma; Tomaspis.

sacchariphagus, Proceras.

Sagaritopsis (Sagaritis) grioti, sp. n., parasite of Thyreion gelotopoeon in Argentina, 109.

Sainfoin (see Onobrychis sativa).

St. John's Wort (see Hypericum perfora-

Saipan, Dacus ferrugineus dorsalis probably introduced into Hawaii from, 249.

Saissetia oleae, in Australia, 74, 177, 216; in Persia, 385; in Trinidad, 319; in U.S.A., 101, 342; on Citrus, 74, 177; on olive, 342; on other plants, 101, 319, 385; natural enemies attempted biological control of, 216, 385; other measures and experiments against, 74, 101, 342, 385. salmani, Coleophora.

Salsola, 99; Eutettix tenellus on, in California, 186, 397, 398.

Salt (Sodium Chloride), not protecting stored beans from Bruchus obtectus, 21; survival of Tyroglyphus nadinus in solution of, 114; effect of, in oil emulsions, 361.

saltensis, Atta (Neoatta) laevigata.

Sambucus, favouring infestation of peach by Acrosternum hilare in Virginia, 373. samiae, Achaetoneura.

Samoa, Pentalonia nigronervosa bunchy-top of banana in, 267; Hyblaea puera on teak in, 267.

sanctae-helenae, Heteronychus.

Sanninoidea exitiosa (see Aegeria). Santomerse D, 224, 226, 228.

Saperda tridentata (see Eutetrapha). Sappaphis (see Anuraphis).

saratogensis, Aphrophora.

Sarcophaga atlanis, parasite of Dendrotettix quercus in New Jersey, 302.

Sarcophaga caridei, parasite of Schisto-cerca paranensis in Argentina, 133; introduced into Peru, 133.

Sarcophaga lambens (sternodontis), parasite of Sacadodes pyralis in Trinidad, 126; bionomics and distribution of, 126.

sauzalitae, Emboloecia.

Saw-toothed Grain Beetle (see Oryzaephilus surinamensis).

Sawdust, in baits for grasshoppers, 305. Sawfly, Apple (see Hoplocampa testudi-

Sawfly, Cherry (see Caliroa limacina). Sawfly, European Spruce (see Gilpinia hercyniae).

Sawfly, Leconte's (see *Neodiprion lecontei*). Sawfly, Mustard (see Athalia proxima). Sawfly, Wheat Stem (see Cephus cinctus). saxeseni, Xyleborus (see X. xylographus). scabrator, Celosterna (Coelosterna).

scalaris, Megaselia.

Scale, California Red (see Aonidiella aurantii).

Scale, European Elm (see Gossyparia spuria).

Scale, Florida Red (see Chrysomphalus ficus).

Scale, Olive (see Parlatoria oleae).

Oyster-shell (see *Lepidosaphes* ulmi).

Scale, Tulip-tree (see Toumeyella liriodendri).

Scale Insects, natural enemies of, 49, 54, 64, 65, 70, 83, 109, 111, 145, 155, 156, 172, 191, 315, 385, 386, 387, 389, 392, 404, 410, 411; lists of parasites and predators of, 270; ants associated with, 74, 110, 111, 404; transmitting cacao viruses, 111; classification and new species of, 69, 110, 316, 401.

Scapteriscus vicinus, measures against, damaging vegetables, etc., in Porto Rico and Georgia, 53, 93, 94.

Sceloporus graciosus, destroying ants in Utah, 24.

schauffussi, Formica pallidefulva.

Schematiza cordiae, proposed introduction of, into Mauritius against Cordia macrostachya, 424; natural enemies and effects on C. macrostachya of, in Trinidad, 424-426.

Schistocerca cancellata, bionomics and control of, in Chile, 108, 109.

Schistocerca gregaria, in eastern Africa, 123–125, 366, 375, 403; in western and north-western Africa, 205, 206, 207; in Egypt, 412, 413; in India, 122, in Persia, 387; 310-312; invading Portugal and Spain, 205; seasonal breeding and migrations of, 123-125, 205-207, 311, 366, 403; effects of temperature on flight of, 205, 375; measurements of body temperature of, 375; other bionomics of, 122, 311; Pastor roseus destroying, 312; measures and experiments against, 403, 412, 413.

Schistocerca paranensis, outbreak of, in Peru, 133; introduction of Argentine parasite against, 133; other measures and experiments against, 133, 134.

Schizanthus, injured by sodium selenate,

Schizonycha, soil treatments against, in tobacco fields in S. Rhodesia, 137.

schlechtendali, Vasates (Phyllocoptes). Schleichera trijuga, host-plant of Laccifer lacca in Bihar, 120.

schoenei, Tetranychus.

schultzei, Frankliniella.

Sciara, insecticides against, on mushrooms in Britain and New South Wales, 284, 365.

Scirpophaga nivella, parasite of, on sugarcane in India and Burma, 119.

Scirtothrips, injury to mango by Aceria mangiferae attributed to, in Egypt, 409. Scirtothrips longipennis, insecticides

against, on Cinchona in Porto Rico, 53.

Scirtothrips signipennis, insecticides against, on banana in Queensland, 306. scitella, Leucoptera (Cemiostoma).

Scolytids (see Bark-beetles).

Scolytus, of Palaearctic Region, 348.

Scolytus multistriatus, not found in Quebec, 320; in U.S.A., 15, 103, 144, 320; transmitting Dutch elm disease, effect of phloem condition on infestation of elm logs by, 15; sprays protecting elm trees and logs against, 103, 144.

Scolytus rugulosus (on fruit trees), in Britain, 358; measures against, in California, 399; factors infestation by, in Italy, 192. factors favouring

scrupulosa, Teleonemia. scutellare, Xanthogramma (Ischiodon). scutellatus, Gonipterus; Monochamus.

Scyllina signatipennis, bionomics of, in Chile. 109.

Scymnus, Stethorus bifidus recorded as, in New Zealand, 427.

Scymnus fenestratus, predacious on mealybugs in Palestine, 410.

Scymnus gracilis, Stethorus pauperculus recorded as, in India, 427.

Scymnus includens, predacious on mealybugs in Palestine, 410.

Scymnus minutulus, 427.

Scymnus punctillum (see Stethorus).

Scymnus suturalis, predacious on mealybugs in Palestine, 410.

Scyphophorus acupunctatus (Sisal Weevil), intercepted in Kenya from Tanganyika, 405.

Seal Oil, in mixture for treating timber against pests, 107.

segetum, Agrotis (Euxoa); Anisoplia. Seius pomi, predacious on Eriophyids, 82.

Selatosomus aeneus (see Corymbites).

Selenaspidus articulatus (on Citrus), measures to prevent introduction of, into California from Mexico, 79; DDT ineffective against, in Peru, 134; in Trinidad, 319.

Selenephera lunigera, identification of excreta of, on conifers in Germany,

seleniticus, Lasiophthicus.

Selenium, soil treatment with selenates for introduction of, into plants, 289-291; risk of poisoning from vegetables containing, 290.

semifumipennis, Uscana.

semipullata, Galerucella (Galeruca).

semipunctata, Lissopimpla.

senio, Phoracantha.

sentis, Urentius.

septemfasciata, Nomadacris. septempunctata, Coccinella

sepulchralis, Hysteropterum.

seriatus, Psallus.

sericeus, Polydrusus,

serpentina, Anastrepha. serricorne, Lasioderma.

sertifer, Neodiprion. servus, Euschistus.

Sesamia, on rice in Solomon Is., 390.

Sesamia inferens, 390.

Sesamia vuteria, sugar-cane Mauritius, 210.

Sesamum orientale (indicum), 310.

Setaria, Cirphis unipuncta on variety of, in Queensland, 307.

setosus, Pseudocneorrhinus.

sexannulatus, Cryptus.

sexdens, Atta (Neoatta).

sexmaculatus, Tetranychus.

sexta, Protoparce.

Shallot, Hylemyia antiqua on, in N. Dakota, 388.

sheldoni, Aceria.

Sida spp., Bemisia tabaci transmitting virus of, in Brazil, 299, 300; Pantomorus godmani on, in New South Wales, 345; Earias spp. on, in Punjab, 312.

Sideridis unipuncta (see Cirphis).

Sierra Leone, Schistocerca gregaria migrating over, 207; Trypetid on cucurbits in, 176; parasite of Dysdercus in, 27. Sigalphus (see Triaspis).

signata, Chlorita (see Empoasca lybica).

signatipennis, Scyllina. signipennis, Scirtothrips.

Silica (Silicon Dioxide), in repellent spray against *Popillia japonica*, 336; forms of, as dust diluents, 329.

Silicates, survey of types of, as dust diluents, 329.

Silicon Dioxide (see Silica).

Silkworm (see Bombyx mori).

Silkworm, Tussore (see Antheraea pernyi). similalis, Loxostege.

similis, Aulacophora; Euproctis (Phalaena); Perilampus.

simmondsi, Adelencyrtus.

simonvi, Lygus.

simplex, Dasus (Gonocephalum); Lissorhoptrus; Taeniothrips.

Sinapis alba (see Mustard).

Sind, Schistocerca gregaria in, 310–312. sinensis, Ceroplastes.

singularis, Otiorrhynchus.

Sipha flava, on sugar-cane in Br. Guiana, 304.

siphonophorae, Pachyneuron.

Siphophyto floridensis (see Epigrimyia). Sirex noctilio, use of Rhyssa persuasoria

against, in pine in New Zealand, 244. Sisal Weevil (see Scyphophorus punctatus).

Sitona cylindricollis, DDT against, on

Melilotus in U.S.A., 81.

Sitona lineatus, DDT against, on peas and beans in Belgium, 291.

Sitophilus (see Calandra).

Sitotroga cerealella (in stored cereals), tests of fumigants against, in Italy, 30, 38; inert dusts ineffective against, in Mauritius, 210; in Uganda, 367; use of eggs of, for rearing Trichogramma minutum, 200.

Slugs, destroying Antholcus varinervis in New Zealand, 244.

smithi, Clemora.

Smoke, repellent to Anastrepha ludens,

Smynthurus viridis, utilisation of Biscirus lapidarius against, in Tasmania, 70.

Soap, spraying with, 52, 75, 176; as an emulsifier, 133, 139; other uses of, in mixed sprays, 4, 19, 50, 52, 75, 102, 104, 199, 218, 247, 286, 305, 327, 334; types of, 218, 247, 50, 102, 104, 199, 218, 247, formulae contains 176, 199, 218, 247; formulae containing, 50, 52, 75, 133, 139, 199, 247, 327: for removing DDT deposits from apples, 135.

Sodium Alkyl Naphthalene Sulphonate,

Wetsit (q.v.) containing, 69.

Sodium Arsenate, in bait for moths, 294; against termites in timber, 169; -Tanalith (q.v.) containing, 414.

Sodium Arsenite, in baits, 4, 74, 294; benzene hexachloride less toxic to domestic animals than, 413.

Bicarbonate, not protecting

stored beans against Bruchus obtectus, 21.

Sodium Carbonate, to prevent precipitation of DDT wax in sprays, 246.

Sodium Chloride (see Salt).

Sodium Chromate, Tanalith (q.v.) con-

taining, 414.

Sodium Cyanide, and sulphuric acid, for generating hydrocyanic acid gas (q.v.), 30, 80, 361, 397.

Sodium Decylbenzenesulphonate, wetting agent for sprays, 224, 226, 228.

Sodium Dinitro-o-cresylate, 111, 272,

Sodium Fluoaluminate (see Cryolite).

Sodium Fluoride, humidity reducing toxicity of dust of, to Lepisma saccharina, 379; DDT less toxic to man than, 162; Tanalith (q.v.) containing, 414.

Sodium Fluosilicate, in baits, 96, 186; effect of carriers on toxicity of dusts of,

to insects, 113.

Sodium Lauryl Sulphate, use of, to remove DDT residues from apples, 135; as an emulsifier, 135.

Sodium Pentachlorphenate, repellency of, to Kalotermes brevis, 233.

Sodium Pentobarbitol, treatment with, of dogs poisoned by Toxaphene, 396.

Sodium Selenate, plants grown in soil containing, against pests, 289, 290, 291.

Sodium Silicate, and Bordeaux mixture. 42.

solani, Epicauta; Macrosiphum (Myzus). solanifolii, Macrosiphum.

Solanum ciliatum, low insecticidal properties of, 52.

Solanum melongena (see Egg-plant).

Solanum rostratum, Paratrioza cockerelli on, in U.S.A., 144.

Solanum tuberosum (see Potato).

Solar Heat, use of, against Scolytids in logs, 103.

Solenopsis geminata rufa, predacious on Eublemma amabilis in Bihar, 120.

Solenopsis molesta, effect of sprays on, destroying Cydia pomonella in W. Virginia, 220.

Solidago, Odynerus tigris nesting in galls on, in Ontario, 167.

Solomon Is., miscellaneous pests in, 267, 390.

Solubea pugnax (Rice Stinkbug), benzene hexachloride against, in U.S.A., 13.

Somalia, Schistocerca gregaria in, 123, 124, 403. Somaliland, British, Schistocerca gregaria

in, 123, 124. Somaliland, French, Schistocerca gregaria

in, 123; Tribolium indicum in, 261. Sonchus oleraceus, destruction of, against

Hercothrips fasciatus in California, 399.

Sooty Mould, associated with Coccids, 293, 387; associated with Hyalopterus arundinis, 192.

Sophora flavescens, insecticidal properties of, **299**.

Sorbus aucuparia, pests on, in Britain, 314, 358; Argyresthia conjugella and, in Sweden, 409.

sordidus, Aphanus; Cosmopolites.

Sorghum, 307: Chilo loftini on, in California and Mexico, 147; pests of, in India, 122, 427; Lygus spp. on, in Uganda, 26.

Sorghum sudanense, Cirphis unipuncta on,

in Oueensland, 307.

Sostroil, in mixture for treating timber against pests, 107.

Sour Clover (see Melilotus indica).

Southern Corn Rootworm (see Diabrotica duodecimpunctata).

Sow Thistle (see Sonchus oleraceus).

Soy Bean, not attacked by Athalia rosae in Italy, 192; pests of, in Peru, 134; pests of, in U.S.A., 18, 228; varieties of, in relation to Popillia japonica, 18; insecticides and injury to, 228, 338; (stored); Bruchus analis in, in Uganda,

Soybean Flour, studies on development of pests of stored products in, 265, 348; in food for rearing clothes moths and carpet beetles, 343; uses and effects of, in sprays, 54, 127, 323, 335, 354,

372, 373.

Soybean Grits, susceptibility of, to infestation by Tribolium confusum, 348. Soybean Oil, uses of, in sprays, 54, 335,

372.

Soybean Phosphatides, in lead-arsenate sprays, 372.

Spain, Earias insulana on cotton in, 197; pests of fruit trees and vines in, 196, 198, 199; locusts in, 202, 205; pests of lucerne in, 203, 293; pests of olive in, 201, 385; pests of vegetables and root crops in, 196, 199, 202, 204; Pediculopsis graminum and fungus infesting wheat in, 202; Thylodrias contractus damaging insect specimens in, 37; beneficial insects in, 196, 197, 200; studies on benzene hexachloride and DDT in, 196, 198, 199, 203, 204, 215, 293.

Spalangia afra, parasite of Dacus in Zanzibar, 177.

sparsa, Aleochara.

Speedsprayer, 322, 323.

spelaea, Coccidella.

Spermophagus subfasciatus, in imported beans in Poland, 385.

Sphaeropsis malorum, not causing leaf spotting of apple in California, 401.

Sphaerosicyos sphaericus, Trypetid in flowers of, in S. Africa, 176.

Sphinx pinastri, identification of excreta of, on pine in Germany, 75, 76.

Spiders, destroying insects, 172, 212, 244, 268, 272; toxicity of thiocyanoacetates to, 253.

Spilochalcis, parasite of Diatraea in Br. Guiana, 304; hyperparasite of Schematiza cordiae in Trinidad, 425.

Spilonota ocellana (on apple), in Canada, 167, 271, 392; in Holland, 276; Eumenid destroying, 167; tests of effects of sprays on, 271, 276.

Spilostethus pandurus, damaging fruits in

India, 123.

Spinach, Haplomyza imitans on, in New South Wales, 73; pests of, in U.S.A., 24, 331; injured by Velsicol AR-60, 421.

Spineless Cactus (see Opuntia).

Spiraea ulmaria, Cladius pectinicornis on, in Belgium, 279.

Spiraea vanhouttei, Aphis spiraecola on, in U.S.A., 335.

spiraecola, Aphis.

splendida, Lonchaea (see L. aurea).

Spodoptera cilium, damaging lawns in Mauritius, 210; disease of, 210. Spodoptera exempta (see Laphygma).

Spodoptera mauritia, in Formosa, 177; destroying grass in Tonga, 267; parasites of, 177, 267.

Sprays, application of, from aircraft, 17, 33, 34, 35, 36, 92, 99, 146, 168, 228, 260; observations on deposits of, from aircraft, 17, 260; equipment for applying, 33, 34, 35, 36, 92, 131, 168, 178, 260, 263, 326; droplet size in relation to distribution and deposits of, 145, 146; equipment for testing, 341, 420; tests with French commercial adjuvants for, 42; dusts compared with, 63, 69, 165, 227, 363.

Spruce, Lymantria monacha on, in Austria, 128; pests of, in Britain, 315; pests of, in Canada, 33, 78, 240, 255, 319, 320, 321; pests of, in Germany, 75, 76; Scolytids on, in Switzerland, 407, 408.

Spruce, Black (see Picea mariana).

Spruce, White (see *Picea glauca*). Spruce Budworm (see *Harmologa fumi*ferana).

Spruce Sawfly, European (see Gilpinia hercyniae).

spumosum, Stibadium.

spuria, Gossyparia.

sputator, Agriotes.

Squash, pests of, in U.S.A., 94, 101, 370; susceptibility of varieties of, to Melittia cucurbitae, 101; effects of insecticides on, 19, 94, 225, 235; DDT mixed with stored seed of, against insect pests, 19.

Squash Bug (see Anasa tristis).

Squash Vine Borer (see Melittia cucurbitae).

stabulans, Muscina.

stali, Idiocerus.

Standesol, 414.

Starch, depletion of, from wood and trees against *Lyctus*, 1, 248.

Stathmopoda auriferella, on mango in

Egypt, **409**.

Statistical Methods, use of, in estimating infestations by *Stephanoderes hampei*, 5; application of, to insecticide tests, 381.

Steel, Stainless, suitability of, for containers for Toxaphene, 395.

steinwedeni, Acaphylla.

Stellaria media, Haplomyza imitans on, in New South Wales, 73.

stellata, Lyda (see Acantholyda pinivora).

stemmator, Xanthopimpla.

Stemphylium, aggravating lead-arsenate spray injury to apple, in California, 401.

Stemphylium congestum, 401.

Stenoma decora, bionomics of, on cacao and kapok in Brazil, 125, 126.

Stenomalina, n.n. (Stenomalus, Thoms.), 316.

Stephanoderes, on coffee in Porto Rico, 300.

Stephanoderes hampei, statistical method of estimating infestation of coffee by, in Brazil, 5.

Stephanotis, Euploea core corinna on, in New South Wales, 344.

sternodontis, Sarcophaga (see S. lambens). Stethorus, feeding habits and lists of species of, 426, 427.

Stethorus bifidus, sp. n., predacious on Paratetranychus pilosus in New Zealand, 427.

Stethorus gilvifrons, predacious on mites in India, 427.

Stethorus pauperculus, predacious on mites in India, 427.

Stethorus picipes, predacious on Tetranychus pacificus in Washington, 150; effect of DDT on, 150.

Stethorus punctillum, predacious on Tetranychus carpini in Switzerland, 282. Stethorus tetranychi, sp.n., predacious on

Tetranychus in Bengal, 427. Stibadium spumosum, parasites of, on

sunflower in U.S.A., 84. sticticalis, Loxostege (Phlyctaenodes).

Stictoptera subobliqua, on Calophyllum inophyllum in Fiji, 266.

stigma, Chilocorus.

Stomatodexia flavipennis, parasite of Diatraea in Br. Guiana, 304.

Strawberry, pests of, in Belgium, 278, 279, 292, 353, 354; pests of, in Britain, 313, 314, 358, 359, 376; *Lachnosterna*

damaging, in Ontario, 164; Agrotis ypsilon on, in Queensland, 308; Cercopids on, in Wisconsin, 160; Aphids and virus diseases of, 312–314, 376; warm water treatment of, against Tarsonemus pallidus, 359; benzene hexachloride not affecting flavour of, 234.

Strawberry Aphis (see Capitophorus fragariae).

Strawberry Crown Moth (see Aegeria bibionipennis).

striana, Tortrix (Archips).

strobi, Chermes (Pineus); Pissodes.

Strophanthus divaricatus, insecticidal properties of, 299.

Strumeta (see Dacus).

Strychnine, ineffective against Agriotes, 286, 287; difference in toxicity of, to insects and vertebrates, 175.

Sturmia, use of, against Gilpinia hercyniae in Canada, 208, 320, 321.

Sturmia inconspicua, parasite of Pompostola hypparchus in Ivory Coast, 67.

Styrene Dibromide, in oil, treatment of maize ears with, against *Heliothis armigera*, 370.

subcoleoptrata, Alophora. subfasciatus, Spermophagus.

subfumatus, Amblymerus.

submucida, Lachnosterna (Phyllophaga).

subnigra, Acutaspis. subobliqua, Stictoptera. subsericea, Formica fusca. subspinosus, Macrodactylus.

Subterranean Clover (see Trifolium subterraneum).

Sudan, Anglo-Egyptian, Anychus latus in, 412; Schistocerca gregaria in, 123, 124, 207, 366; new form of Tribolium indicum in, 261; parasite of Dysdercus in, 27.

Sudan Grass (see Sorghum sudanense).

Sugar, as larval attractant in sprays against *Cydia pomonella*, **296**; in bait-sprays for Diptera, **46**, **47**, **148**, **248**; in sprays against thrips, **53**, **332**; used for feeding insects, **29**, **117**, **120**, **126**, **133**, **148**, **241**; *Tyroglyphus nadinus* unable to survive on, **114**.

Sugar Pine (see *Pinus lambertiana*).

Sugar-cane, pests of, in Br. Guiana, 302, 303, 304, 305; Scirpophaga nivella on, in Burma, 119; Pseudococcus boninsis on, in Hawaii, 316; pests of, in India, 119, 122; pests of, in Mauritius, 209, 210; Chilo loftini on, in Mexico, 146; pests of, in U.S.A., 147, 316; pests of, in W. Indies, 29, 53, 240, 302, 303, 318, 319.

Sugar-cane Aphis, Yellow (see Sipha flava).

Sugar-cane Froghopper (see *Tomaspis saccharina*).

Sugar-cane Froghopper, Demerara (see *Tomaspis flavilatera*).

Sugar-cane Moth Borers (see *Diatraea* saccharalis, *Proceras* sacchariphagus and Sesamia vuteria).

sulcatellus, Campoplex.

sulcatus, Otiorrhynchus (Brachyrhinus). sulcirostris, Panscopus (see Agasphaerops nigra).

sulciventris, Rhoecocoris.

Suleima helianthana, natural enemies of, on Helianthus spp., in U.S.A., 84.

Sulphite Lye, as an emulsifier, 131. Sulphonated Lorol TA, 131.

Sulphonated Petroleum Oil, as adhesive

and spreader for lead arsenate, 143. Sulphur, against Coleoptera, 137, 146, ineffective against Empoasca 338; lybica, 176; against Mirids, 184, 394; against mites, 53, 82, 180, 181, 239, 249, 289, 362, 363, 393; favouring Paratetranychus pilosus and Lepido-saphes ulmi, 271, 392, 393; against Paratrioza cockerelli, 231; insects attacking fleeces of goats dipped in, dusting with, 53, 82, 184, 239, 338, 394; in mixed dusts, 11, 12, 21, 24, 47, 57, 58, 59, 91, 102, 126, 148, 149, 184, 187, 229, 230, 231, 232, 237, 250, 266, 282, 329, 338, 342, 362, 363, 394, 395; spraying with, 82, 249, 392, 393, 394; in mixed sprays, 58, 62, 101, 165, 180, 181, 224, 225, 324, 354, 362, 363; soil treatment with, 137, 148, 149; forms of, 57, 137, 180, 225; formulae containing, 11, 21, 47, 57, 91, 126, 146, 148, 165, 180, 181, 225, 229, 230, 231, 232, 250, 266, 324, 338, 342, 362, 363, 394, 395.

sulphureus, Reuteroscopus.

Sulphuric Acid, survival of *Tyroglyphus* nadinus in, 114; and sodium cyanide, for generating hydrocyanic acid gas (q.v.), 30, 80, 361.

Sumba, on Eleusine in Uganda, 367.

Sunflower (see Helianthus).

Sunflower-seed Oil, emulsions of, as insecticides, 354.

superstitiosus, Dysdercus.

surinamensis, Oryzaephilus.

suturalis, Scymnus.

Sweden, miscellaneous pests in, 408, 409.

Sweet Clover (see Melilotus).

Sweet Potato, new weevil on, in S. Africa, 388; *Trichoptilus wahlbergi* on, in Mauritius, 210; *Bucculatrix* on, in Nyasaland, 44; pests of, in Pacific Is.,

266, 267, 390; Empoasca lybica on, in Palestine, 176; Megastes grandalis on, in Trinidad, 319; Acraea acerata on, in Uganda, 367.

Sweet Potato Weevil (see Cylas formi-

carius).

Switzerland, miscellaneous pests in, 279, 281, 296; orchard pests in, 273, 296; pests of ornamental plants in, 60, 282; Scolytids on spruce in, 407, 408; pests of stored cereals in, 38, 39; beneficial insects in, 273, 282.

syleptae, Eurytoma.

Syncarpia laurifolia (timber of), Phoracantha senio intercepted in, in Fiji, 267; repellent to Kalotermes brevis, 169.

Syntomosphyrum indicum, use of, against fruit-flies in Fiji, 390.

Syrphus balteatus, possible overwintering of adults of, in Switzerland, 273.

Syrphus corollae var. vitiensis, predacious on Aphids in Fiji, 414.

T

tabaci, Bemisia; Thrips. tachardiae, Brachymeria.

Tachardiella cornuta, on Parthenium argentatum in Mexico, 97.

Tachardiella ferrisi, on Adenostoma in California, 187.

Tadpoles, effect of DDT on, 347.

Taeniothrips simplex, experiments against, on gladiolus in U.S.A., 234, 291, 331, 332.

tafoensis, Formicococcus.

Tagetes spp., Homoeosoma electellum on, in Iowa, 84.

Tahiti, Psychodes trilineatus insularis in fig in, 250.

taiwana, Euproctis (Porthesia).

taiwanensis, Apanteles. taiwanus, Euplectrus.

Talc, effects of, on insects, 102, 145, 330, 379; as carrier for dusts, 13, 52, 57, 58, 90, 102, 112, 113, 114, 134, 145, 210, 237, 244, 324, 329, 337, 351, 394, 395, 402, 420; electrostatic charges of dusts of, 393; unsuitable as carrier for dinitro-o-cyclohexylphenol, 288; comparison of pyrophyllite with, 340, 341.

Tamarix, Lachnosterna submucida on, in Texas, 14.

tanaceti, Galeruca.

Tanaemyrmex picipes (see Camponotus).

Tanalith, tests of, for treating timber against *Anobium*, **414**, **415**; composition of, **414**.

Tanganyika, pests of field and stored groundnuts in, 426; locusts in, 123, 124, 294, 403; miscellaneous pests in, 26, 176, 316; Scyphophorus acupunctatus intercepted in Kenya from, 405.

Taniva albolineana, parasites spruce in Quebec, 321. of,

Tanymecus palliatus, on castor and beet in Hungary, 62, 63.

tapetzella, Trichophaga (Tinea).

Tar Distillates, as dormant sprays against Lepidoptera on currants and vines, 286, 297; as dormant sprays against orchard pests, 111, 130, 192, 247, 273, 276, 354; ineffective against eggs of Lygidea mendax, 272; Coniopteryx tineiformis not affected by, 359; experiment with, against Entomoscelis adonidis on rape, 41; ineffective against Ips typographus, 408; tests of toxicity of, to eggs of insects, 111, 112, 273, 274, 276; types of, 111, 112; and dinitro-o-cresol, 273, 274.

Tar Paper Disks, use of, against Hylemyia

brassicae on cabbage, 102. Targionia dearnessi (see Rhizaspidiotus). yuccarum, on Parthenium Targionia argentatum in Mexico, 97.

Taro (see Colocasia).

Tarsonemus pallidus (fragariae), measures against, on strawberry in Belgium and Britain, 353, 359.

Tarsostenus univittatus, predacious on Lyctus brunneus in Africa, 1.

Tartar Emetic, and sugar, spraying with, against thrips, 148, 332; effect of sub-lethal doses of, on Anastrepha ludens, 148.

Tasmania, orchard pests in, 70, 245, 246; Thrips tabaci on beans in, 138; biological control of pests in, 70.

TDE (see DDD).

Tea, pests of, in Ceylon, 251, 368; Xyleborus fornicatus on, in Formosa, 251; pests of, in India, 250, 251; X. fornicatus on, in Netherlands Indies, 251; Chrysomphalus dictyospermi on, in Persia, 386.

Tea Aphis (see *Toxoptera aurantii*).

Tea Shot-hole Borer (see Xyleborus fornicatus).

Teak, Hyblaea puera on, in Samoa, 267. tectus, Ptinus.

Tegenaria domestica, destroying Ephestia elutella in Britain, 212.

telarius, Tetranychus.

Telenomus, parasite of Papilio demodocus in Mauritius, 210.

Telenomus alecto (parasite of Diatraea), in Br. Guiana, 304; established in Porto Rico, 53.

Teleonemia australis, bionomics and control of, on olive in S. Rhodesia, 74, 75.

Teleonemia scrupulosa, establishment of, against Lantana camara in New South Wales and Queensland, 181.

Telephone Poles, treatment of, against

insects in S. Africa, 107.

Temperature, relation of locusts grasshoppers to, 125, 205, 275; locusts and tion of other insects to, 8, 41, 56, 102, 116, 119, 120, 121, 130, 142, 144, 175, 182, 183, 196, 214, 222, 243, 264, 265, 273, 371, 390, 391, 427; relation of mites to, 129, 181, 418; effect of, on action of insecticides, 112, 259, 330, 378, 379, 380, 381. (See Cold Storage and Heat.)

tenebricosa, Capnodis.

Tenebrio molitor, absorption of water

from air by, 375.

mauritanicus, in Tenebroides groundnuts and maize in Africa, 44, 367, 426; predacious on Ephestia cautella, 426; tests with benzene hexachloride against, 235.

tenella, Galerucella. tenellus, Eutettix.

tenera, Leskiomima.

Tent Caterpillars (see *Malacosoma*).

tenthredinis, Mesoleius. tenuicornis, Humbe.

tenuimaculatus, Adoretus.

Tenuipalpus granati, sp. n., on vines and pomegranate in Egypt, 412.

Tenuipalpus obovatus (see Brevipalpus). Tenuipalpus orchidarum, new mite re-

corded as, in Egypt, 412. Tenuipalpus oudemansi (see Brevipalpus). Tepary Beans (see Phaseolus acutifolius). Terminalia, new mite on, in Egypt, 411;

(myrobalans), bibliography of pests of stored, 272.

Terminalia superba, new Coccids on, in Gold Coast, 110; Lyctus in timber of, in Africa, 1, 107, 295.

terminalis, Paratetranychus. terminifera, Chortoicetes.

Termites, attacking living plants, 250, 304, 317; attacking textiles, 324; attacking timber, 232, 233, 249, 295, 300, 317, 414-416; ant destroying, 305; measures and experiments against, 232, 249, 300, 305, 317, 324, 404, 414-416; books on control of, 317.

Terne Plate, suitability of, for containers

for Toxaphene, 395.

Terpineol, as activator for pyrethrins, 380, 381.

terrae-reginae, Empoasca.

terricola, Pristonychus (Laemostenus),

terryi, Pseudococcobius. tertialis, Phlyctaenia. testaceicollis, Pristomerus. testaceus, Rogas. testudinea, Hoplocampa.

tetensi, Dasyneura.

Tetrabutyl Pyrophosphate, toxicity of, to Aphids, 228.

Tetracampe diprioni, parasite of Diprion pini in Germany, 239.

Tetrachlorethane, risk of using, as solvent for DDT, 161.

Tetraethyl Pyrophosphate, toxicity of, to Aphids, 228.

Tetrahydrofurfuryl Pentachlorphenyl Ether, 224.

Tetralin, as auxiliary solvent for DDT, 89. Tetramethylthiuramdisulphide, in spray formula, 131.

Tetramorium caespitum, effect of sprays on, destroying Cydia pomonella in W. Virginia, 220.

tetranychi, Stethorus.

Tetranychus, Stethorus tetranychi predacious on, on jute in Bengal, 427; Tydeus womersleyi recorded as, on Albizzia in Ceylon, 369; in Kirghizia, 118; on beans in New South Wales, 47; on Cinchona in Porto Rico, 53; in U.S.A., 13, 60, 96, 103, 127, 289, 290, 291, 292; on fruit trees, 60, 118, 127; on greenhouse plants, 96, 103, 289, 290, 291, 292; measures and experiments against, 13, 53, 96, 103, 118, 289, 290, 291, 292; DDT increasing infestation by, 47, 60, 127.

Tetranychus bimaculatus, sprays against, on rose and phlox in U.S.A., 154, 228, 335; technique of testing acaricides on, 344.

Tetranychus carpini, effect of sprays on natural enemies and, on Carpinus in Switzerland, 282.

Tetranychus cucurbitacearum, sp. n., foodplants of, in Egypt, 411.

Tetranychus pacificus, effects of sprays on infestation of apple by, in Washington, 149, 150; Stethorus picipes predacious on, 150.

Tetranychus schoenei, DDT increasing infestation of apple by, in Virginia, 257.

Tetranychus sexmaculatus, on avocado and Citrus in California, 343; mite predacious on, 343.

Tetranychus telarius, in Australia, 179, 218; in Britain, 284, 288, 360, 361; in Italy, 253; in Switzerland, 282; in U.S.A., 136, 223, 224; on fruit trees, 179, 218, 253; on greenhouse plants, 224, 282, 284, 288, 360, 361; on hops,

288; measures and experiments against, 223, 224, 253, 282, 284, 288, 360, 361; DDT increasing infestation by, 218.

Tetranychus turkestani, Prangos sprays against, on cotton in Kirghizia, 118. Tetranychus urticae (see T. telarius).

Tetranychus willamettei, DDT and pyrethrum not affecting infestation of vines by, in U.S.A., 147.

Tetrapropyl Pyrophosphate, toxicity of, to Aphids, 228.

tetrarhodus, Capitophorus (Pentatrichopus).

Tetrastichus, parasite of Schematiza cordiae in Trinidad, 425, 426.

Tetrastichus giffardi, parasite of Dacus bivittatus in Cameroons, 177.

Tetrastichus giffardianus, parasitising Dacus in Fiji, 266.

Tetrastichus oophagus, sp. n., parasite of Diprion pini in Germany, 239.

Tetrastichus rapo, parasite of Apanteles glomeratus in Europe, 406, 407; bionomics of, 406.

Tetrops praeusta, bionomics of, on pear in Britain, 358.

Tettigonia spp., damaging cereals in Persia, 388. texana, Anomis.

Texas, Anastrepha serpentina in, 221; cotton pests in, 231; Lachnosterna spp. in, 14; pests of pecan in, 151, 152; Gnorimoschema operculella not a pest of field potatoes in, 56; Pseudaphycus mundus found in quarantine in, on cut flowers from Mexico, 316.

Textiles, treatments of, against insects, 224, 323, 324, 395; stained by pentachlorphenol, 224; not damaged by *Thylodrias contractus*, 37.

textor, Hyphantria.

Thamnotettix argentatus (see Orosius).

Thanasimus formicarius, predacious on Myelophilus in Italy, 195.

Thaumetopoea processionea, on Pistacia in Persia, 386.

theae, Agromyza (Melanagromyza);
Apanteles; Fiorinia.

theivora, Gracilaria; Helopeltis.

Theresia claripalpis (parasite of Diatraea), in Br. Guiana, 304; introduction of, into Porto Rico, 54.

Thermobia domestica, in stored groundnuts in Senegal, 44; treatment of textiles with DDT against, 324; heat and humidity reducing toxicity of DDT deposits to, 378; electricallycharged barrier for confining, 342.

Thersilochus, bionomics of, parasitising Lema melanopa in Italy, 37.

Thiocyanates, against Aphids, 218, 327; against Coleoptera, 101, 284, 315, 396; effect of temperature on toxicity of, to Tribolium castaneum, 380; against Jassids, 165, 325, 326, 398; against Mirids, 102, 184, 272, 394; increasing toxicity of oil to Parlatoria oleae, 17; unsatisfactory against Tetranychus, 103; against thrips, 102, 126; in dusts, 102, 126, 184, 218, 314, 327, 394, 396; in sprays, 103, 165, 272, 284, 325, 326, 394, 398; applied from aircraft, 327, 396; and Bordeaux mixture, 165, 325; and other fungicides, 165; and cryolite, 396; and cubé or rotenone, 327, 395; and DDT, 218, 284, 327, 395; vaporised spray of pyrethrum and, in oil, 398; and sulphur or talc, 394; decreasing yield of cotton, 184.

Thiocyanoacetates, tests with, as insecticides, 252-254.

Thiodiphenylamine (see Phenothiazine). Thisoecetrinus pterostichus, damaged by, in Persia, 388.

Thisoecetrus littoralis, crops damaged by, in Persia, 388.

thoracicus, Triaspis (Sigalphus).

Thrips tabaci, in S. Africa, 45; in Australia, 138, 177, 178; occurrence of males of, on leek in Britain, 362; in U.S.A., 12, 102, 230, 234, 338; on beans, 230; differences in infestation of bean varieties by, 138; on cucurbits, 177; on groundnuts, 338; on onion, 12, 102, 234; on tomato, 178, 234; on tobacco, 178; method of reducing loss of tobacco from spotted wilt transmitted by, 45; insecticides against, 12, 102, 230, 234, 338.

Thrips, Banana Rust (see Scirtothrips

signipennis).

Thrips, Gladiolus (see Taeniothrips simplex).

Thrips, Onion or Tobacco (see Thrips tabaci).

Thyanta custator, measures against, causing deformation of peaches in U.S.A., 235, 373.

Thylodrias contractus, habits of, in houses

in Spain, 37.

Thyraeella collaris, introduced parasite of Plutella maculipennis in New Zealand,

Thyreion gelotopoeon, new parasites of, in Argentina, 109.

Thyreophagus entomophagus, in milled cereal products in Queensland, 310.

Thyridopteryx ephemeraeformis, tests of DDT against, on trees and shrubs in U.S.A., 143, 336.

Thysanoptera, transmitting spotted wilt of tobacco, 45; Coccinellid predacious on, 427.

Thysanus, parasite of Leptomastix flavus in Palestine, 410; experimentally parasitising Clausenia purpurea, 410.

tibialis, Chaetocnema.

tigrina, Coenosia.

tigris, Odynerus.

tiliaria, Erannis.

Timber, Coleoptera in, 1, 107, 245, 248, 295, 414-416; depletion of starch from, against Lyctus, 1, 248; lists of Australian species of, susceptible and immune to L. brunneus, 344; termites in, 169, 232, 245, 249, 295, 300, 317, 414-416; list of species of, resistant to termites, 169; treatment of, with preservatives against pests, 1, 107, 169, 232, 245, 248, 295, 300, 317, 414-416.

Timber Treatments, 414.

Timboril, 300.

Timocratica haywardi, new parasite of, in Argentina, 109.

timocraticae, Notamorphota.

Timothy Grass (see *Phleum pratense*).

Tinea columbariella, infesting fleeces in U.S.A., 146.

Tinea pellionella, infesting fleeces in U.S.A., 146; toxicity of thiocyanoacetates to, 253.

Tinea tapetzella (see Trichophaga).

tineiformis, Coniopteryx. tineivorus, Typhlodromus.

Tineola biselliella, infesting fleeces in U.S.A., 146; not attacking cloth treated with Toxaphene, 395; rearing

tingi, Acutaspis.

of, 343.

Tinplate, suitability of, for containers for Toxaphene, 395.

Tiphia parallela, utilisation of, against Clemora smithi in Mauritius, 210.

Tlascala finitella, dust against, on cranberry in Massachusetts, 100.

Tobacco, thrips on, in S. Africa, 45; pests of, in Australia, 69, 178, 219; Aphids on, in Brazil, 7; Myzus persicae on, in Bulgaria, 5; book on pests of, in France, 405; soil pests of, in S. Rhodesia, 137; Lepidoptera on, in Russia, 75; Plusia orichalcea on, in Uganda, 367; pests of, in U.S.A., 87, 183, 371, 372, 373; experiment with form of Macrosiphum solanifolii and, 362; insects and virus diseases of, 4, 5, 7, 45, 75, 219, 422; dense planting of, to reduce loss from spotted wilt, 45.

Tobacco (Stored), pests in, 309; pests controlled by processing of, 85, 86; effect of, on development of Ephestia elutella, 265.

Tobacco (as an insecticide), spraying with, against Halotydeus destructor, 309; sprays of DDT on residues of, 58. Tobacco Budworm (see Heliothis vires-

cens).

Tobacco Thrips (see Thrips tabaci).

Todd Insecticide Fog Applicator, 147.

Togoland, British, Pseudococcus citri in,

Toluene, in insecticide emulsion concen-

trates, 139, 218.
Tolunitriles, low toxicity of, to Cydia

pomonella, 93.

Tomaspis flavilatera, bionomics and control of, on sugar-cane in Br. Guiana, 302-304; methods of making egg

surveys of, 304.

Tomaspis saccharina, on sugar-cane in Trinidad, 302, 303, 318, 319; period incubation of eggs of, 302, 303, 318; egg surveys in relation to tillage against, 318; Metarrhizium anisopliae

infesting, 303, 319.

Tomato, Trypetid on, in S. Africa, 176; pests of, in Australia, 74, 178, 344, 362, 363; pests of, in Bermuda, 49; pests of, in Brazil, 76, 77; pests of, in Britain, 284, 288, 360, 361; grasshoppers damaging, in Chile, Calocoris norvegicus on, in Italy, 383; Halotydeus destructor on, in Zealand, 309; Empoasca lybica on, in Palestine, 176; pests of, in Peru, 133; Trypetid on, in Uganda, 176; pests of, in U.S.A., 10, 94, 144, 186, 234, 373, **394, 397, 398**; experiments with forms of Macrosiphum solanifolii and, 362; insects and virus diseases of, 5, 7, 66, **67, 344, 398, 422;** effects of insecticides on, 19, 94, 234, 288, 360; DDT mixed with stored seed of, against insects, 19.

Tomato Fruitworm (see Heliothis armigera).

Tomato Pinworm (see Keiferia lycoper-

sicella). Tomato Whitefly (see Trialeurodes vapor-

ariorum).

Tonga, miscellaneous pests in, 267; bunchy-top of banana in, 267; parasites of Spodoptera mauritia in, 267.

torquens, Contarinia (see C. nasturtii). torta, Lachnosterna (Phyllophaga).

tortricis, Anachaetopsis.

Tortrix argyrospila, on blueberry in New Jersey, 137.

Tortrix georgiana, on blueberry in New Jersey, 137.

Tortrix lacheana, dormant sprays against, in orchards in Holland, 276.

Tortrix packardiana, parasites of, on spruce in Quebec, 321.

Tortrix podana, dormant sprays against, in orchards in Holland, 276.

Tortrix reticulana, dormant sprays against, in orchards in Holland, 276.

Tortrix rosaceana, on blueberry in New Jersey, 137.

Tortrix striana, on spruce in Quebec, 321. Torymus anastativorus, sp. n., parasite of Anastatus disparis in Jugoslavia, 117.

Toumeyella liriodendri, on walnut and magnolia in California, 189, 400.

Toxaphene, against grasshoppers, 226, 227, 237; against orchard and vegetable pests, 225; cloth treated with, against clothes moths and carpet beetles, 395, 396; in dusts, 224, 225, 237; in sprays, 224, 225, 226, 237; other insecticides and fungicides compatible with, 224, 225; properties of, 224, 395; toxicity of, to mammals, 396; effect of, on plants, 224, 225; tests relating to storage of, 395.

Toxoptera aurantii, on orange in W. Australia, 74; on Cassia and cacao in Fiji, 413; on Cinchona in Porto Rico. 53; on tea in Travancore, 251; tests of

DDT against, 74, 251.

Toxoptera graminum, bionomics and control of, on cereals and grasses in Uruguay, 49, 50.

trachynotus, Meteorus.

Trachyrhachys kiowa, on Parthenium argentatum in Mexico, 97.

Trachyte, test with, against Calandra oryzae, 210.

transvaalensis, Apanteles.

transversalis, Coccinella repanda.

trapezina, Cosmia (Calymnia).

Traps, for Aphids, 195; for Dacus oleae, 201.

Travancore, pests of tea in, 250, 251.

Triaenodes bicolor, DDT toxic to, in ricefields in Italy, 347.

trialbamaculella, Aroga (Gelechia).

Trialeurodes vaporariorum (on greenhouse plants), in Britain, 361; in New Hampshire, 234; in Switzerland, 282; in New in Tasmania, 70; use of Encarsia formosa against, 70; other measures and experiments against, 234, 282, 361.

Triamylamine, not increasing toxicity of oil to Parlatoria oleae, 17.

Triaspis, 31; introduction of, into California from S. America against Listroderes obliquus, 97.

Triaspis gibberosus, parasite of Bruchus pisorum in Hungary, 62.

Triaspis thoracicus, parasite of Bruchus pisorum in Hungary, 62.

Tribolium, type of bag affecting infestation of groundnuts by, 25: treatment of sacks with DDT against, 324; revision of, 261.

Tribolium castaneum, in E. Africa, 367, 426; in S. Africa, 2; in Australia, 68, 69, 182, 309, 310; parasite of, in Senegal, 44; in U.S.A., 85; in stored cereals and flour, 2, 19, 20, 28, 68, 69, 85, 182, 309, 310, 367; in stored groundnuts, 44, 426; in other stored products, 68, 309, 367; fumigants against, 2, 20, 85, 182; other measures and experiments against, 19, 28, 139, 235, 376; effect of temperature on toxicity of insecticides to, 380.

Tribolium confusum, distribution of, in Australia, 68, 69; parasite of, in stored groundnuts in Senegal, 44; in stored cereals and flour, 14, 15, 19, 68, 348; susceptibility of soybean products to, 348; tests with high frequency electric fields against, 16; other experiments against, 14, 19, 20, 235, 236.

Tribolium indicum, new forms and distribution of, in Africa, 261.

Tribolium madens, tests with benzene hexachloride against, 235.

Trichloracetonitrile, tests with, against pests of stored grain, 20, 38.

2, 2, 3-Trichlorbutyronitrile, tests with, against insects in stored wheat, 20; in mixture with other fumigants, 20.

1, 1, 2-Trichlorethane, tests with, against insects in stored maize, 85.

Trichlorethylene, tests with, against insects in stored grain, 85, 182; ethylene dichloride, 182.

1, 1, 1-Trichlor-2, 2 bis (parabromophenyl) ethane (see Bromo-DDT).

1, 1, 1-Trichlor-2, 2 bis (parachlorphenyl)ethane (see DDT).

1, 1, 1-Trichlor-2, 2 bis (parafluorophenyl)ethane (see Fluoro-DDT).

1, 1, 1-Trichlor-2, 2-bis (paramethoxyphenyl) ethane (see Methoxy-DDT).

Trichlorphenol, insecticidal activity of,

2, 2, 3-Trichlorpropionitrile, tests with, against insects in stored wheat, 20. trichodactyla, Hylemyia (Chortophila).

Trichogramma, introduction of, against Diatraea saccharalis in Porto Rico, 54.

Trichogramma evanescens, parasite Cassida nebulosa in Russia, Lathromeris bruchocida confused with, 242.

Trichogramma minutum, 54; parasite of Diatraea in Br. Guiana, 304; effect of sprays on, parasitising Cydia pomonella in W. Virginia, 219; selection of strains of, 200.

Trichogrammatoidea nana, parasite of Eublemma amabilis in Bihar, 120.

Trichomalus laevinucha, bionomics of, parasitising Megastigmus kuntzei in Poland, 268.

Trichophaga tapetzella, toxicity of thiocyanoacetates to, 253.

Trichoplusia ni, on crucifers in U.S.A., 159, 234, 421; dusts against, 234, 421.

Trichopsidea oestracea, parasite of Chortoicetes terminifera in New South Wales, 344; survival of drought conditions by, 344.

Trichoptilus wahlbergi, on sweet potato in Mauritius, 210.

tridentata, Eutetrapha (Saperda). Triethanolamine Dodecyl Sulphate, DDT spray formula containing, 131.

Trifolium (see Clover).

Trifolium alexandrinum, Hypera brunneipennis on, in Arizona, 55; new mite on, in Egypt, 411.

Trifolium subterraneum, pests of, in Australia, 73, 244.

trifurcata, Cerotoma.

Trigonogastra, parasite of Agromyza theae in Ceylon, 368.

trilineata, Lema.

trilineatus, Ptychodes. trilobitiformis, Pseudaonidia.

Trimerotropis ochraceipennis, bionomics

of, in Chile, 109.

Trinidad, investigations on insects defoliating Cordia macrostachya in, 424-426; miscellaneous pests in, 126, 221, 319; Tomaspis saccharina on sugar-cane in, 302, 303, 318, 319; beneficial insect in, 126: Metarrhizium anisopliae in, and its importation into Br. Guiana, 303, 319.

Trionymus sacchari, on sugar-cane in Br. Guiana, 304; experiment with Leis conformis and, in Egypt, 411; parasite of, in Porto Rico, 316.

Trioxys cirsii (aceris), utilisation of, against Myzocallis annulata in Tas-

mania, 70; parasite of, 70.

Triplochiton scleroxylon, new Coccids on,

in Gold Coast, 110. Tripolitania (see Libya).

Tripolite, as diluent for dusts, 329.

tristigmus, Euschistus.

tristis, Anasa.

tritoma, Loxotropa. Triton, 337.

Triton 720, 146.

Triton B-1956, 335, 339. Triton X-100, 90, 165, 340.

Tritox, 38.

Triumfetta, new Mirid on, in Uganda, 26. Trogosita mauritanica (see Tenebroides). Trybliographa diaphana, parasite of

Hylemyia spp. in England, 174. tryoni, Dacus (Strumeta) ferrugineus.

Trypodendron lineatum, removal of bark from felled spruce against, in Switzerland, 408.

Trypterygium forrestii, insecticidal action and constituents of, 298, 299.

tuberis, Epitrix.

Tulip, mite on bulbs of, in U.S.A., 83. Tulip-tree Scale (see *Toumeyella liriodendri*).

tulipae, Aceria.

Tung (see Aleurites).

Tunisia, Schistocerca gregaria in, 206, 207.

turanicus, Calliptamus; Chrotogonus. turcomana, Ramburiella.

Turf, insects damaging, in Australia, 73, 307; cutworms damaging, in Mauritius, 210; *Lachnosterna* damaging, in Ontario, 164; effect of benzene hexachloride on, 280.

turkestani, Tetranychus (Eotetranychus).

Turkey, tests of DDT against insects in, 352; text-book on economic ento-

mology in, 3.

Turnip, Phyllotreta transmitting new virus disease of, in Britain, 377; Listroderes obliquus on, in California, 187; Helophorus on, in Holland, 190; Calocoris norvegicus on, in Italy, 383; DDT residues on, 135; not injured by Velsicol AR-60, 421.

Turpentine, against Coleoptera in timber, 1, 248, 414, 416; and kerosene, 248,

414, 416.

Tussore Silk Worm (see Antheraea pernyi). Tutsan (see Hypericum androsaemum).

Tychius, not parasitised by Lathromeris bruchocida, 242.

Tychius quinquepunctatus, on peas in Hungary, 63.

Tydeus womersleyi, on Albizzia in Ceylon, 369.

Tylococcus westwoodi, sp. n., on cacao in Gold Coast, 110.

Tyndarichus kuriri, sp. n., parasite of Anastatus disparis in Jugoslavia, 117.

Typhlocyba froggatti (in orchards in Australia), biological control of, 70; sprays against, 74, 246.

Typhlocyba rosae (on rose), in Switzerland, 282; in U.S.A., 335; sprays against, 282, 335.

Typhlodromus tineivorus, predacious on Ephestia elutella in Britain, 212. typicus, Eriophyes quadrisetus.

typographus, Ips.

Tyroglyphus farinae, nature of damage to stored wheat by, in Britain, 417; infesting cheese in New Zealand, 416.

Tyroglyphus nadinus, sp. n., bionomics of, in Italy, 114.

Tyrolichus casei, infesting cheese in New Zealand, 244, 416; measures against, 244.

Tyrophagus, on mushrooms in New South Wales, 365; infesting cheeses in New Zealand and Queensland, 244, 363, 416; measures and experiments against, 244, 363, 365.

Tyrophagus longior, 244, 416. Tyrophagus putrescentiae, 363, 365.

U

Udeopsylla, killed by benzene hexachloride dust in Oklahoma, **100**.

Uganda, cotton pests in, 26, 27, 366, 367; Gonipterus scutellatus on Eucalyptus in, 106, 367; pests of field and stored groundnuts in, 426; miscellaneous pests in, 26, 176, 367; Schistocerca gregaria in, 123, 124, 207, 366; beneficial insects in, 27, 109, 367, 426; failure to establish Plaesius javanus in, 367; new hyperparasite of Pseudococcus in, 109.

Ulex europaeus, establishment of Apion ulicis against, in New Zealand, 244.

ulicis, Apion.

ulmi, Lepidosaphes.

ulmi, auct., Metatetranychus (Oligonychus) (see Paratetranychus pilosus).

ulmifolii, Myzocallis.

Unaspis citri, spray against, on Citrus in New South Wales, 181.

Unaspis euonymi, on Euonymus in California, 400.

undulata, Phyllotreta.

unicolor, Bruchus (Bruchidius).

Union of Soviet Socialist Republics, miscellaneous pests in, 118, 241, 242; Lepidoptera and virus disease of tobacco in, 75; pests of vines in, 3, 4; beneficial insects in, 241, 242; potentialities of biological control in, 241; digest of plant-quarantine restrictions in, 105.

unipuncta, Cirphis (Sideridis).

unispina, Gryllotalpa.

United Provinces, pests of mango in, 110, 123; miscellaneous pests in, 122, 139.

United States of America, cereal pests in, 12, 13, 14, 22, 81; pests of Citrus in, 13, 14, 79, 83, 89, 145, 155, 160, 181, 187, 188, 189, 221, 343; cotton pests in, 12, 13, 14, 156, 183, 184, 231, 370, 371, 401; pests of forest and shade trees in, 14, 33, 36, 87, 92, 143, 144, 158, 220, 234, 243, 254, 301, 320, 333, 334, 336, 388; orchard pests in, 12, 36, 81, 83, 92, 334, 335, 336, 410; pests of ornamental plants in, 84, 190, 335; pests of stored products in, 12, 14, 85, 86, 146, 282; sugar-cane pests in, 147, 316; pests of sunflower and allied plants in, 83, 84; tests of insecticides against miscellaneous pests in, 12-14, 81, 328, 335; review of injurious Eriophyids of, 82; Popillia japonica in, 36, 101, 209, 234, 333, 335, 336, 337, 339; identity of introduced species of Pseudocneorrhinus in, 301; book on termites in, 317; eradication of certain pests in, 219; beneficial insects in, 28, 29, 84, 126, 178, 410; pests intercepted in quarantine in, 233, 240, 315, 316, 400, 401; plant pest legislation in, 71, 105, 184, 209; list of popular names of insects in, 32; directory of pest control in, 347; review of patents relating to pest control in, 240. (See also under the various States.)

univittatus, Tarsostenus.

ununguis, Paratetranychus.

urbana, Ctenolepisma.

Urbanus proteus (Bean Leaf-roller), toxicity of methoxy-DDT to, in U.S.A., 14. urbaurea, Aristotelia.

Urena lobata, Tingid on, in Florida, 24; Earias spp. on, in Punjab, 312.

Urentius sentis, experiments with DDT against, in India, 123.

urichi, Liothrips.

urticae, Tetranychus (see T. telarius).

Uruguay, *Toxoptera graminum* on cereals and grasses in, 49; natural enemies of Aphids in, 50; parasites of *Listroderes obliquus* introduced into California from, 97; summary of plant quarantines in, 105.

Uscana semifumipennis, parasite of Bruchid eggs in Hungary, 62.

ustulatus, Agriotes.

Utah, lizards and birds destroying ants in, 24; miscellaneous pests in, 10-12, 185, 388; western celery mosaic in, 388; hees pollinating lucerne in, 10.

V

vaccinii, Mineola; Ophiola.

Vallota purpurea, Agasphaerops nigra not

feeding on, 142.

Vanessa cardui, bionomics and migrations of, in Eritrea, 382, 383; spray against, 383.

vaporariorum, Trialeurodes.

varia, Desmeocraera.

variabilis, Erythroneura; Hypera (Phyto-

nomus); Lixophaga. variegana, Argyroploce.

variipennis, Phyllotreta.

varinervis, Antholcus. variolarius, Euschistůs.

variolosum, Asterolecanium.

varivestis, Epilachna.

Vasates cornutus, on peach in N. America,

83.

Vasates fockeui, on plum in Canada and U.S.A., 83; dusts against, on greenhouse peach and plum in Holland, 239

Vasates malivagrans (see V. schlechtendali).

Vasates schlechtendali (Apple Rust Mite), described from California as V. malivagrans, sp. n., 80; synonymy of, 83.

Vaseline, use of, to protect timber against *Lyctus*, 248.

Vatsol-OT (see Dioctyl Sodium Sulphosuccinate).

vayssierei, Embia.

Vegetable Marrow, Halotydeus destructor on, in New Zealand, 309.

Vegetable Weevil (see Listroderes obliquus).

vellicatus, Exenterus.

Velsicol 1068 (see Chlordane).

Velsicol AR-60 (mainly methylnaphthalenes), increasing effectiveness of cubé extractives and dusts, 17, 421; as a solvent for DDT, 89, 95, 165, 325, 326, 333, 421; increasing effectiveness of DDT dusts, 420, 421; as solvent for di(4-chlorphenoxy)methane, 155; plants injured by, 421.

velutinana, Eulia.

Velvet-bean Caterpillar (see Anticarsia

gemmatalis).

Venezuela, Sacadodes pyralis on cotton in, 126; digests of plant-quarantine restrictions in, 105.

venilialis, Marasmia.

Ventox, 62.

ventricosus, Pediculoides (Pediculus).

Venturia inaequalis (Apple Scab), sprays against, in Britain, 131, 284, 359; influence of sprays against, on Arthropods in Nova Scotia, 271.

Veralin 3, 274.

Verbascum thapsus, Euschistus on, in Virginia, 334, 373.

Verbesina encelioides, Homoeosoma electellum on, in Missouri, 84.

verditer, Amblymerus. vernata, Paleacrita.

verrucivorus, Decticus.

versicolor, Agonoscelis (see A. pubescens). versicolora, Plagiodera.

versutus, Adoretus.

vesicularis, Eupelmella.

vestitus, Anthonomus; Hylesinus (Chaetoptelius).

Vetches, Cydia nigricana on, in Britain, favouring Euschistus spp. in peach orchards in Virginia, 373.

Viburnum, Coccids on, in California, 400. Vicia (see Vetches).

Vicia faba (see Broad Beans).

vicinus, Homorocoryphus; Scapteriscus.

Victoria, orchard pests in, 178, 179; pests of stored wheat in, 181, 182, 217, 364; failure of biological control of Cydia molesta in, 178; biological control of Hypericum perforatum in,

vigintioctopunctata, Epilachna. Vigna unguiculata (see Cowpeas). villosa, Lagria.

Vine, Grape, pests of, in greenhouses in Belgium and Britain, 278, 288, 292; grasshoppers damaging, in Chile, 109; new mites on, in Egypt, 411, 412; Phalaenoides glycinae on, in South Wales, 108; in Palestine, 176; leafhoppers on, Coccids on, in Punjab, 123; weevils on, in Spain, 198; pests of, in Switzerland, 280, 296; pests of, in Turkmenistan, 3, 4; pests of, in U.S.A., 83, 127, 147, 185, 255, 325, 397, 398; experiment with Cydia molesta and, 239; not attacked by Quadraspidiotus perniciosus, 64; insecticides and injury to, 224, 253, 284, 288; question of DDT residues on fruit of, 81, 135, 147.

Vinegar Flies (see Drosophila).

vinitor, Nysius.

vinnula, Leptopharsa machalana.

Violet, Tetranychus telarius on, in greenhouses in Britain, 288.

virens, Lygus. virescens, Heliothis.

virgata, Ferrisia (Pseudococcus).

Virginia, pests of leguminous crops in, 12, 13, 14, 229, 230, 326, 332, 337, 338, 374, 394; miscellaneous pests in, 96, 331-333, 374; orchard pests in, 256, 257, 334, 372, 373, 374; new parasites and biological control of mealybugs in, 315, 316.

Virginia, West, Cydia pomonella and its natural enemies on apple in, 219, 220, 258.

virginiensis, Anisota. viridifasciata, Chortophaga. viridis, Coccus; Smynthurus. viridissima, Tettigonia. virilis, Blondelia (Lydella).

Virus Disease, of Gilpinia hercyniae, 208, 319.

Virus Diseases (in plants*), relation of Aphids to, 4, 5, 7, 46, 79, 93, 209, 267, 313, 355, 365, 376, 388, 413, 422, 423, 424, 426; relation of Aleurodids to, 299, 300; Coccids transmitting, 110; relation of Jassids to, 77, 186, 219, 231, 344, 350, 355, 398; Piesma quadratum transmitting, 355; Thysanoptera transmitting, 45; Phyllotreta transmitting, 377; relation of Plusia gamma to, 75; mechanical transmission of, 5, 7, 79, 313, 422; book on, 312.

visci, Diaspis.

Vitamin, requirements of, by Ephestia elutella, 265, 266.

viteana, Polychrosis.

Vitex, Trypetid on, in Uganda, 176.

vitiensis, Syrphus corollae. vitifoliae, Phylloxera.

Calepitrimerus (Epitrimerus); vitis, Eriophyes.

vitripennis, Mormoniella.

vittata, Diabrotica (see D. melanocephala);

Epicauta; Phyllotreta. vittiger, Dichroplus.

vollenweiderei, Atta (Neoatta).

vomitoria, Calliphora. vorax, Anthrenus.

vosseleri, Lygus.

Vrilletta, in baled shrubs of Parthenium argentatum in Mexico, 97.

vulgaris, Asaphes; Cremnops (Bracon). vulpinus, Dermestes (see D. maculatus). vuteria, Sesamia.

W

wachtli, Megastigmus. wacri, Newsteadia. wahlbergia, Trichoptilus.

Wallflower, Haplomyza imitans on, in New South Wales, 73.

^{*} These references are also indexed under the individual plants infected.

Walnut, Cydia pomonella on, in Italy, 194; Chromaphis juglandis on, in Kirghizia, 118; pests of, in U.S.A., 80, 155, 188, 189, 399.

Walnut Husk Fly (see Rhagoletis suavis completa).

Walnut-shell Flour, as carrier for dusts, 9, 154, 330.

Walshia amorphella, parasites of, on Amorpha fruticosa in Oklahoma, 9.

Washington, *Hylemyia brassicae* on cabbage in, 222; orchard pests in, 19, 98, 149, 150, 185; beneficial insects in, 98, 150.

Washingtonia, Litoprosopus coachella on, in California and Mexico, 400.

Wasps, attacking grapes in Turkmenistan, 4.

Water, Hot, use of, against Lepidoptera, 186; treatment of strawberry with, against *Tarsonemus pallidus*, 353, 359.

Water Hyacinth (see Eichhornia).

Watermelon, not attacked by Melittia cucurbitae in Massachusetts, 101; Diaphania on, in Peru, 134; effect of insecticides on, 134; DDT mixed with stored seed of, against insects, 19. wauaria, Itame.

Webworm, Fall (see Hyphantria cunea).

West Indies, *Coelosternus* on cassava in, 6; book on pests of food-crops in, 71. (See also under the individual Islands.)

Western Grape Leaf Skeletoniser (see Harrisina brillians).

westwoodi, Tylococcus.

Wetsit, as an emulsifier, 69; composition of, 69.

Wheat, effect of wireworms on yield of, in Britain, 381; Cephus cinctus on, in Canada, 390; pests of, in Italy, 37, 114; Dasus simplex on, in Kenya, 404; Anisoplia spp. on, in Persia, 387; pests of, in Spain, 202; Aphid on, in Uruguay, 49; pests of, in U.S.A., 14, 91, 128, 136; time of sowing of, in relation to Mayetiola destructor in Ohio, 136; probably not injured by benzene hexachloride, 413.

Wheat (Stored), pests in, and their control, 3, 12, 14, 19, 20, 38, 39, 68, 134, 139, 182, 210-214, 217, 264, 282, 364, 365, 417, 418; effect of, on development of *Ephestia elutella*, 265; book on safe storage of, 405; methods and equipment for sampling in silos, 419; studies on sorption of fumigants by, 39, 223.

Wheat Stem Sawfly (see *Cephus cinctus*). White Spirit, as solvent or penetrant in mixtures containing pentachlorphenol

against borers in timber, 1, 107.
White-fringed Beetles (see *Pantomorus*

White-fringed Beetles (see Pantomorus (Graphognathus)).

Whiteflies (Aleurodids), transmitting infectious chlorosis of Malvaceae, 299, 300; natural enemies of, 70, 155; lists of parasites and predators of, 270.

Whitefly, Greenhouse or Tomato (see *Trialeurodes vaporariorum*).

willamettei, Tetranychus.

Willow, Cosmia trapezina on, in Belgium, 277; Prionus coriarius on, in Britain, 215; Plagiodera versicolora on, in Canada, 104; Chrysomphalus dictyospermi on, in Persia, 386; Hyponomeuta on, in Turkey, 352; pests of, in U.S.A., 14, 143, 188, 234, 254.

Wind, effect of, on migration of Schistocerca gregaria, 205; relation of, to dispersion of other insects, 8, 64, 221, 397.

Winter Moth (see Operophtera brumata). Wireworm, False (see Psammodes similis).

Wireworms, populations of, in relation to crop yields, 281, 381, 382; influence of cover crops on injury to tobacco by, 183; measures against, 187, 281; susceptibility of, to stomach poisons, 287.

Wisconsin, Macrosiphum onobrychis on peas in, 12, 153, 229; pests of pine in, 220; Cercopids on miscellaneous crops in, 160.

Wistaria, Rhynchites coeruleus on, in Britain, 358.

womersleyi, Tydeus.

Womersleyia minuta, gen. et sp. n., parasite of Orthoptera in Maldive Is., 427.

Wood Ash (see Ash).

Woodlice, tests of DDT against, 159, 292.

Woollen Cloth (see Textiles).

Woolly Apple Aphis (see Eriosoma lanigerum).

Wyoming, Paratrioza cockerelli in, 144.

X

Xanthogramma grandicorne, predacious on Anuraphis persicae-niger in New South Wales, 247.

Xanthogramma scutellare, predacious on Aphids in Fiji, 414.

xanthomelaena, Disonycha; Galerucella (see G. luteola).

Xanthone, in sprays against orchard pests, 58, 98, 149, 150; not toxic to bees, 222; and DDT, 58, 98, 150; and oil emulsion, 98, 149; and injury to apple, 150.

Xanthopimpla spp., parasites of Proceras sacchariphagus in Mauritius, 210;

life-cycle of, 210.

Xorides albitarsus, parasite of Prionus coriarius in Britain, 215.

Xyleborus, susceptibility of Inga spp. to, in Porto Rico, 300.

Xyleborus dispar, on apple and cherry

in Britain, 358.

Xyleborus fornicatus, on tea in Ceylon, 251, 368; factors affecting damage by, 368; food-plants, distribution and status of subspecies of, 251.

Xyleborus fornicatus fornicatior, **251.**

Xyleborus xylographus (saxeseni), cherry in Britain, 358.

Xylene, as solvent for DDT, 34, 333, 336, 342; emulsified solutions of DDT in, 144, 158, 218, 220, 267, 340; solutions or emulsified solutions of other organic insecticides in, 228, 237, 333, 335, 340, 342.

xylographus, Xyleborus.

Xyloperthodes, in dried cassava roots in

Uganda, 367.

Xylophanes pluto, measures against, on Cinchona in Porto Rico, 53.

Xyloterus (see *Trypodendron*).

Y

Yam Bean (see Pachyrhizus erosus). Yeast, in bait for Cydia pomonella, 132; in food for clothes moths and carpet beetles, 343.

Yellow-flowered Sweet Clover (see

Melilotus officinalis).

Yellow-striped Armyworm (see Prodenia ornithogalli).

Yezabura (see Anuraphis).

Yezabura malifoliae, auct. (see Anuraphis roseus).

ypsilon, Agrotis (Euxoa). yuccae, Puto (Ceroputo). yuccarum, Targionia.

Z

Zanzibar, host fruits and parasites of Dacus bivittatus cucumarius in, 176, 177.

Zeiraphera ratzeburgiana, on spruce in Quebec, 321.

Zenillia caesar, parasite of Harmologa fumiferana in Canada, 207, 320.

Zicrona coerulea, predacious on Leptinotarsa decemlineata in Spain, 196.

Zinc Arsenate, in sprays against Prays oleellus on olive, 141.

Zinc Chloride, treatment of timber with, against Coleoptera and termites, 169, 245, 414, 415, 416.

Zinc Dimethyldithiocarbamate, treatment of wood with, against Kalotermes brevis, 233.

Zinc Fluoarsenate, in sprays against Cydia pomonella on apple, 180, 247; and injury to foliage, 180, 247.

Zinc Fluoborate, treatment of wood with. against Kalotermes brevis, 232.

Zinc Fluoride, treatment of wood with, against Kalotermes brevis, 232.

Zinc Fluosilicate, treatment of wood with, against Kalotermes brevis, 232.

Zinc Naphthenate, treatment of wood with, against Anobium and Kalotermes brouni, 245, 414, 415.

Zinc Oxide, reducing effect on plants of calcium arsenate in soil, 371.

Zinc Pentachlorphenate, treatment of wood with, against Kalotermes brevis, 233.

Zinc Sulphate, doubtful value of, against Lyctus in timber, 1; in DDT spray formula, 165; effects of, with and without lime in lead-arsenate sprays, 151, 152, 180, 373.

zinckenella, Etiella.

zizyphus, Parlatoria.

Zonocerus elegans. on cassava in Nyasaland, 44.

NOTICES.

Secretaries of Societies and Editors of Journals willing to exchange their publications with those of the Institute are requested to communicate with the Director. Authors of papers on economic entomology, whether published in entomological journals or not, are invited to send reprints to the Director for notice in the Review.

The Annual Subscription, in advance, to Volume 38 of the Review, Series A (Agricultural), is 40s. post free; Series B (Medical and Veterinary), 20s. post free. Prices of Back Volumes on application.

Orders and Subscriptions should be sent to the Director, Commonwealth Institute of Entomology, 41, Queen's Gate, London, S.W.7, or through a bookseller.

